



Revista de Biología Marina y Oceanografía

ISSN: 0717-3326

revbiolmar@gmail.com

Universidad de Valparaíso

Chile

Sánchez-Saavedra, M. del Pilar; Licea-Navarro, Alexei; Bernáldez-Sarabia, Johanna  
Evaluation of the antibacterial activity of different species of phytoplankton  
Revista de Biología Marina y Oceanografía, vol. 45, núm. 3, diciembre, 2010, pp. 531-536  
Universidad de Valparaíso  
Viña del Mar, Chile

Available in: <http://www.redalyc.org/articulo.oa?id=47919217019>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System  
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal  
Non-profit academic project, developed under the open access initiative

# Evaluation of the antibacterial activity of different species of phytoplankton

Evaluación de la actividad antibacteriana de diferentes especies de fitoplancton

M. del Pilar Sánchez-Saavedra<sup>1</sup>, Alexei Licea-Navarro<sup>2</sup> and Johanna Bernáldez-Sarabia<sup>1,2</sup>

<sup>1</sup>Departamento de Acuicultura, Centro de Investigación Científica y de Educación Superior de Ensenada (CICESE), Carretera Ensenada-Tijuana 3918, Zona Playitas, Ensenada Baja California, C.P. 22860, México. psanchez@cicese.mx

<sup>2</sup>Departamento de Biotecnología Marina, Centro de Investigación Científica y de Educación Superior de Ensenada (CICESE), Carretera Ensenada-Tijuana 3918, Zona Playitas, Ensenada Baja California, C.P. 22860, México

**Abstract.**— The antibiotic activity of organic extracts from cultures of six species of phytoplankton, *Synechococcus elongatus*, *Synechocystis* sp., *Amphiprora paludosa*, *Porphyridium cruentum*, *Chaetoceros muelleri* and *Dunaliella tertiolecta* was examined. The bacteria used were Gram(-) strains, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Klebsiella pneumoniae*, and the Gram(+) strains were *Bacillus subtilis*, *Staphylococcus aureus*, and *Micrococcus luteus*. All extracts inhibited the growth of *B. subtilis* and effected the generation of inhibition halos, ranging in diameter from 7.06 to 15.23 mm, for several bacteria species. Wide-spectrum antibacterial activity was observed in extracts from *S. elongatus*, *Synechocystis* sp., *A. paludosa*, *P. cruentum*, and *C. muelleri*. Thus, this five phytoplankton species can be used to control bacteria.

**Key words:** Antibiotics, microalgae cultures, cyanobacteriae

## INTRODUCTION

Marine planktonic algae are potentially new sources of specific antibacterial compounds (Patil *et al.* 2009). Diatoms are the major groups of phytoplanktonic class within algae, in which most organic compounds found in them with antibacterial activity have been pigments, lipids, and carbohydrates (Shimizu 1993), cyanobacteria (blue-green algae) have unique, biologically active secondary metabolites due to their ecological and morphological diversity (Kreitlow *et al.* 1999). Cyanobacteria proved to be an excellent source of natural metabolites, possessing antibacterial, antifungal, and cytotoxic-cytostatic activity (Patil *et al.* 2009). The chemical structures of these cyanobacterial metabolites have been determined, including polyketides, amides, alkaloids, lactones, peptides, and lipopeptides (Singh *et al.* 1999).

The use of antimicrobial agents in aquaculture can increase the prevalence of bacteria that are resistant to the organisms that are maintained in the culture; such bacteria could be transmitted and cause infections in humans on consumption of cultured organisms (WHO 2006). In aquaculture, the indiscriminate use of antibiotics influences the rates of diseases and resistance. Several international organizations have restricted the use of certain antibiotics in organisms that are produced for human consumption (FDA 2009).

The antagonism between microbes is a naturally occurring phenomenon through which pathogens are killed or decline in number in the aquaculture environment (Maeda *et al.* 1997). The identification and development of antibiotics are needed continuously due to the increase in resistant pathogens, the evolution of emergent diseases, the existence of naturally resistant bacteria, and the observed toxicity of current medicinal compounds (Demain 1999). In aquaculture, certain bacterial populations govern this equilibrium through their interactions with various pathogenic agents. An understanding of the specific interactions between bacteria and microalgae will allow us to optimize productive systems in aquaculture (Riquelme & Avendaño 2003). Thus, our aim was to examine the antibacterial properties of several strains of marine phytoplankton for their potential use in aquaculture.

## MATERIAL AND METHODS

Marine phytoplankton strains *Amphiprora paludosa* var. *hyalina*, *Synechococcus elongatus*, and *Synechocystis* sp. were obtained from 'Laboratorio de Microalgas, Centro de Investigación Científica y de Educación Superior de Ensenada (CICESE), Ensenada, México'. *Dunaliella tertiolecta* and *Chaetoceros muelleri* strains were obtained from the Oyster Center of Sonora (COES), and

*Porphyridium cruentum* was acquired from the Institute of Applied Microbiology at the University of Tokyo, Japan.

Cultures were maintained in nonaxenic, monospecies batch cultures in triplicate. The bacterial counts of all phytoplankton strains were lower than  $1 \times 10^6$  cells mL<sup>-1</sup>. Cultures were grown in batches in 2.8-L Fernbach flasks that contained 2 L of medium (Guillard 1975), where in silicate and vitamin concentrations were doubled for diatoms and cyanobacteria, respectively. Cultures were maintained at  $20 \pm 2^\circ\text{C}$  with continuous light at  $75 \mu\text{E m}^{-2}\text{s}^{-1}$  and shaken manually every 12 h, no oxygen was supplied by aeration.

To generate the growth curves, optical density (OD) was measured daily at 550 nm on a Hach DR/4000 UV-VIS spectrophotometer. Cell concentrations were determined using a hemacytometer, and size was measured by light microscopy. The cell concentrations were used to obtain the exponential growth rate ( $\mu$ ) and mean doubling time (G) (Fogg and Thake 1987). The number of colony-forming units (CFU) was determined in triplicate sets of each cultured microalgal strain using the method of Colwell *et al.* (1975).

Bacterial strains were obtained from the bacterial culture collection of the Faculty of Chemistry Science and Engineering at 'Universidad Autónoma de Baja California' (UABC), Tijuana, México. The Gram(-) strains were *Escherichia coli*, *Pseudomonas aeruginosa*, and *Klebsiella pneumoniae*, and the Gram(+) strains were *Bacillus subtilis*, *Staphylococcus aureus*, and *Micrococcus luteus*. Bacteria were precultured at  $37^\circ\text{C}$  for 12 h in Petri dishes in 10 mL Luria-Bertani agar (LB agar) medium (Maniatis *et al.* 1982).

The microalgae and cyanobacteria cultures were harvested on day 12 during the stationary phase, and the cells were concentrated by centrifugation at 3500 g for 15 min at  $20^\circ\text{C}$ . The total cell biomass of each cultured strain was lyophilized, and the dry weight value was obtained gravimetrically (Table 1).

The cells were disrupted with an ultrasonic processor at 80 MHz for 2 min with 3 3-second intervals and extracted twice with 10 mL hexane; during the extractions, flasks were shaken at 150 rpm for 2 h in the dark at  $20^\circ\text{C}$  (CLSI 2006). Pellets from the hexane extraction were reextracted with dichloromethane and methanol, as described above. The solvents were evaporated under a nitrogen stream. The dry microalgal extracts were dissolved in 1 mL dimethyl sulfoxide (DMSO).

The antimicrobial tests were performed in vitro on LB agar plates by diffusion method (Collins *et al.* 1989). The plates were inoculated with a standardized suspension of each bacterial strain (1.0 OD, measured at 600 nm as the McFarland equivalent concentration of  $10^8$  CFU mL<sup>-1</sup>) and maintained at  $37^\circ\text{C}$  (CLSI 2006).

The inhibition of bacterial growth was assessed in triplicate using sterilized 6-mm-diameter paper disks (S- & S Filter-Paper-Disks) that contained 30  $\mu\text{L}$  of algal extract in DMSO. Carbenicillin ( $500 \mu\text{g mL}^{-1}$ ), a broad-spectrum antibiotic, was used as a positive control, and seawater and DMSO were used as negative controls (Al-Saimary *et al.* 2007). The inhibition zone was defined as the diameters of the area in which no bacterial growth was detected (CLSI 2006). Measurements were recorded after 24 h of incubation at  $37^\circ\text{C}$ . Pre-assays were run to

**Table 1.** Mean cell size, cell density of the inocula and at harvest, exponential growth rate ( $\mu$ ), mean doubling time (G), dry weight, and colony-forming units (CFU) of marine microalgae and cyanobacterial strains maintained in batch cultures / Tamaño promedio de la célula, densidad de células en el inóculo y en la cosecha, tasa de crecimiento exponencial ( $\mu$ ), tiempo de duplicación (G), peso seco y unidades formadoras de colonias (CFU) de microalgas y cianobacterias marinas mantenidas en cultivos estáticos

Strain	Cell size $\mu\text{m}$	Cell density (inoculated) cells mL <sup>-1</sup> $10^6$	Cell density (harvested) cells mL <sup>-1</sup> $10^6$	$\mu$ divisions day <sup>-1</sup>	G	Dry weight mg mL <sup>-1</sup>	CFU CFU x $10^3$ mL <sup>-1</sup>
<i>Synechococcus elongatus</i>	1 x 2	0.06	24.3	0.7	1.4	1.1	n.d.
<i>Synechocystis</i> sp.	1.3 x 2	0.06	28.6	0.7	1.4	0.7	n.d.
<i>Amphiprora paludosa</i>	5.9 x 2.7	0.02	4.6	0.6	1.6	0.4	0.6
<i>Dunaliella tertiolecta</i>	8-11 x 14-16	0.02	1.8	0.5	2.0	0.5	3.0
<i>Chaetoceros muelleri</i>	8-12 x 7-10	0.02	2.6	0.5	2.0	0.9	2.6
<i>Porphyridium cruentum</i>	5 x 8	0.02	9.7	0.7	1.4	1.3	0.5

n.d. not detected

determine the optimal time and growth of the bacterial strains. Microalgal extracts were preabsorbed for 4 h at 20°C to increase assay sensitivity.

To assess the differences in measurements of the inhibition zone, we performed one-way analysis of variance (ANOVA). Significant differences were analyzed using Tukey's a posteriori test. We considered all of the assumptions of the statistical test (Zar 1996). All statistical analyses were performed using STATISTICA, version 6.0 with  $\alpha = 0.05$ .

## RESULTS AND DISCUSSION

The cultured microalgae and cyanobacteria strains showed taxon-dependent differences in cell size, density, dry weight, and growth rate (Fig. 1, Table 1). As expected, cyanobacteria cultures were the smallest and had the highest density, and *Dunaliella tertiolecta* cultures had the largest cell size and the lowest density. The *Amphiprora paludosa* strain had the lowest cellular mass, expressed as dry weight (Table 1).

In the antibacterial activity assays, certain parameters of growth, such as growth phase, cell density, exponential growth rate, mean doubling time, and dry weight, were relevant, because these variables can influence the




Figure 1. Growth of batch cultures of marine microalgae and cyanobacteria according to the mean optical density (OD) from triplicates measurements / Crecimiento de cultivos estáticos de microalgas y cianobacterias marinas según el valor de densidad óptica promedio (OD) de triplicados

antibacterial activity mentioned above. All microalgae cultures had CFU values that were lower than  $3.0 \times 10^3$  mL<sup>-1</sup> (Table 1).

Based on the screen, the *Chaetoceros muelleri* methanolic extract had the most potent antibacterial activity against *B. subtilis* and *S. aureus*, generating inhibition halos of 8.53-15.86 mm (Table 2). Of the 18 algal extracts that we tested, *S. elongatus*, *Synechocystis* sp., *A. paludosa*, and *P. cruentum* had antibacterial activity against Gram(+) and Gram(-) bacteria. Antibacterial activity against *M. luteus* or *E. coli* was not detected in any extract (Table 2).

The cyanobacterial extracts from *S. elongatus* and *Synechocystis* sp. had antibacterial activity against *B. subtilis* and *S. aureus* (inhibition halos: 7.96-9.86 mm) and the Gram(-) *K. pneumoniae* (inhibition halos: 9.76-9.93 mm) (Table 2).

Extracts from the benthic diatom *A. paludosa* generated inhibition halos of 7.60-11.80 mm against *B. subtilis* and *S. aureus*. Against *P. aeruginosa*, only the methanol extract had antibacterial activity (mean value 10.53 mm) (Table 2).

Extracts of the Rhodophyceae *P. cruentum* in all 3 solvents inhibited the growth of *B. subtilis* (inhibition halos of 7.06-9.80 mm). Only the dichloromethane extract was active against *P. aeruginosa* and *K. pneumoniae*, effecting mean inhibition halos of 9.83 and 11.00 mm, respectively (Table 2).

The extracts that we used might contain compounds that have antibacterial activity. To identify and evaluate the activity of individual compounds, they must undergo further purification and separation. Microalgae comprise a diverse group of photosynthetic organisms that have successfully adapted their metabolism to survive in different habitats. The ability to withstand environmental stress is related to their ability to produce a vast array of secondary metabolites, which have considerable value in the biotechnology, aquaculture, health, and food industries (Andersen 1996). Antibacterial activity has been observed predominantly in the Bacillariophyceae Class, the most notable species of which are *Chaetoceros*, *Nitzschia*, and *Thalassiosira*. Strain selection is important, because biological activities can differ between strains of the same species (Pesando 1990).

Our results confirm the reported antibacterial properties of *Chaetoceros* spp. and *P. cruentum* (Pesando 1990, Ördög et al. 2004). Our data suggest that the antibacterial

Table 2. Mean inhibition halos in organic extracts from batch cultures of marine cyanobacteria and microalgae using hexane (H), dichloromethane (D), and methanol (M). Standard deviation are shown in brackets. a,b,c values indicate statistical significance (ANOVA and Tukey a posteriori test,  $P < 0.05$ :  $a < b < c$ ) / Valores promedio de los halos de inhibición en los extractos orgánicos de cultivos estáticos de cianobacterias y macroalgas marinas utilizando hexano (H), diclorometano (D), y metanol (M). La desviación estándar se indica en paréntesis. Valores a,b,c indican diferencias estadísticas (ANDEVA y prueba a posteriori de Tukey,  $P < 0,05$ :  $a < b < c$ )

Strains	Extracts	<i>Staphylococcus aureus</i>	<i>Bacillus subtilis</i>	<i>Micrococcus luteus</i>	<i>Escherichia coli</i>	<i>Klebsiella pneumoniae</i>	<i>Pseudomonas auruginosa</i>
<i>Synechococcus elongatus</i>	H	7.96 (0.41) a	8.83 (0.32) a	-	-	9.93 (0.11) a	-
	D	9.86 (0.15) c	9.23 (0.09) b	-	-	9.76 (0.25) a	-
	M	8.66 (0.37) b	8.63 (0.35) a	-	-	-	-
<i>Synechosystis</i> sp.	H	9.56 (0.37) a	8.20 (0.20) a	-	-	9.83 (0.28) a	-
	D	9.33 (0.57) a	9.76 (0.25) b	-	-	9.86 (0.23) a	-
	M	9.40 (0.60) a	9.36 (0.55) b	-	-	-	-
<i>Amphiprora paludosa</i>	H	8.66 (0.40) a	11.76 (0.40) c	-	-	-	-
	D	10.76 (0.25) b	10.13 (0.75) b	-	-	-	-
	M	11.80 (0.52) c	7.60 (0.40) a	-	-	-	10.53 (0.41)
<i>Dunaliella tertiolecta</i>	H	10.00 (1.03) a	11.60 (0.60) c	-	-	-	-
	D	9.23 (0.25) a	8.93 (0.11) a	-	-	-	-
	M	9.70 (0.26) a	10.06 (0.11) b	-	-	-	-
<i>Chaetoceros muelleri</i>	H	8.96 (0.32) b	8.53 (0.41) a	-	-	-	-
	D	8.70 (0.36) a	9.70 (0.26) b	-	-	-	-
	M	15.86 (0.30) c	15.23 (0.32) c	-	-	-	-
<i>Porphyridium cruentum</i>	H	-	9.80 (0.62) b	-	-	-	-
	D	-	7.06 (0.40) a	-	-	11.00 (0.20)	9.83 (0.32)
	M	-	7.20 (0.26) a	-	-	-	-
Negative Control		-	-	-	-	-	-
Positive Control		13.13 (0.72)	16.36 (0.11)	10.56 (0.66)	12.00 (0.20)	10.26 (0.11)	13.03 (0.25)

activities of these strains are primarily Gram(+) -specific, perhaps due to the additional membrane that exists in Gram(-) bacteria. The active compound in the extracts might encounter difficulties in penetrating the additional membrane, which contains porins that impede the influx of drugs and a multiple-drug pump efflux system that expels most drugs from the bacteria, a well-documented phenomenon (Schmidt & Hansen 2001).

One of the cyanobacteria (*Synechococcus elongatus*) that we used were isolated from commercial aquaculture farms from a shrimp farm in Nayarit, México (Aguilar-May & Sánchez-Saavedra 2007) and *Synechosystis* sp. from an abalone farm in Eréndira, Baja California, México. The antibacterial activity of local *S. elongatus* and *Synechosystis* sp. strains had not been reported.

Cyanobacteria have the ability to synthesize antibiotic compounds could benefit their high growth rate and survival under extreme conditions in their competitive

natural environment (Patil *et al.* 2009). This capacity might be useful for the isolation and production of new antibacterial compounds. For example, they could be used as a probiotic in shrimp farms to eliminate pathogenic bacteria, such as members of the genus *Vibrio*. *Synechococcus elongatus* has been used to control eutrophication and remove nitrogen and phosphorous by immobilization onto chitosan cells (Aguilar-May & Sánchez-Saavedra 2009) and as food for postlarval shrimp (Sánchez-Saavedra & Moreno-Pérez 2009<sup>1</sup>).

Our data implicate the novel antibacterial activity of the benthic diatom *Amphiprora paludosa*, which is a common strain that is used as feed for postlarval abalone cultures in México (Correa *et al.* 2009). Concentrated cultures of *A. paludosa* can be stored at 4°C in the dark for two months. Under these conditions, the cells maintain their viability due to their tolerance to low temperatures and antibacterial activity, as evidenced by their low CFU

<sup>1</sup>Sánchez-Saavedra MP & MPA Moreno-Pérez. 2009. Effect of monospecies and mixed diets on the growth and survival of shrimp larvae *Litopenaeus vannamei* (Bonne, 1931). World Aquaculture Symposium, Veracruz, México, 25 al 29 de septiembre. p. 93.



values ( $1 \times 10^4$  mL<sup>-1</sup>) in storage cultures and low bacterial colony counts by epifluorescence microscopy (Sánchez-Saavedra 2006).

*Dunaliella tertiolecta*, *Choetoceros muelleri*, and *Porphyridium cruentum* strains showed antibacterial activity against *B. subtilis*, and only the dichloromethane extract of *P. cruentum* inhibited the growth of *K. pneumoniae* and *P. auruginosa*. Antibacterial activity of *P. cruentum* has been reported against *Vibrio* spp. (Cooper *et al.* 1983), and *C. muelleri* has been tested against *Vibrio* and *Pseudomonas* (Pesando 1990). Ethanolic *D. salina* extracts have shown antimicrobial activity against *S. aureus* and *E. coli*. In these extracts, 15 volatile compounds and several fatty acids (primarily palmitic,  $\alpha$ -linolenic, and oleic acids) were identified (Herrero *et al.* 2006).

Antibacterial activity, observed in diatoms *Chaetoceros* and *Skeletonema*, has been associated with several fatty acids, primarily those with 10 carbon atoms, which induced lysis in bacterial protoplasts (Naviner *et al.* 1999). The strains *S. elongatus*, *Synechosystis* sp., *A. paludosa*, *C. muelleri*, and *P. cruentum* can potentially be used to control Gram(+) and Gram(-) bacteria. Our results showed that several microalgal strains could constitute a biological method of bacterial control under controlled culture conditions due to their antibacterial activity. Yet, the specificity of the chemical compounds that are produced from the microalgae cultures for a specific bacterial group must be considered. All antibacterial activities in the microalgae cultures were attributed to the cells -not to the bacteria- based on the low or undetectable CFU counts from the microalgal culture samples. In the sea, the bacterial concentration approaches  $10^6$  cells mL<sup>-1</sup> and exceeds  $10^6$  cells mL<sup>-1</sup> in aquaculture systems (Maeda *et al.* 1997).

In conclusion, this work demonstrates the vital role of certain marine microalgae and cyanobacteria in the production of antibacterial substances and lays the foundation for future applications by the aquaculture and pharmaceutical industries. Future studies should identify the compounds that mediate the antibacterial activity that we have observed.

## ACKNOWLEDGMENTS

This work was supported by Consejo Nacional de Ciencia y Tecnología de México (CONACyT, -Project-SAGARPA-239 and CN-04-127). Thanks to Darla-Torres-Ariño, Juan-Fernández-Apango, and Mario-Navarro-Armenta for

technical assistance and Lilia-Hurtado-Ayala for supplying bacteria strains. Thanks to Dr. Ellis-Glazier and Blue-Pencil-Science for editing this English-language text.

## LITERATURE CITED

- Aguilar-May B & MP Sánchez-Saavedra. 2009.** Growth and removal of nitrogen and phosphorous by free-living and chitosan-immobilized cells of the marine cyanobacterium *Synechococcus elongatus*. *Journal of Applied Phycology* 21: 353-360.
- Aguilar-May B, MP Sánchez-Saavedra, J Lizardi & D Voltolina. 2007.** Growth of marine *Synechococcus* sp. immobilized in chitosan with different times of contact with NaOH. *Journal of Applied Phycology* 19(2): 181-183.
- Al-Saimary IE, SS Bakr, BY Khudair & YA Abass. 2007.** Efficiency of antibacterial agents extracted from *Thymus vulgaris* L. (Lamiaceae). *The Internet Journal of Nutrition and Wellness* 4(1): [on line] <<http://www.ispub.com/ostia/index.php?xmlFilePath=journals/ijnw/vol4n1/thymus.xml>>
- Andersen RA. 1996.** Maintaining cultures for biotechnology and industry. In: Hunter-Cervera JC & Belt A (eds). *Algae*, pp. 29-64. Academic Press, London.
- Collins CH, PM Lynes & JM Granje. 1989.** Antimicrobial sensibility and assay test. In: Collins CH & PM Lynes (eds). *Microbial methods*, pp. 155-168. Butterworth, London.
- Colwell RR, RK Zisemore, JF Carmey, RY Morita, JDJ Nelson, JH Pickard, JR Schwatz, V Valkenburg, SD Walker & TR Wright. 1975.** Marine and estuarine microbiology. Laboratory manual, 96 pp. University Park Press, Baltimore.
- Cooper S, A Battat, P Marsot & M Sylvestre. 1983.** Production of antibacterial activities by two Bacillariophyceae grown in dialysis culture. *Canadian Journal of Microbiology* 29: 338-341.
- Correa RJG, MP Sánchez-Saavedra, MT Viana, N Flores-Acevedo & C Vásquez-Peláez. 2009.** Effect of eight benthic diatoms as feed on the growth of red abalone postlarvae (*Haliotis rufescens*). *Journal of Applied Phycology* 21(4): 387-393.
- CLSI. 2006.** Performance standards for antimicrobial disk susceptibility test; Approved Standard-Ninth Edition. National Committee for Clinical Laboratory Standards, Wayne, Pennsylvania Document M2-A9 26(1): 7-9.
- Demain AL. 1999.** Pharmaceutical active secondary metabolites of microorganisms. *Applied Microbial Biotechnology* 52: 455-463.
- FDA Agents. 2009.** Food and Drug Administration.-[on line] <[http://en.citizendium.org/wiki/Food\\_and\\_Drug\\_Administration](http://en.citizendium.org/wiki/Food_and_Drug_Administration)>
- Fogg GA & B Thake. 1987.** Algal cultures and phytoplankton ecology, pp. 12-42. The University of Wisconsin Press, Madison.

- Guillard RLL. 1975.** Culture of phytoplankton for feeding marine invertebrates. In: Smith ML & MH Chanley (eds). Culture of marine invertebrates animals, pp. 29-60. Plenum Press, New York.
- Herrero M, E Ibáñez, A Cifuentes, G Reglero & S Santoyo. 2006.** *Dunaliella salina* microalga pressurized liquid extracts as potential antimicrobial. Journal of Food Protection 69(10): 2471-2477.
- Kreitlow S, S Mundt & U Lindequist. 1999.** Cyanobacteria - a potential source of new biologically active substances. Journal of Biotechnology 70: 61-63.
- Maeda M, K Nogami, M Kanematsu & K Hirayama. 1997.** The concept of biological control in aquaculture. Hydrobiologia 358: 285-290.
- Maniatis T, EF Fritsh & J Sambrook. 1982.** Molecular cloning. A laboratory manual, 458 pp. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York.
- Naviner M, JP Bergé, P Durand & H Le Bris. 1999.** Vibriostatic bacteria isolated from rearing seawater of oyster brood stock: Potential as biocontrol agent for vibriosis in oyster larvae. Fish Pathology 34: 139-144.
- Ördög V, WA Stirk, R Lenobel, M Bancírová, M Strnad, J Staden, J Szigeti & LNémeth. 2004.** Screening microalgae for some potentially useful agricultural and pharmaceutical secondary metabolites. Journal of Applied Phycology 16: 309-314.
- Patil LS, MV Kulkarani & PR Puranik. 2009.** Anthelmintic effects of extracts of some indigenously isolated culturable cyanobacterial species. Pharmacologyonline 2: 879-886.
- Pesando D. 1990.** Antibacterial and antifungal activities of marine algae. In: Akatsuka I (ed). Introduction to applied phycology, pp. 3-26. SPB Academic Publishing, Amsterdam.
- Riquelme C & RE Avendaño-Herrera. 2003.** Interacción bacteria-microalga en el ambiente marino y uso potencial en acuicultura. Revista Chilena de Historia Natural 76: 725-736.
- Sánchez-Saavedra MP. 2006.** The effect of cold storage on cell viability and composition of two benthic diatoms. Aquacultural Engineering 34: 131-136.
- Schmidt LE & PH Hansen. 2001.** Allelopathy in the prymnesiophyte *Chrysochromulina polylepis*: effect of cell concentration, growth phase and pH. Marine Ecology Progress Series 216: 67-81.
- Shimizu Y. 1993.** Microalgal metabolites. Chemistry Review 93: 1685-1698.
- Singh IP, KE Milligan & WH Gerwick. 1999.** Tanikolide, a toxic and antifungal lactone from the marine cyanobacterium *Lyngbya majuscula*. Journal of Natural Products 62: 1333-1335.
- WHO. 2006.** Antimicrobial use in aquaculture and antimicrobial resistance. Consultations and Workshops. Report of a Joint FAO/OIE/WHO. Expert Consultation, Seoul, Republic of Korea. 13-16 June 2006, 97 pp. World Health Organization, Geneva.
- Zar JH. 1996.** Biostatistical analysis, 718 pp. Prentice Hall, Englewood Cliffs.

---

Recibido el 8 de diciembre de 2009 y aceptado el 14 de julio de 2010