



Revista de Biología Marina y Oceanografía

ISSN: 0717-3326

revbiolmar@gmail.com

Universidad de Valparaíso

Chile

Galleguillos, Ricardo; Canales-Aguirre, Cristian B.; Ferrada, Sandra; Barrera, Andrea
SSRs in *Octopus mimus*: development and characterization of nine microsatellite loci
Revista de Biología Marina y Oceanografía, vol. 46, núm. 3, diciembre, 2011, pp. 491-494
Universidad de Valparaíso
Viña del Mar, Chile

Available in: <http://www.redalyc.org/articulo.oa?id=47922575019>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

SSRs in *Octopus mimus*: development and characterization of nine microsatellite loci

SSRs en *Octopus mimus*: desarrollo y caracterización de nueve loci microsatélites

Ricardo Galleguillos¹, Cristian B. Canales-Aguirre¹,
Sandra Ferrada¹ and Andrea Barrera¹

¹Laboratorio de Genética y Acuicultura, Departamento de Oceanografía, Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción, Casilla 160-C, Concepción, Chile. cristiancanales@udec.cl

Abstract. - Nine microsatellite loci were developed for *Octopus mimus*, a cephalopod of commercial importance for artisanal fishermen. Genetic variation at these loci was examined in samples from Clavelito, a Benthic Resources Management Area (AMERB, in Spanish). All nine loci were highly polymorphic, with the number of alleles per locus ranging from 4 to 28 and the expected heterozygosity ranging from 0.651 to 0.946. These markers will be useful to address issues of population genetics, ecology, conservation and fisheries management related to that species.

Key words: Population genetics, AMERBs, cephalopod

INTRODUCTION

To evaluate how genetic variability is distributed it is necessary to use polymorphic molecular markers. In the last few years, microsatellites have been some of the most used molecular markers for determining genetic variability and genetic structure levels (Barbará *et al.* 2008). These codominant markers are highly polymorphic, with motifs of 1 to 6 nucleotides in length, organized in tandems, widely distributed along the genome (Tautz 1989).

In cephalopods as in other marine groups, few studies have been devoted to identify and characterize polymorphic microsatellite loci. This is unusual, due to the low genetic diversity that has been observed using other molecular markers in cephalopods (*i.e.*, Allozymes: Brierley *et al.* 1995, Triantafillos *et al.* 2004 and mitochondrial DNA: Oosthuizen *et al.* 2004).

Octopus mimus (Gould, 1852), is a member of Octopodidae Family, it inhabits rocky shore systems in the Southeast Pacific from northern Perú to central Chile. Biological traits such as separate sexes, internal fertilization and a semelparous reproductive strategy (Leite *et al.* 2008) make this species a good model to answer questions about genetic connectivity. Most studies in the Genus *Octopus* have focused on determining the phylogeny (*e.g.*, Warnke 2004) and only a few focus on their population biology. Even a smaller number of these studies used microsatellite loci as a tool for determining genetic structure (*e.g.*,

Greatorex *et al.* 2000, Cabranes *et al.* 2008, Doubleday *et al.* 2009). Up until now, there is no information about *O. mimus* population genetics. In this paper, we describe the isolation, characterization and utility of nine microsatellite loci for *O. mimus* in one Chilean resource management area.

MATERIALS AND METHODS

High molecular weight DNA was extracted from arm tissue of 20 individuals from both sexes using a salting-out protocol (Jowett 1986). The genomic library was developed by Genetic Identification Services¹. Briefly, the DNA was partially digested with a cocktail of seven blunt-end restriction enzymes (*Hae*III, *Stu*I, *Eco*RV, *Sca*I, *Bsr*BI, *Pvu*II, *Hin*CII). Fragments between 350 and 700 bp were selected by gel extraction and ligated to a 20 bp oligonucleotide adaptor containing a Hind III restriction site at the 5' end. Microsatellite enrichment was achieved using streptavidin-coated magnetic beads and 5'-biotinylated CA₁₅, AAC₁₂, TACA₈ and TAGA₈ oligonucleotide probes. The captured molecules were amplified by PCR using a primer complementary to the adaptor, digested with *Hind*III to remove the adaptor, and ligated into the *Hind*III site of the pUC19 vector. The plasmids were then electroporated into *Escherichia coli* DH5. Recombinant clones, identified by blue-white

¹Genetic Identification Services, Chatsworth. <<http://www.genetic-id-services.com>>

selection, were chosen arbitrarily for sequencing on an ABI 377 using the Big Dye Terminator Cycle Sequencing methodology (Applied Biosystems). Specific primers flanking the identified microsatellite sequences were designed using Designer PCR version 1.03 (Research Genetics) (Table 1). The microsatellite loci were amplified in 10 µl reactions containing 1X PCR buffer, 2 mM MgCl₂, 0.2 mM forward primer (fluorescently labeled), 0.2 mM reverse primer, 0.2 mM dNTPs, 0.1 U µl⁻¹ Taq DNA polymerase (Invitrogen), and 20 ng of genomic DNA template. PCR was performed in a PTC-200 (MJ Research) thermal cycler with the following parameters: 94°C for 3 min, followed by 34 cycles of 94°C for 40 s, 56°C for 40 s, 72°C for 30 s, and a final extension at 72°C for 4 min. All loci successfully amplified under the same conditions. The PCR products were analyzed on an ABI 3330 DNA sequencer. Alleles were scored using Peak Scanner v1.0, with GS500 (Applied Biosystems) as the internal size standard.

To characterize the polymorphism of each locus we used 100 individuals from Clavelito, Chile. To evaluate the potential presence of null alleles and genotyping artifacts (stutter bands or large dropout alleles) we first

checked the data set with MICRO-CHEKER v2.2.3 software (Van Oosterhout *et al.* 2004). The number of alleles (Na), the expected (He) and observed (Ho) levels of heterozygosity were obtained using GENALEX v6 (Peakall & Smouse 2006). Deviations from Hardy-Weinberg equilibrium (H&W) and gametic disequilibrium between markers were tested using ARLEQUIN v3.1 (Excoffier *et al.* 2005). All probability values were estimated using 10000 permutations.

RESULTS AND DISCUSSION

We observed a significant departure from HWE in four loci (*i.e.*, *OmimA2*, *OmimB111*, *OmimC1* and *OmimC122*). None of the studied loci showed gametic disequilibrium. On the other hand, the results did not show any score errors like large allele dropout and stutter bands. Two loci showed heterozygotes deficiency: *OmimA105* and *OmimB111* (Table 1) which, in this case, could be attributed mainly to: a) admixture of individuals captured over a large geographical area that may include more than one panmictic unit (Wahlund effect), b) null alleles, as some of the sequences (allele) could not be amplified, the number of heterozygotes is sub estimated, c) inbreeding

Table 1. Characteristics of the microsatellite loci isolated for the *Octopus mimus*. The values reported for each marker are the allele size range (bp), the number of alleles (Na), the observed (Ho) and the expected (He) heterozygosity, and the GenBank accession number. *Omim*: *Octopus mimus*. *: $P < 0.05$; **: $P < 0.01$ / Características de los loci microsatélites aislados en *Octopus mimus*. Los valores reportados para cada marcador son el rango del tamaño alélico (pb), el número de alelos (Na), la heterocigosidad observada (Ho) y esperada (He), y el número de acceso a GenBank. *Omim*: *Octopus mimus*. *: $P < 0,05$; **: $P < 0,01$

Locus	N° Genbank	Primer sequence (5'-3')	Repeat unit	Size range (bp)	Clavelito		
					Na	Ho	He
<i>OmimA2</i>	GU584076	F: TCAATGGATGTTGATGTCAGA G R: TTCCAGGCAAGGTAATATCTTG	(AT) ₄ (GT) ₃ N ₂ (GT) ₆ N ₄ (GT) ₃ N ₇ (AT) ₃ N ₂ (GT) ₁₄	268-306	8	0.893	0.816*
<i>OmimA103</i>	GU584077	F: CATTGCTGAACGAAGAACTC R: GCTTATCCCGAACAGGTAAT	(ATCT) ₆	218-350	28	1	0.939
<i>OmimA104</i>	GU584078	F: TGGTTGATAGAAACTGTGTCG R: GTTGTGGTGATGGTGACTG	(CCA) ₆ N ₃ (CCA) ₂	186-276	13	0.936	0.880
<i>OmimA105</i>	GU584079	F: GCCCTTTTGTGTAATAATACG R: TTCTCGCTAATGGCAGTTC	(AC) ₂₈	194-270	24	0.850	0.946
<i>OmimB101</i>	GU584080	F: CACCAAGTCACAGACATCTTG R: GCCAGATAAACAAACAATAGCTG	(CA) ₅₄	206-256	4	0.467	0.651
<i>OmimB111</i>	GU584081	F: TCCTGATTCAAGTCATTTACAGG R: GGGTCTGTTTCTTTGACATG	(GT) ₃ (GA) ₁₅	186-300	8	0.273	0.826**
<i>OmimC1</i>	GU584082	F: CGTGTGTCAACCAATTTC R: GGTGTAGTTAAATGGCATACTC	(ATGT) ₁₃ (GT) ₁₂ N ₂ (GT) ₅ N ₂ (GT)	133-213	14	0.938	0.805**
<i>OmimC106</i>	GU584083	F: TGCCATCAAAGTGACACTG R: AACCACAGATACTCGAAGAATG	(TGTA) ₂₀ N ₄ (TGTA) ₉	188-300	10	0.862	0.747
<i>OmimC122</i>	GU584084	F: ATTGTTGAAATCGCAACTGT R: AATTCTCATAATCGCTGTTGTG	(ACT) ₁₉	118-256	17	0.942	0.800**

and d) selection. According to our results, null alleles are the most plausible cause of heterozygotes deficiency at these two loci, as suggested by the excess of homozygotes obtained by the Micro-Checker software. Heterozygosity deficits due to null alleles have been recorded in *Octopus* species, (e.g., *O. vulgaris*, Casu *et al.* 2002, Cabranes *et al.* 2008, *O. maorum*, Doubleday *et al.* 2008, *O. maya*, Juárez *et al.* 2010). The presence of heterozygosity deficit could support the low variability recorded in other molecular markers used in cephalopods (Allozyme, Maltagliati *et al.* 2002; mitochondrial DNA, Oosthuizen *et al.* 2004). Moreover, life history traits like semelparity, low fecundity and territorial behavior could promote inbreeding and consequently the heterozygosity deficit.

The Na per locus ranged from 4 to 28, the Ho from 0.273 to 1 and the He varied from 0.651 to 0.946 (Table 1). Polymorphism found in microsatellite loci of *O. mimus* is comparable with other studies in other members of the genus *Octopus*. He in *O. mimus* (mean He = 0.823) was less than He in *O. vulgaris* (mean He = 0.874, Cabranes *et al.* 2008; mean He = 0.91, Murphy *et al.* 2002) and *O. maorum* (mean He = 0.85, Doubleday *et al.* 2008). On the other hand, average He in *O. mimus* was greater than He in *O. maya* (mean 0.645, Juárez *et al.* 2010) and *O. vulgaris* (mean He = 0.765, Grotorex *et al.* 2000).

In summary, this work provides a significant genetic tool for future population genetic studies, which should improve government management policies to maintain the sustainability of these important fisheries, and helping not only to promote the conservation of the fishing stock, but also to conserve genetic diversity of this cephalopod along the Chilean coast.

ACKNOWLEDGMENT

Sandra Ferrada and Cristian B. Canales-Aguirre were supported by Doctoral Fellowships for the 'Programa de Doctorado en Sistemática y Biodiversidad', from the graduate school of the Universidad de Concepción. Sandra Ferrada was supported by a CONICYT doctoral fellowship. This work forms part of the FIP 2008-39 Project.

LITERATURE CITED

Barbará T, C Palma-Silva, GM Paggi, F Bered, MF Fay & C Lexer. 2007. Cross-species transfer of nuclear microsatellite markers: potential and limitations. *Molecular Ecology* 16: 3759-3767.

Brierley AS, JP Thorpe, GJ Pierce, MR Clarke & PR Boyle. 1995. Genetic variation in the neritic squid *Loligo forbesi* (Myopsida: Loliginidae) in the northeast Atlantic Ocean. *Marine Biology* 122: 79-86.

Cabranes C, P Fernandez-Rueda & JL Martínez. 2008. Genetic structure of *Octopus vulgaris* around the Iberian Peninsula and Canary Islands as indicated by microsatellite DNA variation. *ICES Journal of Marine Science* 65: 12-16.

Casu M, F Maltagliati, M Meloni, D Casu, P Cossu, G Binelli, M Curini-Galletti & A Castelli. 2002. Genetic structure of *Octopus vulgaris* (Mollusca, Cephalopoda) from the Mediterranean Sea as revealed by a microsatellite locus. *Italian Journal of Zoology* 69: 295-300.

Doubleday ZA, GT Pecl, JM Semmens & LV Danyushevsky. 2008. Using styler elemental signatures to determine the population structure of *Octopus maorum*. *Marine Ecology Progress Series* 360: 125-133.

Doubleday ZA, JM Semmens, AJ Smolenski & PW Shaw. 2009. Microsatellite DNA markers and morphometrics reveal a complex population structure in a merobenthic octopus species (*Octopus maorum*) in south-east Australia and New Zealand. *Marine Biology* 156: 1183-1192.

Excoffier L, G Laval & S Schneider. 2005. Arlequin 3.01: An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* 1: 47-50.

Grotorex E, C Jones, J Murphy, L Key, A Emery & P Boyle. 2000. Microsatellite markers for investigating population structure in *Octopus vulgaris* (Mollusca: Cephalopoda). *Molecular Ecology* 9: 641-642.

Jowett T. 1986. Preparation of nucleic acids. In: Roberts DB (ed). *Drosophila: A practical approach*, pp. 275-286. IRL Press, Oxford.

Juárez OE, C Rosas & L Arena. 2010. Heterologous microsatellites reveal moderate genetic structure in the *Octopus maya* population. *Fisheries Research* 106: 209-213.

Leite T, M Haimovic, W Molina & K Warnke. 2008. Morphological and genetic description of *Octopus insularis*, a new cryptic species in the *Octopus vulgaris* complex (Cephalopoda: Octopodidae) from tropical southwestern Atlantic. *Journal of Molluscan Studies* 74: 63-74.

Maltagliati F, P Belcari, D Casu, M Casu, P Sator, G Vargiu & A Castelli. 2002. Allozyme genetic variability and gene flow in *Octopus vulgaris* (Cephalopoda, Octopodidae) from the Mediterranean Sea. *Bulletin of Marine Sciences* 71: 437-486.

Murphy JM, E Baguerias, LN Key & PR Boyle. 2002. Microsatellite DNA markers discriminate between two *Octopus vulgaris* (Cephalopoda: Octopoda) fisheries along the northwest African coast. *Bulletin of Marine Science* 71: 545-553.

- Oosthuizen A, M Jiwaji & P Shaw. 2004.** Genetic analysis of the *Octopus vulgaris* population on the coast of South Africa. *South African Journal of Science* 100: 603-607.
- Peakall R & PE Smouse. 2006.** GENALEX 6: genetic analysis in Excel. Population software for teaching and research. *Molecular Ecology Notes* 6: 288-295.
- Tautz D. 1989.** Hypervariability of simple sequences as a general source of polymorphic DNA markers. *Nucleic Acids Research* 17: 6463-6471.
- Triantafillos L, GD Jackson, M Adams & BL McGrath-Steer. 2004.** An allozyme investigation of the stock structure of arrow squid *Nototodarus gouldi* (Cephalopoda: Ommastrephidae) from Australia. *ICES Journal of Marine Science* 61: 829-835.
- Van Oosterhout C, WF Hutchinson, DPM Wills & P Shipley. 2004.** MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* 4: 535-538.
- Warnke K, R Söller, D Blohm & U Saint-Paul. 2004.** A new look at geographic and phylogenetic relationships within the species group surrounding *Octopus vulgaris* (Mollusca, Cephalopoda): indications of very wide distribution from mitochondrial DNA sequences. *Journal of Zoological Systematics and Evolutionary Research* 42: 306-312.

Received 12 February 2011 and accepted 6 October 2011