



Revista de Biología Marina y Oceanografía

ISSN: 0717-3326

revbiolmar@gmail.com

Universidad de Valparaíso

Chile

Ortega, Karla J.; Sáez, Claudio A.; Macaya, Erasmo C.  
Changes in invertebrate assemblages inhabiting *Lessonia spicata* (Phaeophyceae) holdfasts after the  
2010 earthquake-mediated coastal uplift in Chile  
Revista de Biología Marina y Oceanografía, vol. 49, núm. 1, abril, 2014, pp. 129-134  
Universidad de Valparaíso  
Viña del Mar, Chile

Available in: <http://www.redalyc.org/articulo.oa?id=47931255014>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System  
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal  
Non-profit academic project, developed under the open access initiative

RESEARCH NOTE

## Changes in invertebrate assemblages inhabiting *Lessonia spicata* (Phaeophyceae) holdfasts after the 2010 earthquake-mediated coastal uplift in Chile

Cambios en los ensambles de invertebrados asociados a discos del huiro negro *Lessonia spicata* (Phaeophyceae) después del levantamiento costero producto del terremoto de 2010 en Chile

Karla J. Ortega<sup>1</sup>, Claudio A. Sáez<sup>2,4</sup> and Erasmo C. Macaya<sup>1,3</sup>

<sup>1</sup>Laboratorio de Estudios Algales ALGALAB, Departamento de Oceanografía, Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción, Casilla 160-C, Concepción, Chile. emacaya@oceanografia.udec.cl

<sup>2</sup>Plymouth University, School of Marine Science & Engineering, Faculty of Science & Environment, Drake Circus, Plymouth, PL4 8AA, United Kingdom

<sup>3</sup>Interdisciplinary Center for Aquaculture Research (INCAR), Universidad de Concepción, P.O. Box 160-C, Concepción, Chile

<sup>4</sup>Present address: Departamento de Biología, Facultad de Química y Biología, Universidad de Santiago de Chile, Casilla 40 Correo 33, Santiago, Chile

**Abstract.** On February 27th 2010, a mega-earthquake struck Chile causing coastal coseismic uplift. In this study we compared the invertebrate fauna from uplifted (UH) and non-uplifted (NH) *Lessonia spicata* kelp holdfasts. Multivariate analysis of the assemblage structure revealed different faunal composition between holdfasts, the anemone *Phymactis clematis* and the mussel *Perumytilus purpuratus* displayed the highest contribution to these differences, being most abundant in UH. A preference to inhabit in UH is most likely to be related to food availability and capacity to thrive with high salinity and desiccation, while NH became habitat for a rare primary ecological succession. The present study helps to understand ecological implications of such rare events.

**Key words:** Coastal coseismic uplift, kelps, associated fauna, assemblages analysis

### INTRODUCTION

On February 27<sup>th</sup> 2010 (27F), one of the strongest earthquakes ( $M_w$  8.8) in the last 150 years struck Chile along the central-southern coast (Farías *et al.* 2010, Vargas *et al.* 2011) causing coastal coseismic uplift of up to ~3 m at locations in proximity to the earthquake rupture (Castilla *et al.* 2010, Vargas *et al.* 2011, Melnick *et al.* 2012). This event produced massive mortality of intertidal communities suddenly exposed to drastic changes in biotic and abiotic factors such as predation, desiccation, salinity and temperature stress (Castilla *et al.* 2010, Jaramillo *et al.* 2012).

Previous studies have demonstrated the effect of sudden rocky shore uplifts on kelps and its associated assemblages (Castilla 1988, Castilla & Oliva 1990). Castilla & Oliva (1990) observed that after a coastal uplift caused by an earthquake in 1985 in central Chile, the intertidal kelp belt expanded downwards in approximately 1 m; moreover, the old uplifted middle-intertidal was colonized by species of barnacles, organisms characteristic of the upper-intertidal. Kelps are ecosystem bio-engineers; they are main primary producers and play an important

ecological role in coastal rocky shores of temperate environments (Vásquez & Santelices 1984, Smith *et al.* 1996, Smith 2000). The complex and compact holdfasts of kelps provide a valuable habitat for a high diversity of small invertebrates (Tuya *et al.* 2011); these micro-ecosystems give shelter, access to food, as well as nursery and spawning sites (Cancino & Santelices 1984, Ojeda & Santelices 1984, Villouta & Santelices 1986, Thiel *et al.* 2007). Along the central Chilean coast, the kelp *Lessonia spicata* is the dominant algal species in the low intertidal, and their holdfasts host high species richness (Vásquez & Santelices 1984). This species was previously recognized as *L. nigrescens* until recent molecular, morphological and nomenclatural studies led to the recognition of a separate species between 29°-41°S in central-southern Chile (Tellier *et al.* 2009, González *et al.* 2012).

Many studies have highlighted the use of kelp holdfasts faunal assemblages as bio-monitor following environmental perturbations (*e.g.*, Jones 1972, Smith & Simpson 1993, Smith *et al.* 1996, Smith 2000, Sáez *et al.*


2012). In this study we took advantage of the coseismic uplift in Chile to study the effects of changes in the vertical positioning of the *L. spicata* belts on the structure of fauna inhabiting uplifted holdfasts (UH), to that occurring in new non-uplifted holdfasts (NH) (Fig. 1 C, D).

## MATERIALS AND METHODS

The study was carried out at ‘La Desembocadura’, Biobío Region, Chile (36°48’25.92”S, 73°10’38.08”W), a site known to be uplifted by the 27F mega-earthquake (see Fig.1 A,B). Along the *L. spicata* belt, 5 UH separated by several metres were collected; those holdfasts were very damaged (Fig. 1D), as also observed by Castilla & Oliva (1990) after an earthquake in central Chile in 1985. Additionally, 5 NH were taken. In all cases and after cutting-off the stipes, NH and UH holdfasts with similar sizes ( $20 \pm 2$  cm of base diameter) were detached using a

crowbar and stored in bags. Sampling was carried out 14 months after earthquake, in April 2011.

In the laboratory, holdfasts were rinsed with seawater to promote the detachment of organisms; the remnant was sieved through a 0.5 mm mesh and retained. Organisms were identified to the highest possible level of taxonomic resolution. We calculated the total number of species (S) and Shannon-Wiener Index (H’). In order to compare differences, one-way ANOVA was carried out. Prior to the analyses, tests for normality (Kolmogorov-Smirnov test) and homogeneity of variances (Bartlett test) were performed. For multivariate analysis, data were transformed by  $\log(x+1)$  and a similarity matrix based on Bray-Curtis was constructed (Clarke *et al.* 2006). In order to graphically represent holdfast assemblages between UH and NH, non-metric multidimensional scaling (nMDS) was applied. Analysis of similarities (ANOSIM) was used



**Figure 1.** A) *Lessonia spicata* belt at sampling site ‘La Desembocadura’, arrows display the uplifted *L. spicata* individuals; B) Kelp upper limit, before and after the 27 February 2010 mega-earthquake; C) Non-uplifted *L. spicata* holdfast; D) Uplifted *L. spicata* holdfast (Scale = 10 cm) / A) Cinturón de *Lessonia spicata* en ‘La Desembocadura’, las flechas muestran los individuos levantados; B) Límite superior de *L. spicata* antes y después del mega-terremoto del 27 febrero de 2010; C) Discos de *L. spicata* No-levantados; D) Discos de *L. spicata* levantados (Escala = 10 cm)

to identify significant differences between assemblages belonging to UH and NH. *P*-values were calculated using the Spearman product-moment correlation method with a maximum of 9999 unrestricted permutations (Clarke 1993). A similarities percentages (SIMPER) analysis was used to identify the species that contributed the most to the differences in assemblage's structure between UH and NH. All multivariate procedures were carried out using PRIMER v.6 software package (Plymouth Marine Laboratory 1997).

## RESULTS AND DISCUSSION

The fauna associated with *L. spicata* holdfasts comprised 26 taxa from 5 phyla (Table 1): Annelida, Arthropoda, Mollusca, Cnidaria and Nemertea. The polychaetes had the highest number of families, with 7 representatives. In UH we documented 24 species, from 12 orders and 8 classes; whereas in NH were documented 19 species, from 12 orders and 8 classes. The faunal composition was different between the two types of holdfasts as

**Table 1. Associated fauna inhabiting holdfasts of *L. spicata*, including abundance for each species, family or phylum / Fauna asociada a discos de *L. spicata*, incluyendo abundancia por cada especie, género, familia o phylum**

Uplifted Holdfasts - UH	Non-Uplifted Holdfasts - NH
<b>ANNELIDA</b>	<b>ANNELIDA</b>
Terebellidae (50)	Terebellidae (8)
Nereididae (287)	Nereididae (184)
Syllidae (36)	Syllidae (18)
Phyllodocidae (6)	Phyllodocidae (6)
Polynoidae (9)	Polynoidae (2)
Sabellariidae (13)	Sabellariidae (61)
Lumbrineridae (5)	
<b>ARTHROPODA</b>	<b>ARTHROPODA</b>
Decapoda	Decapoda
<i>Acanthocyclus gayi</i> (74)	<i>Acanthocyclus gayi</i> (12)
<i>Homalaspis crenulatus</i> (1)	<i>Homalaspis plana</i> (4)
Cirripedia (15)	Cirripedia (18)
Isopoda (51)	Isopoda (7)
Amphipoda (293)	Amphipoda (32)
Pycnogonida (1)	Pycnogonida (1)
<b>MOLLUSCA</b>	<b>MOLLUSCA</b>
Bivalvia	Bivalvia
<i>Perumytilus purpuratus</i> (1077)	<i>Perumytilus purpuratus</i> (82)
<i>Semimytilus algosus</i> (1532)	<i>Semimytilus algosus</i> (828)
<i>Brachidontes granulata</i> (4)	<i>Brachidontes granulata</i> (6)
Gastropoda	Gastropoda
<i>Scurria scurra</i> (90)	<i>Scurria scurra</i> (17)
<i>Lottia orbigny</i> (2)	
<i>Fissurella</i> sp. (2)	
Polyplacophora	Polyplacophora
<i>Chiton granosus</i> (7)	<i>Chiton granosus</i> (1)
<i>Chaetopleura peruviana</i> (2)	
<b>CNIDARIA</b>	<b>CNIDARIA</b>
<i>Phymactis clematis</i> (119)	<i>Phymactis clematis</i> (3)
<i>Anthothoe chilensis</i> (6)	
<b>NEMERTEA</b> (308)	<b>NEMERTEA</b> (330)


demonstrated by the nMDS (Fig. 2) and by the results of ANOSIM (R-statistic= 0.432, *P*= 0.0024). SIMPER analysis revealed that the anemone *Phymactis clematis* and the bivalve *Perumytilus purpuratus* contributed the most to the differences between assemblages, each accounting for 19.34% of the differences (Table 2). Abundances of *P. clematis* were 119 in UH versus 3 in NH holdfasts, whereas abundances of *P. purpuratus* were 1077 in UH versus 82 in NH holdfasts (Table 1.). The number of species (S) was significantly different between the two types of holdfasts (*F*= 13.1, *P*< 0.01) but there were no significant differences in the Shannon-Wiener index (*H'*) (*F*= 4.59, *P*= 0.064).

The structure of the assemblages differed clearly between holdfasts in the old and new intertidal. There are only a few studies presenting data on the effects of earthquake-mediated coastal uplifts on intertidal organisms, although these studies have mostly been descriptive, using organisms such as sessile invertebrates and coralline algae as bioindicators of coastal uplift (e.g., Bodin & Klinger 1986, Castilla & Oliva 1990, Awata *et al.* 2008, Castilla *et al.* 2010). Kelp holdfasts have been recognized as 'micro-ecosystems' (Thiel & Vásquez 2000), and the structure of their assemblages has commonly been used as a biomonitoring tools for assessing environmental stress (e.g., Jones 1972, Smith & Simpson 1992, Sáez *et al.* 2012). However, to the extent of our

**Table 2. SIMPER analysis of the species contributing most to dissimilarities in holdfast assemblages structure. Average Bray-Curtis similarity was 23.18%. XAbund 1 = average abundance of Uplifted Holdfasts; XAbund2 = average abundance of Non-Uplifted Holdfasts; XDiss = similarity average; % Contrib = percent contribution to overall similarity between the 2 types of holdfasts / Análisis SIMPER de las especies que contribuyen a las diferencias entre los ensambles asociados a discos. El promedio de disimilitud de Bray-Curtis fue 23,18%. XAbund 1 = promedio de abundancia de UH; XAbund 2 = promedio de abundancia de NH; XDiss = similitud promedio; % Contrib = contribución porcentual de similitud general entre los 2 tipos de discos**

Species	XAbund 1	XAbund 2	XDiss	%Contrib	%Cum
<i>Phymactis clematis</i>	4.79	1.39	2.56	11.03	11.03
<i>Perumytilus purpuratus</i>	6.98	4.42	1.93	8.31	19.34
Amphipoda	5.68	3.5	1.64	7.09	26.43
<i>Anthothoe chilensis</i>	1.95	0	1.46	6.31	32.74
Isopoda	3.95	2.08	1.41	6.07	38.8
Lumbrineridae	1.79	0	1.35	5.81	44.61
<i>Acanthocyclus gayi</i>	4.32	2.56	1.32	5.68	50.29
Terebellidae	3.93	2.2	1.3	5.62	55.92
<i>Scurria scurra</i>	4.51	2.89	1.22	5.25	61.17
<i>Homalaspis plana</i>	0	1.61	1.21	5.22	66.39





**Figure 2. A) Non-parametric multidimensional scaling plot (nMDS) and ANOSIM of the invertebrate assemblages associated with UH (Uplifted) and NH (Non-Uplifted) holdfasts. B) Species number. C) Shannon-Wiener index of associated fauna inhabiting the 2 types of *L. spicata* holdfasts. Bars indicate standard error / A) Análisis no paramétrico de escalamiento multidimensional (nMDS) y análisis de similitudes (ANOSIM) que describen los ensambles de invertebrados de Discos Levantados (UH) y Discos No-levantados (NH). B) Número de especies. C) Índice de Shannon-Wiener correspondiente a la fauna asociada en los 2 tipos de discos de *L. spicata*. Barras indican error estándar**

knowledge, this is the first study showing effects of coastal uplift on the composition of invertebrate assemblages inhabiting kelp holdfasts. It is also the first to assess a rare primary succession occurring in a new intertidal belt.

The bivalve *P. purpuratus* and the anemone *P. clematis* contributed the most to the differences in the structure of the assemblages, mainly because these species were more abundant in UH. One possible reason is that the assemblages in NH were much younger than their UH counterpart and so the stage of succession may drive the differences (Smith *et al.* 1996). However, it has been observed that *P. purpuratus* is capable of withstanding long periods of desiccation and high salinity; moreover, they have been observed to occasionally occupy positions in the high intertidal, where seawater reach is only sporadic and limited (Vial *et al.* 1992). *Phymactis clematis* has also been found to be tolerant to long periods of desiccation and high salinity (Acuña & Zamponi 1995). Furthermore, and more importantly, *P. clematis* preferentially preys on mussels such as *Brachydontes rodriguezi* (Acuña & Zamponi 1996), which might

suggest predator-prey links between *P. clematis* and *P. purpuratus*. In the intertidal, *L. spicata* holdfasts provide refuge from wave action during high tide, and from desiccation during low tide due to water retention in its complex structure (Vásquez & Santelices 1984). Our data suggest that a drastic coastal uplift, produced by natural events such as the mega-earthquake in 2010, may benefit holdfast species with high tolerances to desiccation and high salinity, and that UH still provide suitable refuge for their survival and development.

The NH in the new intertidal belt had lower species richness in relation to the UH. These results are in agreement with the observations made by Vásquez & Santelices (1984) and Villouta & Santelices (1986), who found that new holdfasts contained lower species richness. Our data indicates that only one species, the polychaete worm Sabellariidae sp., had higher abundance in NH than in UH. Sabellariids are fast-growing, detritivorous and filter-feeding worms, which need high humidity and regular water exposure (see Knight-Jones 1981). Their high abundance in NH might suggest they are pioneer species of this primary succession.

This investigation provides new insights into the effects of natural events such as drastic coastal uplift in littoral communities. Our data suggest that holdfasts lifted within the ranges observed in the 2010 earthquake in 'La Desembocadura' can still remain as active 'micro-ecosystems'; however, the species benefiting will be those that can survive and withstand long periods of desiccation and high salinity. On the other hand, young holdfasts in the new intertidal belt provide a new refuge for invertebrate assemblages and provide the possibility for a rare primary succession. In Chile, coastal uplifts caused by telluric events can cover extended regions of coastline and might induce massive changes in the structure of assemblages that still remain unknown. In the future, replication of this study using greater sampling efforts and higher taxonomic resolution will help to better elucidate the macro-ecological implications of such rare events.

#### ACKNOWLEDGMENTS

We thank the contribution of two anonymous referees that helped improving this manuscript. This research was partially supported by FONDECYT Project 11110437 (CONICYT) to ECM. We also thank CONICYT Becas Chile Scholarship (72110557) for financial support for doctoral studies to CAS. We are grateful to Ricardo Jeldres, Ariel Cáceres, María Monsalvez and Solange Pacheco for their help in the field and laboratory analyses.

#### LITERATURE CITED

- Acuña FH & MO Zamponi. 1995.** Ecology of intertidal sea anemones, density, dispersion and autoecology of *Phymactis clematis* Dana, 1849 (Anthozoa, Actiniaria). *Ciencias Marinas* 21: 1-12.
- Acuña FH & MO Zamponi. 1996.** Trophic ecology of the intertidal sea anemones *Phymactis clematis* Dana, 1849, *Aulactinia marplatensis* (Zamponi, 1977) and *A. reynaudi* (Milne-Edwards, 1857) (Actiniaria: Actiniidae): Relationships between sea anemones and their prey. *Ciencias Marinas* 22: 397-413.
- Awata Y, S Toda, H Kaneda, T Azuma, H Horikawa, M Shishikura & T Echigo. 2008.** Coastal deformation associated with the 2007 NotoHanto earthquake, central Japan, estimated from uplifted and subsided intertidal organisms. *Earth Planets and Space* 60: 1059-1062.
- Bodin P & T Klinger. 1986.** Coastal uplift and mortality of intertidal organisms caused by the September 1985 Mexico earthquakes. *Science* 233: 1071-1073.
- Cancino JM & B Santelices. 1984.** Importancia ecológica de los discos de *Lessonia nigrescens* Bory (Phaeophyta) en Chile central. *Revista Chilena de Historia Natural* 57: 23-33.
- Castilla JC. 1988.** Earthquake-caused coastal uplift and its effects on rocky intertidal kelp communities. *Science* 242: 440-443.
- Castilla JC & D Oliva. 1990.** Ecological consequences of coseismic uplift on the intertidal kelp belts of *Lessonia nigrescens* in central Chile. *Estuarine Coastal and Shelf Science* 31: 45-56.
- Castilla JC, PH Manríquez & A Camaño. 2010.** Effects of rocky shore coseismic uplift and the 2010 Chilean mega-earthquake on intertidal biomarker species. *Marine Ecology Progress Series* 418: 17-23.
- Clarke KR. 1993.** Non-parametric multivariate analyses of changes in community structure. *Australian Journal of Ecology* 18: 117-143.
- Clarke KR, PJ Somerfield & MG Chapman. 2006.** On resemblance measures for ecological studies, including taxonomic dissimilarities and a zero-adjusted Bray-Curtis coefficient for denuded assemblages. *Journal of Experimental Marine Biology and Ecology* 330: 55-80.
- Fariás M, G Vargas, A Tassara, S Carretier, S Baize, D Melnick & K Bataille. 2010.** Land-Level changes produced by the Mw 8.8 2010 Chilean earthquake. *Science* 329: 916-916.
- González A, J Beltrán, L Hiriart-Bertrand, V Flores, B de Reviers, JA Correa & B Santelices. 2012.** Identification of cryptic species in the *Lessonia nigrescens* complex (Phaeophyceae, Laminariales). *Journal of Phycology* 48: 1153-1165.
- Jaramillo E, JE Dugan, DM Hubbard, D Melnick, M Manzano, C Duarte, C Campos & R Sanchez. 2012.** Ecological implications of extreme events: footprints of the 2010 earthquake along the Chilean coast. *PLoS ONE* 7: e35348.
- Jones DJ. 1972.** Changes in ecological balance of invertebrate communities in kelp holdfast habitats of some polluted North-Sea waters. *Helgoländer Wissenschaftliche Meeresuntersuchungen* 23: 248-260.
- Knight-Jones PE. 1981.** Behavior, setal inversion and phylogeny of Sabellida (Polychaeta). *Zoologica Scripta* 10: 183-202.
- Melnick D, M Cisternas, M Moreno & R Norambuena. 2012.** Estimating coseismic coastal uplift with an intertidal mussel: calibration for the 2010 Maule Chile earthquake (M-w=8.8). *Quaternary Science Reviews* 42: 29-42.
- Ojeda FP & B Santelices. 1984.** Ecological dominance of *Lessonia nigrescens* (Phaeophyta) in central Chile. *Marine Ecology Progress Series* 19: 83-91.

- Sáez CA, A Pérez-Matus, MG Lobos, D Oliva, JA Vásquez & M Bravo. 2012.** Environmental assessment in a shallow subtidal rocky habitat: Approach coupling chemical and ecological tools. *Chemistry and Ecology* 28: 1-15.
- Smith SDA. 2000.** Evaluating stress in rocky shore and shallow reef habitats using the macrofauna of kelp holdfasts. *Journal of Aquatic Ecosystem Stress and Recovery* 7: 259-272.
- Smith SDA & RD Simpson. 1992.** Monitoring the shallow Marine Biology: An Annual Review 45: 195-344.
- Tuya F, K Larsen & V Platt. 2011.** Patterns of abundance and assemblage structure of epifauna inhabiting two morphologically different kelp holdfasts. *Hydrobiologia* 658: 373-382.
- Vargas G, M Fariás, S Carretier, A Tassara, S Baize & D Melnick. 2011.** Coastal uplift and tsunami effects associated to the 2010 Mw8.8 Maule earthquake in Central Chile. *Andean Geology* 38: 219-238.
- Vásquez JA & B Santelices. 1984.** Comunidades de macroinvertebrados en discos de adhesión de *Lessonia nigrescens* en Chile central. *Revista Chilena de Historia Natural* 57: 131-154.
- Vial MV, RW Simpfendorfer, DA Lopez, ML Gonzalez & K Oelckers. 1992.** Metabolic responses of the intertidal mussel *Perumytilus purpuratus* (Lamarck) in emersion and immersion. *Journal of Experimental Marine Biology and Ecology* 159: 191-201.
- Villouta E & B Santelices. 1986.** *Lessonia trabeculata* sp. nov (Laminariales, Phaeophyta), a new kelp from Chile. *Phycologia* 25: 81-86.

---

Received 02 January 2013 and accepted 20 November 2013  
Associate Editor: Pilar Muñoz M.