



Revista Árvore

ISSN: 0100-6762

r.arvore@ufv.br

Universidade Federal de Viçosa
Brasil

Gallo, Ricardo; Xavier, Aloisio; Coelho de Moura, Luciana; de Almeida Oliveira, Brenner;
Rocha do Nascimento, Heloisa; Campos Otoni, Wagner

IBA AND MICROCUTTING COLLECTIONS IN THE MICROPROPAGATION OF
EUCALYPTUS SPP HYBRID CLONES.

Revista Árvore, vol. 41, núm. 6, 2017, pp. 1-13

Universidade Federal de Viçosa
Viçosa, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=48853637004>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

IBA AND MICROCUTTING COLLECTIONS IN THE MICROPROPAGATION OF *EUCALYPTUS* SPP HYBRID CLONES.¹

Ricardo Gallo^{2*}, Aloisio Xavier³, Luciana Coelho de Moura⁴, Brenner de Almeida Oliveira⁵, Heloisa Rocha do Nascimento⁶ and Wagner Campos Otoni⁷

¹ Received on 08.08.2014 accepted for publication on 26.09.2017.

² Universidade Federal de Viçosa, Programa de Pós-Graduação em Ciência Florestal, Viçosa, Minas Gerais, Brasil. E-mail: <gallo.florestal@gmail.com>.

³ Universidade Federal de Viçosa, Departamento de Engenharia Florestal, Viçosa, MG, Brasil. E-mail: <xavier@ufv.br>.

⁴ Universidade Federal de Viçosa, Doutorado em Ciência Florestal, Viçosa, MG, Brasil and <lucianacm2005@yahoo.com.br>.

⁵ Universidade Federal de Viçosa, Mestrado em Ciência Florestal, Viçosa, MG, Brasil. E-mail: <b.oliveiraeft@gmail.com>.

⁶ Universidade Federal do Rio Grande do Norte, Unidade Acadêmica Especializada em Ciências Agrária, Macaíba, Brasil. E-mail: <gualterges@yahoo.com.br>.

⁷ Universidade Federal de Viçosa, Programa de Pós-Graduação em Fitotecnia, Viçosa, MG, Brasil. E-mail: <helornasc@gmail.com>.

⁷ Universidade Federal de Viçosa, Departamento de Biologia Vegetal, Viçosa, MG, Brasil. E-mail: <wcotoni@gmail.com>.

*Corresponding author.

ABSTRACT – This study aims to evaluate the effect of IBA concentrations and microcuttings successive collections in the micropropagation of *Eucalyptus grandis* x *E. urophylla* and *Eucalyptus urophylla* x *E. globulus* clones. Clumps containing six to eight buds of clones established *in vitro* were transferred to a 250 mL glass flask in JADS semisolid medium. Successive collections were performed every 20 days for *Eucalyptus grandis* x *E. urophylla* clone and every 30 days for *Eucalyptus urophylla* x *E. globulus* clone. The following variables were evaluated under *in vitro* conditions: number of shoots > 0.5 cm, number of microcuttings > 2 cm, length of the longest microcutting, and shoots vigor. Under *ex vitro* conditions, in the greenhouse and shade house, the following variables were evaluated: seedling height, percentage of survival, stem diameter, percentage of root observed at the lower end of the tube, and seedling vigor. In full sun (*ex vitro*), the following variables were analyzed: seedling height, stem diameter, survival, number of roots, root volume, seedling vigor, and shoot and root dry matter. Good *in vitro* microcuttings productivity was observed over the successive collections. IBA levels were adjusted for each clone, ranging from 0.25 to 0.50 mg L⁻¹ for *Eucalyptus grandis* x *E. urophylla* clone, and from 0.75 to 1.0 mg L⁻¹ for *Eucalyptus urophylla* x *E. globulus* clone. IBA concentrations led to residual effects under *ex vitro* conditions, providing good rooting and survival for *Eucalyptus grandis* x *E. urophylla* and *Eucalyptus urophylla* x *E. globulus* clones at IBA concentrations between 0.25 and 0.50 mg L⁻¹ and between 0.50 and 1.0 mg L⁻¹, respectively.

Keywords: *Cloning; In vitro propagation; Vegetative propagation.*

AIB E COLETAS DE MICROESTACAS NA MICROPROPAGAÇÃO DE CLONES HÍBRIDOS DE EUCALYPTUS SPP.

RESUMO – O presente trabalho tem como objetivo avaliar o efeito de doses de AIB e de coletas sucessivas de microestacas na micropropagação de clones de *Eucalyptus grandis* x *E. urophylla* e *Eucalyptus urophylla* x *E. globulus*. Tufos contendo de seis a oito gemas dos clones *in vitro*, foram transferidos para frascos de vidro - 250 mL, em meio semissólido JADS, sendo feitas coletas a cada 20 dias para o clone *Eucalyptus grandis* x *E. urophylla* e a cada 30 dias para o *Eucalyptus urophylla* x *E. globulus*. Nas condições *in vitro*, foram avaliados: número de brotos > 0,5 cm, número de microestacas > 2 cm, comprimento da maior microestaca e vigor dos brotos. Nas condições *ex vitro*, na casa de vegetação e casa de sombra, foram avaliados: altura da muda, sobrevivência, diâmetro do colo, porcentagem de emissão de raiz na extremidade inferior do tubete, vigor das mudas. Em pleno sol (*ex vitro*), foram avaliados: altura da muda, sobrevivência, diâmetro do



colo, número de raízes, volume de raiz, vigor das mudas, e matéria seca da parte aérea e da raiz. Observou-se boa produtividade de microestacas *in vitro* no decorrer das coletas, sendo ajustados níveis de AIB para cada clone, variando de 0,25 a 0,50 mg L⁻¹ para o clone *Eucalyptus grandis* x *E. urophylla* e de 0,75 a 1,0 mg L⁻¹ para o clone *Eucalyptus urophylla* x *E. globulus*. Nas condições *ex vitro*, a dosagem de AIB apresentou efeito residual, proporcionando bom enraizamento e sobrevivência, entre 0,25 a 0,50 mg L⁻¹ de AIB para o clone *Eucalyptus grandis* x *E. urophylla* e 0,50 a 1,0 mg L⁻¹ para o clone *Eucalyptus urophylla* x *E. globulus*.

Palavras-chave: Clonagem; Propagação *in vitro*; Propagação vegetativa.

1. INTRODUCTION

The successful productivity of *Eucalyptus* plantations in Brazil is a result of the combination of several factors, such as the well-established clonal forestry programs, the constant development of genetic improvement strategies, the production of specific hybrids, the selection of elite clones, and the improvement of clonal propagation technologies (Ferrari et al., 2004; Xavier et al., 2013).

Micropropagation is a clonal technique that produces microcuttings and provides countless advantages to the production process of *Eucalyptus* seedlings, such as faster mass propagation of clones; greater nutritional, environmental, and phytosanitary control; and longer storage periods; besides retention of hybrid vigor and the possibility of transport over long distances without damaging the material (Bisht et al., 1999; Xavier et al., 2013).

In addition, microcuttings production by micropropagation improves the cloning process due to the rejuvenation and reinvigoration technique used as a mechanism to overcome rooting problems of the cuttings, which is mainly observed in the rescue of adult trees (Joshi et al., 2003; Xavier et al., 2013).

Specific innovations have been proposed for the application of large-scale micropropagation technologies in *Eucalyptus* (Xavier et al., 2013), such as the manipulation of the *in vitro* atmosphere and/or environment (e.g., photoautotrophic propagation) (Kozai, 2010); the use of temporary immersion and liquid medium bioreactors (Oliveira et al., 2011, 2014); the replacement of the semisolid medium with alternative substrates (Kirdmanee, et al., 1995); the automation and mechanization of systems operations (Penchel et al., 2007); and the use of *in vitro* microclonal hedge.

Several factor can influence the productivity of the traditional miniclinal hedge, such as climatic variables, the genetic material, the production system (Cunha et al., 2005), the mineral nutrition (Cunha et al., 2009a),

the management of miniclinal hedge seedlings (Mafia et al., 2005), among others. Cunha et al. (2009b) concluded that the increase in temperature favors the production of minicuttings in eucalyptus miniclinal hedge, regardless of the miniclinal hedge type.

The major advantage of transferring the miniclinal hedge from *ex vitro* to *in vitro* environment is the greater control of climatic variables, such as low temperatures, besides the *in vitro* elimination of pests and diseases. Thus, the existence of efficient micropropagation protocols for *Eucalyptus* would enable the use of *in vitro* microclonal hedge and make the microcuttings available to the forest industry (Xavier et al., 2013).

The knowledge of the use of auxins is fundamental for the establishment of efficient protocols. This fact is because these growth regulators are directly related to several physiological processes, such as activation of cambium cells, apical dominance, plant growth promotion, lateral and adventitious root formation, foliar abscission, floral buds and fruits development, and induction of vascular differentiation (Bresinsky et al., 2012; Kerbaudy, 2012; Taiz and Zeiger, 2013).

IBA (3-indole butyric acid) is widely used in *in vitro* and *ex vitro* propagation processes as this auxin does not damage explants. However, the concentration to be applied must be tested for each species and propagule (Titon et al., 2003a; Iacona and Muleo, 2010).

Studies have reported the need for using IBA for rooting in the micropropagation of several morphological processes, such as: rooting in *Oncidium baueri* (Camargo et al., 2015); increase in number and length of roots in nodal segments of *Campomanesia adamantium* (Rossato et al., 2015); *in vitro* elongation of hybrid clones of *Eucalyptus globulus* (Oliveira et al., 2016); and *ex vitro* acclimatization and rooting of *Ilex paraguariensis* microcuttings (Tronco et al., 2015).

The *in vitro* environment enables the production of microcuttings. The knowledge of the number of collections or production time in this system is crucial

for the establishment of an *in vitro* microclonal hedge. The information on the the number of collections or production can provide the amount of microcutting that a clump of shoot can produce in a given time and and culture medium, providing practical benefits to the implantation of *in vitro* microclonal hedge.

The objective of the present work was to evaluate the effect concentrations of IBA and successive collections of microcutting on clumps of shoots in multiplication in the micropropagation of clones of *Eucalyptus grandis* x *E. urophylla* and *Eucalyptus urophylla* x *E. globulus*.

2. MATERIAL AND METHODS

2.1 Plant material and *in vitro* cultivation conditions

The experiments were conducted at the Tissue Culture Laboratory II of the Institute of Applied Biotechnology for Agriculture (BIOAGRO), at the Federal University of Viçosa, located in the municipality of Viçosa/MG.

The material used in this work resulted from the multiplication by micropropagation of *Eucalyptus grandis* x *E. urophylla* and *Eucalyptus urophylla* x *E. globulus* clones, with 25 and 72 subcultures, respectively. Shoots originated from clumps containing six to eight differentiated buds were cultivated in test tubes containing 10 mL JADS culture medium (Correia et al., 1995), added with 30 g L⁻¹ of sucrose (VetecTM), 100 mg L⁻¹ of myo-inositol (SigmaTM), 800 mg L⁻¹ of PVP-30 (polyvinylpyrrolidone - VetecTM), 0.5 mg L⁻¹ of BAP (6 - benzylaminopurine – SigmaTM), 0.01 mg L⁻¹ of NAA (naphthaleneacetic acid – SigmaTM), and 7 g L⁻¹ of agar (MerckTM).

Subsequently, clumps were transferred to test tubes with JADS culture medium added with 0.3 mg L⁻¹ of BAP. The culture medium was adjusted to pH 5.8 and autoclaved in 1.5 atm pressure, at 121 °C, for 20 minutes.

Cultures were maintained in a growth room at 25 ± 2 °C, for 16 h photoperiod, and 33 μmol m⁻²s⁻¹ irradiances (quantified by the LI-CORTM radiometer, LI-250A Light Meter), provided by two tubular fluorescent lamps (Special Daylight, 40 W, OsramTM, Brazil).

At 30 days after incubation in the culture medium, with BAP reduction, clumps of each clone were

transferred to 250 mL glass flasks containing 40 mL of JADS culture medium, added with 30 g L⁻¹ of sucrose, 100 mg L⁻¹ of myo-inositol, 800 mg L⁻¹ of PVP-30, 7 g L⁻¹ of agar, 0.05 mg L⁻¹ of IBA, varying the concentrations of indole 3 butyric acid (IBA) (SigmaTM) between 0.0; 0.25; 0.50 and 1.00 mg L⁻¹.

In each flask, four clumps containing six to eight shoots were inoculated at the four IBA concentrations for each clone and maintained in a growth room with the same characteristics previously described. The experiment consisted of a completely randomized design, in a 5x4 factorial scheme, with five microcutting collections and four IBA concentrations (0.0; 0.25; 0.50; and 1.00 mg L⁻¹), with four replications; each plot contained four clumps of shoots.

2.2 Microcuttings *ex vitro* rooting

Experiments were conducted at the Research Center of the Forest Engineering Department of UFV.

Microcuttings > 2 cm from the clones and from treatments with IBA concentration under *in vitro* conditions were placed in Petri dishes containing two sheets of water-moistened filter paper and subsequently transferred to the nursery.

Microcuttings of each treatment (IBA concentrations) and clones (*Eucalyptus grandis* x *E. urophylla* and *Eucalyptus urophylla* x *E. globulus*) were planted in 55 cm³ plastic tubes containing commercial substrate (Tropstrato Vida VerdeTM) and vermiculite (medium particle size) at a ratio of 1:1, added with 5 kg m⁻³ of simple superphosphate (HeringerTM), and placed in a greenhouse (at 20-30°C and relative humidity e" 80%).

On the first ten days in the greenhouse, microcuttings were covered with Aluminet (50%), at a distance of 20 cm from the tubes. At 20 days after greenhouse cultivation, microcuttings were transferred to a shade house, where they received topdressing by applying 2 mL of monoammonium phosphate (MAP) per seedling, at a concentration of 2 g L⁻¹. Seedlings were maintained in the shade house (50%) for ten days and then transferred to full sun, when they received 5 mL of the NPK formulation 20-5-20 (HeringerTM) per seedling, at a concentration of 6 g L⁻¹, remaining under this condition for another 30 days.

The experiment consisted of a completely randomized block design, with four IBA concentrations

(0.0, 0.25, 0.50, and 1.00 mg L⁻¹) from the *in vitro* applications, in five blocks (collections), with a variable number of microcuttings, based on the *in vitro* production of each treatment.

2.3 Experimental evaluations and data analysis

Growth characteristics evaluation and microcuttings collection occurred every 20 days of *in vitro* cultivation for the *Eucalyptus grandis* x *E. urophylla* clone and every 30 days of *in vitro* cultivation for the *Eucalyptus urophylla* x *E. globulus* clone. Afterward, clumps of shoots were transferred to a new culture medium containing the same treatments. For both clones, five collections were performed, totaling 100 days of cultivation for *Eucalyptus grandis* x *E. urophylla* clone and 150 days for *Eucalyptus urophylla* x *E. globulus* clone.

Number of shoots > 0.5 cm per clump of shoots (NS > 0.5 cm), number of microcuttings > 2 cm (NM > 2 cm), length of the longest microcutting (cm) (LLM), and shoot vigor (SV) [based on a scale of grades ranging from 1 (Low), 2 (Intermediate), and 3 (High)] were analyzed on the day of each *in vitro* collection.

Under the *ex vitro* condition, at the moment of transfer from the greenhouse to the shade house, seedling height (H, cm), stem diameter (SD, mm), percentage of survival (PS, %), percentage of root observed at the lower end of the tube (PRT, %), and seedling vigor (SEV) [based on a scale of grades ranging from 1 (Low), 2 (Intermediate), and 3 (High)] were evaluated. After 30 days in full sun, seedling height (H, cm), stem diameter (SD, mm), percentage of survival (PS, %), number of roots (NR), root volume (RV, cm³), and seedling vigor (SEV) [based on a grade scale ranging from 1 (Low), 2 (Intermediate), and 3 (High)], shoot dry matter (SDM, g), and root dry matter (RDM, g) were evaluated.

Data were analyzed in the R software, version 3.2.4 (R Core Team, 2016), with the aid of the ExpDes package, version 1.1.2 (Ferreira et al., 2013). Percentage of survival and seedling height under *ex vitro* conditions did not present a normal distribution by the Shapiro-Wilk test at 5% significance, and thus were transformed into $\arcsen \sqrt{x}$ and , respectively. Regression equations were generated for the variables that were significant by the analysis of variance.

3. RESULTS

3.1 *Eucalyptus grandis* x *E. urophylla* clone

Based on the results obtained in the *in vitro* evaluations, number of shoots > 0.5 cm (Figure 1A) showed a quadratic trend with the number of collections. An increase in the number of shoots was observed from the second collection, followed by stabilization. Number of shoots > 0.5 cm presented the quadratic trend (Figure 1E), with a decrease in production at the highest IBA concentration.

The production projection of microcutting > 2 cm (Figure 1B) tended to increase linearly. However, the maximum microcutting production occurred at the IBA concentration of 0.44 mg L⁻¹ (Figure 1F).

The longest microcuttings were formed in the first and last collections (Figure 1C), increasing size from the fourth to the fifth collection. Conversely, shoot vigor over the collection presented a linear decrease (Figure 1D). The use of IBA concentrations for these variables had the same trend as that observed for the other evaluated characteristics, being the point that provides the longest microcutting, estimated in 0.54 mg L⁻¹ of IBA (Figure 1G). For shoot vigor, this point was estimated to be 0.45 mg L⁻¹ of IBA (Figure 1H).

The optimum range of IBA concentration for better results for shoot and microcutting production, as well as for size and vigor is from 0.25 to 0.50 mg L⁻¹.

Ex vitro evaluations at the moment of transfer from the greenhouse to the shade house had the same results as those reported for the *in vitro* environment. The characteristics treated with IBA presented quadratic results with *in vitro* IBA concentrations of 0.25 mg L⁻¹ (Figure 2B), and higher percentage of root observed at the lower end of the tube (Figure 2C).

At the moment of transfer from the greenhouse to the shade house, IBA concentrations did not affect the characteristics evaluated in the seedlings. However, in the last evaluation in full sun, treatments with IBA concentrations applied *in vitro* significantly influenced percentage of survival (PS, %), number of roots (NR), shoot dry matter (SDM, g), root dry matter (RDM, g), and root volume (RV, cm³) (Figure 2D-H).

RV showed a decreasing linear trend in function of the *in vitro* IBA treatments. RV reduced at the highest IBA concentration (1.0 mg L⁻¹) (Figure 2D). The highest




Figure 1 - Characteristics observed in the *in vitro* cultivation of *Eucalyptus grandis* x *E. urophylla* clone in function of successive collections. (A) Number of shoots > 0.5 cm; (B) Number of micro-cuttings > 2 cm; (C) Length of the longest micro-cutting (cm); and (D) Shoot vigor; and in function of IBA concentrations (0.0, 0.25, 0.5 and 1.0 mg L⁻¹). (E) Number of shoots > 0.5 cm; (F) Number of micro-cuttings > 2 cm; (G) Length of the longest micro-cutting (cm); and (H) Shoot vigor.

Figura 1 - Características observadas no cultivo *in vitro* do clone *Eucalyptus grandis* x *E. urophylla* em função das coletas sucessivas. (A) Número de brotos > 0,5 cm; (B) Número de microestacas > 2 cm; (C) Comprimento da maior microestaca (cm); e (D) Vigor dos brotos; e em função das doses de AIB (0,0; 0,25; 0,5 e 1,0 mg L⁻¹). (E) Número de brotos > 0,5 cm; (F) Número de microestacas > 2 cm; (G) Comprimento da maior microestaca (cm); e (H) Vigor dos brotos.

values for PS (Figure 2E), NR (Figure 2F), and for SDM (Figure 2G) and RDM (Figure 2H) were observed at intermediate IBA concentrations (0.25 and 0.5 mg L⁻¹), with a quadratic response at the end of the full sun stage (60 days).

The use of IBA under *in vitro* conditions also provided residual effects in the *ex vitro* conditions, directly influencing seedlings quality. Seedling height, seedling vigor, and percentage of root observed at the lower end of the tube were significant at the moment of transfer from the greenhouse to the shade house. However, these characteristics are statistically similar at the other stages (shade house and full sun). Stem

diameter did not present statistical difference with treatments application in any of the *ex vitro* environment.

In the greenhouse, *Eucalyptus grandis* x *E. urophylla* clone had mean value of 85.2% microcuttings survival without the influence of IBA concentrations applied under *in vitro* conditions. At the moment of transfer from the greenhouse to the shade house, and without the influence of the *in vitro* concentrations, percentage of survival decreased (78.45%). Conversely, after 60 days in full sun, microcuttings showed 77.8% survival under the influence of the IBA concentrations applied in the *in vitro* environment. The point of maximum survival was estimated using 0.52 mg L⁻¹ IBA in the culture medium, under *in vitro* condition.




Figure 2 - Characteristics observed at the moment of transfer from the greenhouse to the shade house (A) Seedling height (cm); (B) Seedling vigor; and (C) Roots observed at the lower end of the tube (%); and characteristics observed in the seedling in full sun (D) Root volume (cm³); (E) Percentage of survival (%); (F) Number of roots; (G) Shoot dry matter (g); and (H) Root dry matter (g). For *E. urophylla* x *Eucalyptus grandis* clone in function of IBA concentrations (0.0, 0.25, 0.5 and 1.0 mg L⁻¹).

Figura 2 - Características observadas na saída da casa de vegetação (A) Altura da muda (cm); (B) Vigor das mudas; e (C) Raiz observada na extremidade inferior do tubete (%); e características observadas em pleno sol em mudas (D) Volume de raiz (cm³); (E) Porcentagem de sobrevivência (%); (F) Número de raiz; (G) Massa seca da parte aérea (g); e (H) Massa seca da raiz (g). Para o clone *Eucalyptus grandis* x *E. urophylla* em função das doses de AIB (0,0; 0,25; 0,5 e 1,0 mg L⁻¹).

3.2 *Eucalyptus urophylla* x *E. globulus* clone

The analysis of variance for *Eucalyptus urophylla* x *E. globulus* clones revealed interaction in the collection. IBA concentrations were significant for number of shoots > 0.5 cm (NS > 0.5 cm) [with significance ($p = 0.0102$)] and shoot vigor ($p < 0.01$). For the number of microcuttings > 2 cm (NM > 2 cm), the variables the variables were independent ($p < 0.01$). For length of the longest microcuttings (LLM), only the IBA concentrations were significant ($p < 0.01$).

IBA concentrations for NS > 0.5 cm in function of the collections (Figure 3A) showed a second-degree

polynomial behavior. From the first collection, all IBA concentrations followed an increasing trend; however, from the third to the fourth collection, the IBA concentrations of 0.0; 0.25 and 0.5 mg L⁻¹ tended to reduce shoot production. For the highest IBA concentration (1.0 mg L⁻¹), this trend only occurred from the fourth to the fifth collection.


For shoot vigor (SV), the response was in function of the interaction between treatments, showing a significant ($p < 0.01$) decreasing linear behavior for IBA concentrations of 0.25; 0.50 and 1.00 mg L⁻¹ in function

of the collections (Figure 3B), with an evident decrease in vigor with the advance of the monthly collections.

Production of NM > 2 cm increased in function of the successive collections (Figure 3C), and the IBA concentration of 0.65 mg L⁻¹ was the optimum point

(Figure 3D). LLM presented increasing linear response in function of the IBA concentrations (Figure 3E).

Results obtained under *ex vitro* conditions indicated the effect of IBA concentrations applied in the *in vitro* environment, with differences in the evaluations performed in greenhouse, shade house, and in full sun.



(A) NS > 0.5 cm

- 0.0 mg L⁻¹ $\hat{y} = 1.7 + 8.3857^{(p<0.01)} \times \text{Collections} - 1.2143^{(p<0.01)} \times \text{Collections}^2$ $R^2 = 0.82$
- △ 0.25 mg L⁻¹ $\hat{y} = 2.45 + 7.8643^{(p<0.01)} \times \text{Collections} - 1.0357^{(p<0.01)} \times \text{Collections}^2$ $R^2 = 0.87$
- 0.50 mg L⁻¹ $\hat{y} = 3.55 + 7.6214^{(p<0.01)} \times \text{Collections} - 0.9286^{(p<0.01)} \times \text{Collections}^2$ $R^2 = 0.92$
- ◇ 1.0 mg L⁻¹ $\hat{y} = 7.15 + 5.5464^{(p<0.01)} \times \text{Collections} - 0.5536^{(p<0.01)} \times \text{Collections}^2$ $R^2 = 0.82$

(B) SV

- △ 0.25 mg L⁻¹ $\hat{y} = 2.3 - 0.1875^{(p<0.01)} \times \text{Collections}$ $r^2 = 0.48$
- 0.50 mg L⁻¹ $\hat{y} = 2.225 - 0.15^{(p<0.01)} \times \text{Collections}$ $r^2 = 0.65$
- ◇ 1.0 mg L⁻¹ $\hat{y} = 2.6188 - 0.3188^{(p<0.01)} \times \text{Collections}$ $r^2 = 0.93$

- ▲ (C) NM > 2.0 cm $\hat{y} = 2.9375 - 0.6607^{(p=0.186)} \times \text{Collections} + 0.2142^{(p=0.01)} \times \text{Collections}^2$ $R^2 = 0.89$

- (D) NM > 2.0 cm $\hat{y} = 1.3668 + 10.4064^{(p<0.01)} \times \text{IBA} - 7.9454^{(p<0.01)} \times \text{IBA}^2$ $R^2 = 0.91$

- (E) LLM (cm) $\hat{y} = 2.4230 + 3.2235^{(p<0.01)} \times \text{IBA}$ $r^2 = 0.93$

Figure 3 - Characteristics observed in *in vitro* cultivation of *Eucalyptus urophylla* x *E. globulus* clone in function of collections (every 30 days) and IBA concentrations (0.0, 0.25, 0.5 and 1.0 mg L⁻¹). (A) Number of shoots > 0.5 cm; (B) Shoot vigor; (C) Number of micro-cuttings > 2 cm in function of the successive collections (D) and in function of the IBA concentrations; and (E) Length of the longest micro-cutting in function of the IBA concentrations.

Figura 3 - Características observadas no cultivo *in vitro* do clone de *Eucalyptus urophylla* x *E. globulus* em função das coletas (a cada 30 dias) e doses de AIB (0,0; 0,25; 0,5 e 1,0 mg L⁻¹). (A) Número de brotos > 0,5 cm; (B) Vigor dos brotos; (C) Número de microestacas > 2 cm em função das coletas sucessivas (D) e em função das doses de AIB; e (E) Comprimento da maior microestaca em função das doses de AIB.

Seedling height at the moment of transfer from the greenhouse to the shade house had an increasing linear trend, and the highest IBA concentration (1.0 mg L⁻¹) was associated with higher mean values for seedling height (Figure 4A). However, under shade house and in full sun conditions, seedling height

increased with the application of 0.91 and 0.80 mg L⁻¹ of IBA, respectively.

Seedlings had a greater stem diameter in the *ex vitro* stages - greenhouse, shade house, and full sun - when the concentrations of 0.5 and 0.75 mg L⁻¹ were used in the *in vitro* condition (Figure 4B). However,




Figure 4 - Characteristics observed at the moment of transfer from the greenhouse (GH) to the shade house (SH) and in full sun (FS) in seedlings of *Eucalyptus urophylla* x *E. globulus* clones in function of the IBA concentrations (0.0, 0.25, 0.5 and 1.0 mg L⁻¹). (A) Seedling height (cm); (B) stem diameter (mm); (C) Seedling vigor; (D) Percentage of survival (%); and (E) Root observed at the lower end of the tube (%).

Figura 4 - Características observadas na saída da Casa de vegetação (GH); Casa de sombra (SH) e Pleno sol (FS) em mudas do clone de *Eucalyptus urophylla* x *E. globulus* em função das doses de AIB (0,0; 0,25; 0,5 e 1,0 mg L⁻¹). (A) Altura da muda (cm); (B) Diâmetro do colo (mm); (C) Vigor das mudas; (D) Porcentagem de sobrevivência (%); e (E) Raiz observada na extremidade inferior do tubete (%).

for seedling vigor (SEV), higher values were observed at higher IBA concentrations used in the *in vitro* condition (Figure 4C).

The percentage of survival presented a quadratic trend for all the *ex vitro* stages, and the highest percentage was observed with the *in vitro* application of IBA concentration between 0.81 and 0.94 mg L⁻¹ (Figure 4D). The percentage of root observed at the lower end of the tube presented a quadratic response to the regression equation. Consequently, this variable presented the highest values when the application of IBA ranged between 0.67 and 0.65 mg L⁻¹ (Figure 4E) in the greenhouse and shade house, respectively.

NR (Figure 5A) and VR (Figure 5B) evaluated at the end of the full sun stage presented a quadratic trend in function of the IBA concentrations, with similar maximum points: for NR, the estimated point was 0.67 mg L⁻¹ of IBA; and for the VR, the estimated point was 0.70 mg L⁻¹ of IBA.

SDM (Figure 5C) and RDM (Figure 5D) showed second-degree polynomial behavior in function of the IBA concentrations, with different maximum point of matter production: 0.75 mg L⁻¹ for SDM and 0.92 mg L⁻¹ for RDM.

Percentage of survival during microcuttings acclimatization at the moment of transfer from the greenhouse to the shade house presented a mean of 74.7%, being influenced by the IBA concentrations applied in the *in vitro* environment. The highest percentage of survival was estimated in 96.3% with the IBA concentration of 0.81 mg L⁻¹. At the moment of transfer from the shade house to the full sun, microcuttings survival (64.2%) continued to be influenced by the concentrations that had been applied in the *in vitro* environment, with an optimal IBA concentration estimated at 0.70 mg L⁻¹, which resulted in 84.0% microcuttings survival. In full sun, the mean value of microcuttings survival was 62.1%, still influenced by residual IBA, presenting a maximum survival point estimated at 0.89 mg L⁻¹ of IBA, leading to 84.0% microcuttings survival.

4.DISCUSSION

Under *in vitro* experimental conditions, the number of shoots of the *Eucalyptus grandis* x *E. urophylla* clone increased. Production decreased from the fourth collection, which was also observed in *Eucalyptus*




Figure 5 - Characteristics observed in seedlings of *Eucalyptus urophylla* x *E. globulus* clone in full sun in function of IBA concentrations (0.0, 0.25, 0.5, and 1.0 mg L⁻¹). (A) Number of roots; (B) Root volume (cm³); (C) Shoot dry matter (g); and (D) Root dry matter (g).

Figura 5 - Características observadas em pleno sol em mudas do clone de *Eucalyptus urophylla* x *E. globulus* em função das doses de AIB (0,0; 0,25; 0,5 e 1,0 mg L⁻¹). (A) Número de raiz; (B) Volume de raiz (cm³); (C) massa seca da parte aérea (g); e (D) massa seca da raiz (g).

urophylla x *E. globulus* clone, regardless of the IBA concentration. Oliveira (2016) and Gómez et al. (2007) state that the variation of the multiplication rate is related to the *in vitro* conditions and consequently to the excessive exposure to growth regulators in the subcultures. It is also related to the balance of cytokinins and auxins in the culture medium since this balance varies depending on the species and explant type.

For all *in vitro* variables, an increasing trend was observed until the IBA concentration of 0.5 mg L⁻¹. A decrease was observed from this value for *Eucalyptus grandis* x *E. urophylla* clone. This trend is explained for this is a rejuvenated material, in which the internal hormonal balance favors regeneration. Negative responses to additional hormonal applications may occur under certain conditions. Similar results were

reported by Bezerra et al. (2014) for *Mimosa caesalpinhiifolia* treated with BAP concentrations in *in vitro* multiplication. According to Hartmann et al. (2011), excessive concentrations of auxin and the exposure time to this growth regulator may inhibit shoots and roots development, causing yellowing and leaf fall, necrosis, and even plant death.

In the present work, the use of growth regulators was essential to achieve the desired multiplication rates. Auxin, such as IBA, and cytokinins, such as BAP, are used in several combinations for morphogenic control during adventitious shoots multiplication in *Eucalyptus* (Del Ponte et al., 2001; Dutra et al., 2009). Similar results were obtained using IBA in shoots of *Eucalyptus globulus* ssp. *Maidenii*, with a higher multiplication rate in culture medium containing 0.2 mg L⁻¹ BAP + 0.02 mg L⁻¹ IBA (Sotelo and Monza, 2007).

Microcuttings production increased over the collections. No studies have reported on *in vitro* microcuttings production or elongated shoots in successive collections. However, this condition is analogous to the rejuvenation observed in plant materials that undergo several *in vitro* subcultures (Xavier et al., 2013). These are relevant results regarding the formation of *in vitro* microclonal hedge, and work as preliminary study for a mass and economic production of *Eucalyptus* microcuttings.

In addition, in the case of an *ex vitro* environment, microcuttings produced from microstump presents a cyclical trend and may be related to the temporary depletion of the latter, leading to lower productions (Titon et al., 2003b), besides the temperature change (Xavier and Comério, 1996).

The absence of IBA for both clones reduced growth, corroborating the work of Oliveira (2016), in which the absence of IBA provided lower growth rates for *Eucalyptus urophylla* x *E. globulus* clone and *Eucalyptus grandis* x *E. globulus* clone. The best IBA concentration for *Eucalyptus globulus* ssp. *maidenii* elongation was 0.50 mg L⁻¹, resulting in four elongated shoots per explant (> 2 cm) (Sotelo and Monza, 2007), which is in agreement with the present results. Different shoot length responses were observed by Oliveira (2016), with optimal IBA concentrations ranging from 0.25 mg L⁻¹ to 1.0 mg L⁻¹.

For both clones, shoots vigor reduced over the collections. This may have occurred due to the increase

of the callus at the base of the *in vitro* explant. Callus is defined as a mass of disjointed and disorganized cells (Hartmann et al., 2011), which may have hindered the water and nutrients transport to the clump and reduced shoots vigor. Wendling et al. (2003) evaluated subculture of ministump and concluded that the subcultures did not increase ministumps vigor, evidencing a certain reduction trend with the increase in the subcultures.

Shoot vigor results in function of IBA concentrations were very different between the clones, evidencing genetic control for this variable. For *Eucalyptus grandis* x *E. urophylla* clone, the material that received the treatments with the intermediate concentration presented greater vigor. For *Eucalyptus urophylla* x *E. globulus* clone, the material that received the highest concentrations had the highest vigor; however, vigor decreased with the increase in the number of collections. Oliveira (2016) confirmed that the highest vigor for the tested *Eucalyptus* hybrid clones varied with the IBA concentrations between 0.36 and 0.80 mg L⁻¹ in MS medium. In JADS culture medium, the maximum vigor points were higher at the IBA concentration of 0.60 mg L⁻¹.

Evaluations under *ex vitro* environment conditions for *Eucalyptus grandis* x *E. urophylla* and *Eucalyptus urophylla* x *E. globulus* clones showed that treatments containing IBA in the culture medium influenced microcuttings quality. For the *Eucalyptus grandis* x *E. urophylla* clone, the intermediate IBA concentrations applied in the *in vitro* environment provided the best results. For *Eucalyptus urophylla* x *E. globulus* clones, higher IBA concentrations presented satisfactory results; however, at some concentrations, the percentage of microcuttings rooting decreased. In some cases, the plant response to the endogenous or applied auxin may vary depending on the tissue and the IBA concentration already present in the propagule. In addition, auxin may increase the rhizogenic response to a certain point, depending on the concentration, after which inhibitory effect occurs (Hartmann et al., 2011).

For the present study, high mean percentage of survival was observed for both clones in the nursery, which makes feasible the proposed microcuttings technique. Studies with *Bowdichia virgilioides* (Moura et al., 2012), *Aegiphila verticillata* (Almeida et al., 2015), and *Azadirachta indica* (Houllou et al., 2015)

have also presented high survival rates for *in vitro* microcuttings.

Further studies should be carried out to define the set of factors that influence micropropagation/microcuttings of these *Eucalyptus* spp. clones or other genetic materials and hybrids of commercial interest, aiming to facilitate *in vitro* miniclinal hedges for seedlings production by the forestry industry.

5. CONCLUSIONS

For the clones evaluated in this study, results revealed that 1) the production of microcuttings for *Eucalyptus grandis* x *E. urophylla* and *Eucalyptus urophylla* x *E. globulus* clones increased over the five *in vitro* collections; 2) the IBA concentrations for each clone were adjusted, ranging from 0.25 to 0.50 mg L⁻¹ for *Eucalyptus grandis* x *E. urophylla* clone and from 0.75 to 1.0 mg L⁻¹ for *Eucalyptus urophylla* x *E. globulus* clone; 3) the *ex vitro* conditions were influenced by the residual effect of the *in vitro* IBA concentrations; 4) IBA concentration between 0.25 and 0.50 mg L⁻¹ and between 0.50 to 1.0 mg L⁻¹ provided good rooting for *Eucalyptus grandis* x *E. urophylla* clone and *Eucalyptus urophylla* x *E. globulus* clone, respectively.

6. ACKNOWLEDGMENTS

The authors thank the Brazilian sponsoring agencies CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico, Brazil), FAPEMIG (Fundação de Amparo à Pesquisa do Estado de Minas Gerais), and CAPES (Coordenação de Aperfeiçoamento de Pessoal de Ensino Superior) for the financial support; and BIOAGRO (Instituto de Biotecnologia Aplicada à Agropecuária) for the laboratory infrastructure.

7. REFERENCES

- Almeida LMS, Morais LE, Resende CF, Braga VF, Pereira PF, Silva RAC et al. Micropropagation and acclimatization of *Aegiphila verticillata* Vell.: na endangered woody species. *Revista Árvore*. 2015;39:305-14.
- Bezerra RMF, Aloufa MAI, Freire FAM, Santos DD. Efeito de 6-benzilaminopurina sobre a propagação *in vitro* de *Mimosa caesalpiniiifolia* Benth. (Fabaceae). *Revista Árvore*. 2014;38:771-8.
- Bisht P, Sharma VK, Joshi I, Kapoor ML. Micropropagation of newly produced F1 hybrid of *Eucalyptus* (*E. tereticornis* Sm. x *E. camaldulensis* Dehn. Southern Form). *Silvae Genetica*. 1999;48:104-8.
- Bresinsky A, Körner C, Kadereit JW, Neuhaus G, Sonnenwald U. *Tratado de botânica de Strasburger*. 36ª ed. Porto Alegre: Artmed; 2012. 1166p.
- Camargo SS, Rodrigues DB, Rodrigues CM, Assis AM, Faria RT, Schuch MW. Fitorreguladores e espectros de luz na micropropagação de *Oncidium baueri* Lindl. *Ciência Rural*. 2015;45(11):2007-12.
- Correia D, Gonçalves AN, Couto HZ, Ribeiro MC. Efeito do meio de cultura líquido e sólido no desenvolvimento de gemas de *Eucalyptus grandis* x *Eucalyptus urophylla* na multiplicação *in vitro*. *IPEF*. 1995(48/49):107-16.
- Cunha ACMCM, Paiva HN, Leite HG, Barros NF, Leite FP. Influência do estado nutricional de minicepas no enraizamento de miniestacas de eucalipto. *Revista Árvore*. 2009a;33:607-15.
- Cunha ACMCM, Paiva HN, Leite HG, Barros NF, Leite FP. Relações entre variáveis climáticas com produção e enraizamento de miniestacas de eucalipto. *Revista Árvore*. 2009b;33:195-203.
- Cunha ACMCM, Wendling I, Souza Júnior L. Produtividade e sobrevivência de minicepas de *Eucalyptus benthamii* Maiden et Cabbage em sistema de hidroponia e em tubete. *Ciência Florestal*. 2005;15:307-10.
- Del Ponte EM, Mattei VL, Peters JA, Assis TF. Multiplicação e enraizamento *in vitro* de *Eucalyptus globulus* subsp. *Globulus* Labill. *Revista Árvore*. 2001;25:1-8.
- Dutra LF, Wendling I, Brondani GE. A micropropagação de eucalipto. *Pesquisa Florestal Brasileira*. 2009;58:49-59.
- Ferrari MP, Grossi F, Wendling I. Propagação vegetativa de espécies florestais. Colombo: Embrapa Florestas; 2004.
- Ferreira EB, Cavalcanti PP, Nogueira DA. ExpDes: Experimental designs package. R

package version 1.1.2. 2013.

Gómez C, Ríosa D, Sánchez-Olate M. Efecto del subcultivo sucesivo sobre la caulogénesis adventicia de *Eucalyptus globulus*. Bosque. 2007;28:13-7.

Hartmann HT, Kester DE, Daves Jr FD, Geneve RL. Plant propagation: principles and practices. 8th ed. New Jersey: Prentice-Hall; 2011.

Houllou LM, Souza RA, Santos ECP, Silva JJP, Barbosa MR, Sauvé JPG et al. Clonal propagation of nem (*Azadirachta indica* A. Juss.) via direct and indirect *in vitro* regeneration. Revista Árvore. 2015;39:439-45.

Iacona C, Muleo R. Light quality affects *in vitro* adventitious rooting and *ex vitro* performance of cherry rootstock colt. Scientia Horticulturae. 2010;125:630-6.

Joshi I, Bisht P, Sharma VK, Uniyal DP. *In vitro* clonal propagation of mature *Eucalyptus* F1 hybrid (*Eucalyptus tereticornis* Sm. x *E. grandis* Hill ex. Maiden). Silvae Genetica. 2003;52(n.3/4):110-3.

Kerbaui GB. Fisiologia vegetal. 2^a ed. Rio de Janeiro: Guanabara Koogan; 2012.

Kirdmanee C, Kitaya Y, Kozai T. Effects of CO₂ enrichment and supporting material *in vitro* on photoautotrophic growth of *Eucalyptus* plantlets *in vitro* and *ex vitro*. In Vitro Cellular and Developmental Biology - Plant. 1995;31:144-9.

Kozai T. Photoautotrophic micropropagation—environmental control for promoting photosynthesis. Propagation of Ornamental Plants. 2010;10:188-204.

Mafia RG, Alfenas AC, Ferreira EM, Zarpelon TG, Siqueira L. Crescimento de mudas e produtividade de minijardins clonais de eucalipto tratados com rizobactérias selecionadas. Revista Árvore. 2005;29:843-51.

Moura LC, Titon M, Fernandes JSC, Santana RC, Oliveira MLR. Micropropagação de sucupira preta por meio de gemas axilares. Pesquisa agropecuária brasileira. 2012;47(12):1691-8.

Oliveira LS, Xavier A, Lopes AP, Takahashi EK, Otoni WC. Multiplicação e alongamento *in vitro* de clones híbridos de *Eucalyptus globulus*. Ciência Florestal. 2016;26:235-47.

Oliveira ML, Xavier A, Penchel RM, Santos AFS. Multiplicação *in vitro* de *Eucalyptus grandis* x *E. urophylla* cultivado em meio semissólido e em biorreator de imersão temporária. Scientia Forestalis. 2011;39(91):309-15.

Oliveira ML, Xavier A, Penchel Filho RM, Reis JP. Efeito do intervalo de imersão e de injeção de ar na multiplicação *in vitro* de *Eucalyptus grandis* x *Eucalyptus urophylla* em biorreator de imersão temporária. Ciência Florestal. 2014;24:37-45.

Penchel RM, Otoni WC, Xavier A. Tecnologia de biorreatores e propagação *in vitro*. In: Borém A. editor. Biotecnologia florestal. Viçosa, MG: Universidade Federal de Viçosa, 2007. p.75-92.

R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna. Disponível em: <http://www.R-project.org/>. 2016.

Rossato M, Schumacher PV, Costa Netto APD, Souza GC, Reis EF, Stein VC. Multiplication and *in vitro* rooting of *Campomanesia adamantium* Camb. Plant Cell Culture & Micropropagation. 2015;11(2):70-7.

Sotelo M, Monza J. Micropropagation of *Eucalyptus maidenii* elite trees. Agrociencia. 2007;9(6):81-9.

Taiz L, Zeiger E. Fisiologia vegetal. 5^a ed. Porto Alegre: Artmed; 2013.

Titon M, Xavier A, Otoni WC, Reis GG. Efeito do AIB no enraizamento de miniestacas e microestacas de *Eucalyptus grandis* W. Hill ex Maiden. Revista Árvore. 2003a;27:1-7.

Titon M, Xavier A, Reis GG, Otoni WC. Eficiência das minicepas e microcepas na produção de propágulos de clones de *Eucalyptus grandis*. Revista Árvore. 2003b;27:619-25.

Tronco KMDQ, Bisognin DA, Fleig FD, Horbach MA. *Ex vitro* rooting and acclimatization of *Ilex paraguariensis* A. St Hil. microcuttings. Cerne. 2015;21:371-8.

Wendling I, Xavier A, Paiva HN. Influência da miniestaquia seriada no vigor de minicepas de clones de *Eucalyptus grandis*. Revista Árvore. 2003;27:611-8.

Xavier A, Comério J. Microestaquia: uma

maximização da micropropagação de *Eucalyptus*. Revista Árvore. 1996;20:9-16.

Xavier A, Wendling I, Silva RL. Silvicultura clonal: princípios e técnicas. 2ªed. Viçosa, MG: Universidade Federal de Viçosa; 2013.