



Archivos de Zootecnia

ISSN: 0004-0592

pa1gocag@lucano.uco.es

Universidad de Córdoba

España

Carvalho, M.A.; Quesenberry, K.H.

AGRONOMIC EVALUATION OF ARACHIS PINTOI (KRAP. AND GREG.) GERMPLASM IN FLORIDA

Archivos de Zootecnia, vol. 61, núm. 233, marzo, 2012, pp. 19-29

Universidad de Córdoba

Córdoba, España

Available in: <http://www.redalyc.org/articulo.oa?id=49523362003>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

# AGRONOMIC EVALUATION OF *ARACHIS PINTOI* (KRAP. AND GREG.) GERMPLASM IN FLORIDA

## EVALUACIÓN AGRONÓMICA DE GERMOPLASMA DE *ARACHIS PINTOI* EN FLORIDA

Carvalho, M.A.<sup>1\*</sup> and Quesenberry, K.H.<sup>2</sup>

<sup>1</sup>Brazilian Agriculture Research Corporation (Embrapa). Embrapa Cerrados. Planaltina, DF. Brazil.

\*marcelo@cpac.embrapa.br

<sup>2</sup>University of Florida. Agronomy Department. Gainesville. Florida. USA.

### ADDITIONAL KEYWORDS

Tropical legume. Dry matter yield. Seed production. Nutritive value.

### PALABRAS CLAVE ADICIONALES

Leguminosa tropical. Materia seca. Producción de semillas. Valor nutritivo del forraje.

### SUMMARY

*Arachis pintoi* germplasm displayed great variability with respect to adaptation, dry matter yield, nutritive value, seed production, and nematode reaction. Average forage dry matter yield (FDMY) during the year 2003 was 4.35 Mg/ha, and ranged from zero to 9.10 Mg/ha. Crude protein and *in vitro* organic matter digestibility (IVOMD) were high and confirmed the fact that *A. pintoi* has excellent nutritive value. Average CP was 180 g/kg of DM and ranged from 130 to 220 g/kg of DM. IVOMD averaged 670 g/kg of DM and ranged from 600 to 730 g/kg of DM. Some accessions produced high seed yields reaching values above 1 Mg/ha, 18 and 30 months after sowing. The average seed production was 0.32 Mg/ha in 2003, and 0.43 Mg/ha in 2004. Overall, accessions PI 604817, 497574, 604815, 604810, 604808, and 604857 were the best adapted to the north central Florida environment and with the best agronomic characteristics. This research strength that general conclusions about the adaptation and agronomic value of a species to a particular environment should not be based on a single or a few germplasm lines.

### RESUMEN

El germoplasma de *Arachis pintoi* muestra una gran variabilidad con respecto a la adaptación, producción de materia seca, valor nutritivo, producción de semillas, y la reacción a los nematodos. El rendimiento medio de materia seca de forraje (FDMY) durante el año 2003 fue de 4,35 Mg/ha, y

osciló entre cero y 9,10 Mg/ha. La proteína bruta y la digestibilidad *in vitro* de la materia orgánica (IVOMD) fueron altas confirmando que *A. pintoi* tiene excelente valor nutritivo. La proteína bruta promedio fue de 180 g/kg de MS y osciló entre 130 a 220 g/kg de MS. La IVOMD fue de 670 g/kg de MS, promedio y osciló entre 600 y 730 g/kg de MS. Algunas cepas de semillas de alto rendimiento alcanzaron valores superiores a 1 Mg/ha, 18 y 30 meses después de la siembra. El promedio de producción de semillas fue 0,32 Mg/ha en 2003, y 0,43 Mg/ha en 2004. En general, las cepas PI 604817, 497574, 604815, 604810, 604808 y 604857 fueron las que mejor se adaptaron al medio ambiente de la zona norte-centro de Florida siendo las de mejores características agronómicas. Este estudio demuestra, que en general las conclusiones sobre la adaptación y el valor agronómico de una especie a un medio ambiente no deben estar basadas en una o unas pocas líneas de germoplasma.

### INTRODUCTION

Agronomic evaluation is a very important step in germplasm characterization programs. Although molecular and morphological characteristics are relevant, plant breeders and ultimately producers have their attention focused on the potential of the plant to grow well in their environment and to produce forage, grain, or other economic

Recibido: 10-6-09. Aceptado: 12-5-10.

Arch. Zootec. 61 (233): 19-29. 2012.

products. Thus, agronomic evaluation will always be a key component in selection and breeding programs.

When evaluating a species outside its original environment, it is important to assess its adaptation to the new ecosystem. Emphasis must be given to how soils, climate, and rainfall conditions will impact the growth of this *new species*. Along with adaptation, several agronomic characteristics can be measured. The importance of each variable will be defined by the use of the plant and by the environment where it will be cultivated. In the case of a forage crop such as *Arachis pintoi* growing in a subtropical environment like Florida, forage yield, forage nutritive value, seed production and winter survival are some of the characteristics that should be evaluated.

*Arachis pintoi* is native of and well adapted to certain tropical environments. According to Pizarro and Rincón (1994), *A. pintoi* was evaluated by the International Tropical Pastures Evaluation Network (RIEPT) in Brazil, Uruguay, Bolivia, Colombia, Peru, and Venezuela under savanna and humid tropical conditions. They concluded that it presented a wide range of adaptation and grows best under humid tropical conditions with total annual rainfall ranging from 2000 to 4000 mm.

Pizarro and Rincón (1994), however, reported that plants growing in a subtropical environment in Pelotas-Brazil, exposed to severe frosts (temperature < 0°C), presented a reduced growth. However these conditions did not result in a drastic reduction of the plant stands. In fact, plants recovered after the return of warm and rainy conditions.

The literature has abundant *A. pintoi* forage yield data collected in the tropics. In evaluations performed in Bolivia, Brazil, Ecuador, Colombia, and Peru, accession CIAT 17434 produced between 0 and 2.7 Mg/ha of dry matter (DM) during the rainy season and 0.04 to 2.8 Mg/ha during the dry season with a growing period of 12 weeks (Pizarro and Rincón, 1994). In Costa Rica,

Argel and Valerio (1993) reported forage yields of 7, 12 and 7 Mg/ha of dry matter for accessions CIAT 17434, 18744, and 18748, with 20 months of growth in Guapiles and San Isidro. In Puerto Rico, forage dry matter yields of 2.1 Mg/ha were harvested 16 weeks after planting from the accession CIAT 17434 (Argel, 1994).

As stated before, the literature has several examples of research work done in the past where agronomic characterization of *Arachis pintoi* was the primary goal. However, most of this work was done with a single germplasm accession, that latter was released as commercial cultivars in several different countries. These studies also have in common the fact that most of them were carried out in tropical regions. Therefore, there is a lack of information about other accessions of *A. pintoi* stored in germplasm banks, and there are little or no existing data regarding the performance of the same germplasm in subtropical conditions.

The goal of this research was to evaluate the agronomic adaptation, forage yield, forage nutritive value and seed yield of several *A. pintoi* germplasm accessions stored at the USDA-NPGS germplasm bank.

## MATERIAL AND METHODS

### THE GERMPLASM

Accessions of *A. pintoi* stored in the Southern Regional Plant Introduction Station of the National Plant Germplasm System (NPGS) located at Griffin, Georgia were transferred to the University of Florida in 2001 and 2002. A list of these accessions with information related to the respective PI numbers and sites of collection is presented in **table I**.

### ADAPTATION AND FORAGE DRY MATTER YIELD

In September 2001, 4 m<sup>2</sup> (2 m x 2 m) field plots of the original 24 accessions were established with four rooted cuttings in a randomized complete block design with two

## AGRONOMIC EVALUATION OF *ARACHIS PINTO*/GERMPLASM IN FLORIDA

replications at the Agronomy Department Forage Research Unit of the University of Florida near Gainesville. The experiment was located on a Pomona sand (siliceous, Hyperthermic Typic Psammaquents- Entisol) soil type, a flatwood soil characteristic of north central Florida. The soil was prepared by disking and 190 kg/ha of 0-10-20 fertilizer was applied and incorporated into the soil. *Arachis glabrata* Florigraze and Arbrook were used as local standards and were also planted following the same scheme.

During the years 2001, 2002, and 2003 plots were maintained with periodical hand weeding and applications of the herbicide Cadre (Imazapic), in accordance with the

recommended dosage for cultivated peanut (100 g/ha). Annual applications of 800 kg/ha of 0-10-20 were made in May 2002 and 2003.

Visual evaluations were made following the winter 2001/2002. In these evaluations number of plants per plot, plant survival and rate of spread were assessed. Plant survival was expressed as percentage of total plants alive and rate of spread in accordance with a scale of 1 to 5 (1 < 10% plot coverage, 2= 20%, 3= 30%, 4= 40%, and 5 > 50% plot coverage). These parameters were intended as indications of adaptation and winter survival. Average annual temperature and rainfall data during the experimental period is presented in **table II**. No dry matter harvest was made in 2002.

Forage dry matter yield (FDMY) was estimated on three dates during 2003, with an interval of 8 weeks between harvests. These dates were June 13<sup>th</sup> (Harvest 1), August 13<sup>th</sup> (Harvest 2), and October 21<sup>st</sup> (Harvest 3). To estimate FDMY a sample of 0.25 m<sup>2</sup> was harvested at ground level using battery-powered hand clippers as a cutting tool. Pinto peanut and weeds were manually separated and the samples were placed in a 65°C forced-air drier for 72 h. Dry weight was measured using a digital scale. Plot borders were uniformly cut at soil level with a rotary lawn mowing tractor after each harvest.

FDMY was analyzed with Proc GLM of SAS (SAS, 1989) using the following model:

$$Y_{ij} = \mu + B_i + A_j + e_{ij}$$

Where:

$\mu$ = mean;

$B_i$ = block effect;

$A_j$ = accession effect;

$e_{ij}$ = experimental error.

LSD at 5% significance level was used as the mean separation test.

This same model was used for forage nutritive value and seed production.

**Table I.** *Arachis pinto* accessions characterized in this study. (Cepas de *Arachis pinto* caracterizadas en este trabajo).

Accession number	Lat. (South)	Long. (West)	Altitude (meter)
476132	16° 08'	47° 12'	690
497541	18° 38'	44° 04'	640
497574	13° 23'	44° 05'	450
604798	16° 18'	46° 58'	630
604799	16° 19'	46° 51'	580
604800	16° 41'	46° 29'	540
604801	16° 42'	46° 25'	560
604803	14° 25'	44° 22'	510
604804	14° 20'	44° 25'	560
604805	16° 59'	45° 57'	570
604807	13° 18'	46° 48'	510
604808	13° 18'	46° 42'	500
604809	13° 02'	46° 45'	610
604810	13° 06'	46° 45'	600
604811	13° 51'	46° 52'	490
604812	14° 28'	46° 29'	500
604813	14° 27'	47° 00'	480
604814	15° 52'	39° 08'	50
604815	15° 49'	47° 58'	1080
604817	18° 38'	44° 04'	630
604856	16° 53'	42° 07'	360
604857	13° 23'	44° 05'	450
604858	15° 26'	47° 21'	700
604859	17° 03'	42° 21'	360

## CARVALHO AND QUESENBERRY

## FORAGENUTRITIVEVALUE

Forage samples harvested on August 13<sup>th</sup> 2003 were ground in a Wiley Mill, to

pass through a 1 mm screen. This ground tissue was analyzed for crude protein and *in vitro* organic matter digestibility (IVOMD).

**Table II.** Climatological data at the Forage Research Unit in Gainesville, FL, during the period of the agronomic evaluation field trial. (Datos climatológicos en la Forage Research Unit de Gainesville, FL, durante el periodo de evaluación agronómica).

	Temperature °C			Rainfall (mm)	N° days drought	N° of days below temperature		
	Min.	Max.	Mean			0°C	4°C	12°C
2001								
Sep	23.22	10.18	34.34	292.2	13	0	0	1
Oct	19.46	-0.11	31.48	2.9	29	1	4	11
Nov	17.51	-0.22	28.88	27.1	26	1	3	14
Dec	15.21	-5.44	28.24	38.4	27	6	9	18
2002								
Jan	12.77	-7.31	29.11	133.0	19	6	14	19
Feb	12.41	-8.09	28.8	27.7	25	6	16	24
Mar	17.45	-6.37	31.65	83.1	22	4	8	19
Apr	21.92	3.64	33.15	9.5	23	0	1	4
May	23.11	6.42	35.57	39.5	26	0	0	7
Jun	24.89	15.08	37.47	104.7	10	0	0	0
Jul	25.71	18.05	36.84	138.3	13	0	0	0
Aug	25.05	11.94	35.61	265.3	13	0	0	1
Sep	25.66	17.13	33.99	128.2	19	0	0	0
Oct	22.14	6.82	34.88	41.6	21	0	0	4
Nov	14.07	-4.31	30.08	133.1	22	7	13	23
Dec	10.96	-3.01	26.44	183.1	21	11	18	27
2003								
Jan	8.22	-8.33	24.17	4.5	27	17	23	31
Feb	13.42	-0.82	28.42	172.5	17	4	12	23
Mar	18.69	2.03	30.64	194.4	19	0	1	14
Apr	19.22	-0.26	30.99	41.4	26	1	2	15
May	24.05	11.64	34.81	50.9	24	0	0	1
Jun	25.27	16.44	34.49	297.8	12	0	0	0
Jul	25.92	19.91	36.04	121.4	11	0	0	0
Aug	25.53	19.55	34.04	206.2	9	0	0	0
Sep	23.87	13.62	33.17	211.0	17	0	0	0
Oct	20.66	7.61	31.34	103.7	25	0	0	7
Nov	17.39	-2.63	31.62	34.1	25	2	4	15
Dec	10.28	-6.46	25.16	18.9	24	12	21	31
2004								
Jan	10.98	-6.16	27.29	33.9	24	13	17	28
Feb	12.35	-1.76	27.82	143.6	15	2	9	24
Mar	16.54	-1.35	29.00	25.5	27	1	6	25
Apr	17.95	1.69	31.09	40.3	25	0	6	23
May	23.92	5.41	36.13	16.4	28	0	0	4
Jun	25.77	18.03	36.15	174.3	16	0	0	0
Jul	25.99	18.22	35.28	254.8	13	0	0	0
Aug	25.69	20	34.94	150.7	12	0	0	0

## AGRONOMIC EVALUATION OF *ARACHIS PINTOI* GERMPLASM IN FLORIDA

The nitrogen (N) analysis started with sample digestion using a modification of the aluminum block digestion procedure (Gallaher *et al.*, 1975). In this procedure samples of 0.25 g and catalyst of 1.5 g of 9:1  $K_2SO_4$ : $CuSO_4$  were used during a 4 h digestion at 375°C using 6 mL of  $H_2SO_4$  and 2 mL of  $H_2O_2$ . Digestate was then filtrated and N was determined by semiautomated colorimetry (Hambleton, 1977).

IVOMD was performed by a modified two-stage technique (Moore and Mott, 1974). Both crude protein and IVOMD were expressed on an organic matter basis.

### SEED PRODUCTION

In February 2003 and 2004 plots were sampled to assess seed production. Single samples were taken in each plot using a soil core sampler of 12.5 cm diameter and 24 cm depth (0.01227 m<sup>3</sup>). The soil was screened on a sieve with a 0.6 x 0.6 cm mesh to remove the pods from the soil. The pods were then dried at room temperature for 6 weeks and weighed. Calculations were made to extrapolate the values in terms of production per hectare.

## RESULTS AND DISCUSSION

### ADAPTATION AND FORAGE DRY MATTER YIELD

The evaluations performed in 2002, eight, eleven, and twelve months after planting, revealed that although *A. pinto* presented great reduction of green tissue during the winter, most plants did not die, and in fact, they fully recovered when the temperature warmed up and soil moisture increased (**table III**). During the 2001/2002 winter, temperatures reached 0°C or less on 24 different occasions.

The average plant survival (PS%) in Evaluation 1 was 79%, which shows that *A. pinto* can tolerate winters where freezing and frosting are normal occurrences. However, when rate of spread (RS) was analyzed, most of the accessions were covering less than 50% of the plot area.

Cultivars Florigraze and Arbrook presented 100% of plant survival, but they also covered less than 50% of the plot area.

This same trend was displayed in Evaluations 2 and 3, where average plant survival recorded was superior to 80% and plot coverage inferior to 50%. Actually, this is one of the biggest problems that *A. pinto* and also *A. glabrata* display, the fact that both species required a long period to establish themselves and cover the area where they were planted.

Even with general small plot cover, differences among accessions were noted regarding plant survival and plot coverage, which is an indication of variation in adaptation to the north Florida environment. Accessions PI 497574, 604798, 604800, 604803, 604807, 604814, 604817, and 604857 were the ones that appeared better adapted, with values of plant survival and rate of spread superior to the others.

Pizarro and Rincón (1994) affirmed that *A. pinto* slow establishment is a limitation to adoption as producers want to see quick results. They suggested that when seeds are used as the propagation material the establishment time is shortened compared to vegetative propagation. French *et al.* (1994) stated that although adequate establishment has been achieved in Florida with rhizoma peanut (*A. glabrata*) in a single year, a typical planting demands 2 to 3 yr for full development. Cook *et al.* (1994) agreed with the authors previously cited. They believe that the major constraint to farmer acceptance of these species is the complexity and cost of establishment.

Slow establishment is particularly significant where weed pressure is high. This fact will require extra resources to assure an adequate initial coverage that will extend the period of utilization of the pasture or hay field. Research is necessary to determine the best planting method, using the different factors that could result in a fast and inexpensive pasture establishment.

A careful analysis of these evaluations

**Table III.** Winter survival evaluations of *Arachis pintoi* at the Forage Research Unit in Gainesville-FL. (Evaluación de la supervivencia invernal de *Arachis pintoi* en la Forage Research Unit de Gainesville-FL).

Accession PI number	Evaluation 1 (04/25/2002)			Evaluation 2 (07/03/2002)			Evaluation 3 (08/23/2002)		
	PL	PS (%)	RS	PL	PS (%)	RS	PL	PS (%)	RS
476132	3.0	75	1.0	3.0	75	2.4	4.0	100	2.0
497541	2.0	50	0.5	2.0	50	1.0	4.0	100	1.0
497574	4.0	100	2.5	3.0	75	1.5	4.0	100	2.5
604798	3.5	87	2.0	3.0	75	3.0	4.0	100	3.0
604799	3.0	75	3.0	3.0	75	2.5	4.0	100	2.0
604800	3.5	87	2.0	3.0	75	2.0	3.5	87	2.5
604801	1.0	25	0.5	1.0	25	1.0	3.5	87	2.0
604803	3.5	87	2.0	3.5	87	2.8	4.0	100	3.0
604804	4.0	100	1.5	3.0	75	1.5	3.0	75	1.5
604805	3.0	75	2.0	3.0	75	2.0	4.0	100	1.5
604807	3.5	87	2.5	3.5	87	2.0	4.0	100	3.0
604808	4.0	100	4.0	4.0	100	4.5	4.0	100	2.0
604809	3.0	75	2.0	3.0	75	3.0	3.5	87	2.5
604810	3.0	75	2.5	2.0	50	2.0	3.5	87	1.5
604811	4.0	100	2.5	4.0	100	2.0	4.0	100	1.5
604812	2.0	50	1.0	1.5	37	1.5	4.0	100	2.0
604813	4.0	100	1.5	4.0	100	1.5	4.0	100	1.5
604814	1.5	37	0.5	3.0	75	2.4	3.5	87	4.0
604815	2.0	50	2.0	0.5	12	1.0	4.0	100	2.0
604817	4.0	100	2.5	3.8	94	3.5	4.0	100	3.5
604856	2.5	62	1.0	3.0	75	1.5	4.0	100	2.0
604857	3.5	87	2.0	3.8	94	3.2	4.0	100	3.0
604858	3.5	87	3.0	3.3	81	3.5	4.0	100	2.5
604859	1.0	25	1.0	3.8	94	3.2	1.5	37	1.5
Arbrook	4.0	100	4.0	3.3	81	3.5	4.0	100	4.5
Florigraze	4.0	100	2.5	3.0	75	3.0	4.0	100	2.5
LSD	2.2	55	2.0	2.6	65	2.2	0.9	21	1.8

PL: number of plants per plot; PS (%): plant survival; RS: Rate of spread.

demonstrates that there are large differences among accessions in relation to environmental adaptation. Accessions PI 604801, 604804, 604812, 604814, and 604859, presented small plot coverage in each one of the three evaluations.

Forage dry matter yield (FDMY) was assessed three times during summer 2003 with an 8-week interval between harvests. Significant differences ( $p < 0.01$ ) were present among accessions in all three FDMY harvests, indicating that accessions showed

variability in their environmental adaptation, an attribute that will consequently impact their forage production potential.

Forage dry matter yield at each harvest date in 2003 is presented in the **table IV**. A large amount of variability was present among these *A. pintoi* accessions with respect to their FDMY. The average FDMY at harvest 1 was 1.34 Mg/ha, ranging from 0.06 to 2.86 Mg/ha. More than half of the accessions presented superior performance producing above the average, but



# AGRONOMIC EVALUATION OF *ARACHIS PINTOI* GERMPLASM IN FLORIDA

accessions PI 604801, 604804, 604812, and 604814 had very low yields. These low yields are a reflection of poor plot coverage for these accessions. In comparison, accessions PI 604817, 604815, 604803, 604808, 604810, 604857, and 497574 were the most productive, showing no differences in FDMY to the rhizoma peanut cultivars.

At harvest 2, the average FDMY of *A. pinto* germplasm showed a 20% increase

**Table IV.** Forage dry matter yield (Mg/ha) of *Arachis pinto* germplasm Gainesville, FL in 2003. (Rendimiento de materia seca de forraje (Mg/ha) de germoplasmas de *Arachis pinto* en Gainesville, FL, en 2003).

Accession	Harvest*			Annual FDMY
	1	2	3	
476132	0.79	0.75	0.98	2.52
497541	1.18	0.94	0.44	2.56
497574	1.85	3.00	2.50	7.36
604798	1.51	1.52	1.38	4.42
604799	1.35	1.74	1.60	4.69
604800	1.39	1.98	1.33	4.71
604801	0.34	1.10	0.93	2.37
604803	1.99	1.93	1.74	5.67
604804	0.06	0	0	0.06
604805	1.58	1.48	0.88	3.93
604807	1.38	0.78	0.56	2.73
604808	1.98	2.62	1.99	6.59
604809	1.47	0.38	1.14	2.99
604810	1.96	2.54	2.14	6.63
604811	1.06	1.45	1.63	4.14
604812	0.19	0.6	0.82	1.61
604813	0.98	1.72	1.01	3.71
604814	0.10	0	0	0.10
604815	2.32	2.12	2.30	6.74
604817	2.86	3.45	2.79	9.10
604856	1.12	1.90	1.56	4.58
604857	1.85	2.26	2.24	6.35
604858	1.59	2.25	1.87	5.71
604859	1.38	2.29	1.63	5.30
Arbrook	3.16	4.75	1.76	9.67
Florigraze	2.14	1.89	0.95	4.98
LSD	0.79	0.75	0.98	3.74

\*Harvest 1: 13 Sep. 2003; Harvest 2: 13 Aug. 2003; Harvest 3: 21 Oct. 2003.

reaching 1.62 Mg/ha. Accessions PI 604804 and 604814 showed no production, which reflects small plot coverage and poor adaptation to the north Florida environment. Here once more, half of the accessions presented superior performance with FDMY above the average. Accessions PI 604817 and 497574 were the most productive, showing no difference to Arbrook, which produced 4.75 Mg/ha. Florigraze however, was intermediate in FDMY and similar to a group of *A. pinto* accessions. This is somewhat surprising, since it is regarded by many as the most adapted and productive of the rhizoma peanut cultivars.

Harvest 3 presented mean FDMY of 1.39 Mg/ha, which corresponded to a drop of about 20% in relation to the preceding harvest. Arbrook and Florigraze exhibited a 50% reduction in FDMY compared to previous period. *A. pinto* germplasm also exhibited this reduction, however in a lower degree. This reduction in FDMY could represent a change in nutrient partitioning as a process to overcome restriction in growth due to the shortening days in autumn. Accessions PI 604817, 497574, 604815, and 604857 had the highest FDMY and were significantly higher than Florigraze and Arbrook rhizoma peanut.

Annual average FDMY for *A. pinto* accessions was 4.36 Mg/ha (table III). Arbrook produced 9.67 Mg/ha and was the most productive. Accessions PI 604817, 497574, 604815, 604810, 604808, and 604857 were the most productive among the *A. pinto* germplasm and were not different from Arbrook.

Several *A. pinto* accessions yielded more than 5 Mg/ha which can be considered a satisfactory legume production in north central Florida. In fact, this number is similar to yields displayed by other warm-season legumes used in Florida, and even could be compared to *A. pinto* or others legume species yields obtained in tropical regions.

Kretschmer and Wilson (1988) reported 3-year average FDMY for *Desmodium*



*heterocarpon* Florida, *Macroptilium atropurpureum* Siratro, and *Arachis kretschmeri* Pantanal of the order of 2.97, 2.39, and 1.16 Mg/ha respectively. Annual yields of *Stylosanthes guianensis* Savanna, *Alysicarpus vaginalis* and *Indigofera hirsuta*. Flamingo ranged from 5 to 7 Mg/ha, in plantings at Brooksville, FL during years 1989 and 1990 (Williams *et al.*, 1993). Mislavy and Martin (2001) reported similar annual FDMY (5.70 Mg/ha) for *Aeschynomene evenia* at Ona, FL.

In tropical areas, comparable yields were presented by Souza *et al.* (1992) who evaluated *Stylosanthes* and *Centrosema* germplasm in central Brazil. Average FDMY of 1.50 Mg/ha were harvested in a 12-wk growing period during the summers of 1989 and 1990. Pizarro *et al.* (1996) reported annual FDMY of *Calopogonium mucunoides* of the order of 1.72 Mg/ha in research conducted at several different locations in South America.

Data from diverse agronomic characterization of *A. pinto* are also similar to those presented in this work. Argel and Pizarro (1992) stated that cv. Amarillo produced 2.30 Mg/ha in Planaltina, Brazil, during the rainy season. In Australia, Cook *et al.* (1990) stated that Amarillo yielded 6.50 and 7.30 Mg/ha/yr in unirrigated and irrigated conditions respectively. Also in Australia, Cook *et al.* (1994) presented average annual yields of 5.80 Mg/ha for Amarillo in two years of evaluation.

#### FORAGE NUTRITIVE VALUE

Crude protein and IVOMD values of *A. pinto* accessions from this research confirm those reported in the literature, and support the information that this species produces high nutritive value forage. The results endorse the claim of some that *A. pinto* could be considered a *tropical alfalfa* (table V).

Average crude protein concentration across *A. pinto* accessions was 180 g/kg ranging from 139 (PI 604800) to 225 g/kg (PI 604858). Florigraze CP was 172 g/kg while

that of Arbrook was 153 g/kg. When CP of the *A. pinto* germplasm was compared to the rhizoma peanut cultivars, the majority of the accessions demonstrated higher values than either rhizoma peanut cultivars.

Average IVOMD of the *A. pinto* germplasm was 670 g/kg, and ranged from 600 to 730 g/kg. Accession PI 604812 had the lowest value, while PI 604801 was the highest. Florigraze presented the highest

**Table V.** Crude protein (CP) and in vitro organic matter digestibility (IVOMD) of 8-wk regrowth of *Arachis pinto* at the Forage Research Unit in Gainesville, FL in 2003. (Proteína bruta y digestibilidad *in vitro* de la materia orgánica de rebrotes de 8 semanas de *Arachis pinto* en la Forage Research Unit en Gainesville, FL en 2003).

Accession	CP (g kg <sup>-1</sup> )	IVOMD (g kg <sup>-1</sup> )
476132	178	690
497541	147	630
497574	184	640
604798	170	680
604799	189	700
604800	139	700
604801	188	730
604803	195	670
604805	151	690
604807	156	680
604808	189	620
604809	148	620
604810	187	610
604811	189	690
604812	183	600
604813	166	660
604815	194	690
604817	212	670
604856	180	710
604857	222	720
604858	225	720
604859	173	710
Arbrook	153	690
Florigraze	172	740

LSD CP (0.05)= 30.3 g kg<sup>-1</sup>; LSD IVOMD (0.05)= 65 g kg<sup>-1</sup>.

## AGRONOMIC EVALUATION OF *ARACHIS PINTOI* GERMPLASM IN FLORIDA

IVOMD value (740 g/kg), however several *A. pinto* accessions were not different from it.

CP and IVOMD displayed in this research could be considered high and comparable to those reported for *A. pinto* in other investigations. Average values of CP in the leaves of accession CIAT 17434 varied from 122 to 218 g/kg in Colombia, during dry and rainy seasons, respectively. In stems the same accession presented values of 93 and 135 g/kg during the same seasons (Argel and Pizarro, 1992). In Brazil, Purcino and Viana (1994) reported total-herbage CP values of 183, 157 and 161 g/kg, for accessions BRA-013251 (PI 338447), BRA-015253 (PI 604859) and BRA-015598 (PI 604815), respectively. Rincon *et al.* (1992) reported CP values for the whole plant of 130 and 180 g/kg, during the dry and rainy season, respectively. Average values of IVOMD during the same periods were 67 and 62 g/kg.

Average IVOMD of 168 days growth forage from three *A. pinto* accessions in Brazil was 610 g/kg (Pizarro *et al.*, 1992). In Australia, Amarillo IVDMD was 730 g/kg in 2 yr of evaluation (Cook *et al.*, 1994).

### SEED PRODUCTION

Another important factor for which genetic variability should be investigated is seed production. In February 2003 and 2004, respectively, 18 and 30 mo after planting, samples were collected in each plot to assess this trait.

In 2003, the average seed production was 0.32 Mg/ha, ranging from zero to 2 Mg/ha (**table VI**). Only 60% of the *A. pinto* germplasm yielded seeds, however, if we consider only those accessions with production above 0.1 Mg/ha this number falls below 50%. Accession PI 604857 was the most productive yielding 2 Mg/ha.

In 2004, with the exception of three accessions, all accession produced seeds. The average seed production was higher than the previous year reaching 0.43 Mg/ha, ranging from zero to 1.58 Mg/ha (**table**

**Table VI.** Seed production (Mg/ha) of *Arachis pinto* at the Forage Research Unit in Gainesville, FL in 2003 and 2004. ( Producción de semillas (Mg/ha) de *Arachis pinto* en la Forage Research Unit en Gainesville, FL en 2003 y 2004).

Accession	2003*	2004*
476132	0.01	0.07
497541	0.00	0.03
497574	0.00	0.36
604798	0.00	0.69
604799	0.75	1.58
604800	0.03	0.16
604801	0.00	1.10
604803	0.76	0.20
604804	0.00	0.00
604805	0.59	1.21
604807	0.29	0.36
604808	0.30	1.01
604809	0.91	0.43
604810	0.07	0.06
604811	0.00	0.02
604812	1.30	1.32
604813	0.00	0.00
604814	0.00	0.10
604815	0.47	0.17
604817	0.00	0.00
604856	0.08	0.05
604857	2.00	0.62
604858	0.06	0.27
604859	0.00	0.63
Arbrook	0.00	0.00
Florigrase	0.00	0.00
LSD	0.48	0.90

\*pod weight.

**V**). In general, yields of most accessions increased compared to 2003 evaluation. Exceptions were PI 604803, 604809, 604815, and 604857. Curiously, these accessions were among the most productive in 2003. Although 87% of the accessions produced some seeds, six of them yielded less than 0.10 Mg/ha, amount that would not be useful for commercial harvest. PI 604799 had the highest yield (1.58 Mg/ha) in 2004 although it was intermediate in seed yield in 2003.

Seed production obtained in this research for some of the accessions was comparable to that obtained by other authors working with *A. pinto* in tropical conditions. For legumes this trait has particular importance because it can be related to persistence. A species with high seed production capacity may have an advantage over one lacking this characteristic, because in theory, pasture establishment by seeds is simpler and cheaper than when vegetative material is used. Also, plants established by seeds will grow faster (Baruch and Fisher, 1992), and consequently will have higher ground covered area. *A. pinto* has also a notable characteristic that differentiates it from other legumes in relation to seed production, in that the seeds are located below ground.

As previously stated, yields in this research were comparable to others found in the literature. Cook and Franklin (1988) reported mechanized harvest yield for Amarillo in Australia of 1.40 Mg/ha 12 mo after planting. In other work, seed yield of Amarillo in Australia reached an average production of 2.80 Mg/ha (Cook and Loch, 1993).

In a Costa Rica location with 4260 mm annual rainfall, Diulgheroff *et al.* (1990) obtained seed yields of 1.95 Mg/ha, 12 months after sowing the cv. Maní Forrajero perenne. Also in Costa Rica, Argel and Valerio (1993) reported average yields of 0.59, 0.54, 0.50, and 0.52 Mg/ha over the accessions CIAT 17434, 18744, and 18748, at 8, 12, 16, and 20 mo after sowing, respectively. Rincon *et al.* (1992) stated that seed yields of 2 Mg/ha were obtained for cv. Maní Forrajero perenne in mixed pasture with *Brachiaria* under grazing. This supports the idea of excellent persistence potential for *A. pinto* due to seedling recruitment.

## REFERENCES

Argel, P.J. 1994. Regional experience with forage *Arachis* in Central America. In: P.C. Kerridge and

## CONCLUSIONS

*Arachis pinto* germplasm displayed large variability with respect to its adaptation, dry matter yield, nutritive value and seed production.

Annual average dry matter yield for some *A. pinto* accessions was close to the forage production presented by both rhizoma peanut cultivars. Crude protein and IVOMD were high and confirmed the fact that *A. pinto* has superior nutritive value.

Some accessions produce elevated seed yields reaching values superior to 1 Mg/ha 18 and 30 mo after sowing, which would give a pinto peanut cultivar a great advantage over a rhizoma peanut cultivar, in terms of pasture establishment and seedling recruitment, if similar forage yields and persistence were present.

The information generated by this research reaffirms once more for plant selectors and plant breeders that conclusions about a species genetic diversity, adaptation to environments, and agronomic characteristics must not be based on a single or small number of germplasm accessions. This information will also be useful for genetic resources management, germplasm selection and plant breeding programs, and genetic studies in general. Ultimately, this work will contribute for the utilization of the genetic resources of *A. pinto* stored in the germplasm banks.

## ACKNOWLEDGEMENTS

This work was funded partially by the National Council for Scientific and Technological Development (CNPq), a foundation linked to the Ministry of Science and Technology (MCT), to support Brazilian research, and the Brazilian Agriculture Research Corporation (Embrapa).

B. Hardy (ed.). Biology and agronomy of forage *Arachis*. CIAT. Cali. Colombia. pp. 134-143.

## AGRONOMIC EVALUATION OF *ARACHIS PINTOI* GERMPLASM IN FLORIDA

- Argel, P.J. and Valerio, A. 1993. Effect of crop age on seed yield of *Arachis pintoi* at two sites in Costa Rica, Central America. In: Proc. Int. Grassl. Congr., 17<sup>th</sup>. Palmerston North. New Zealand. 8-21 Feb. New Zealand Grassland Association. pp. 1696-1698.
- Argel, P.J. and Pizarro, E.A. 1992. Germplasm case study: *Arachis pintoi*. In: Pastures for the tropical lowlands-CIAT's Contribution. CIAT. Cali. Colombia. pp. 57-76.
- Baruch, Z. y Fisher, M. 1992. Efecto del método de siembra y de la textura del suelo sobre el crecimiento y desarrollo de *Arachis pintoi*. Proc. Reunión Sabanas. Red Internacional de Evaluación de Pastos Tropicales (RIEPT). CIAT. Brasília. Brazil.
- Cook, B.G. and Loch, D.S. 1993. Commercialisation of *Arachis pintoi* cv. Amarillo in northern Australia. Proc. Int. Grassl. Congr., 17<sup>th</sup>. Palmerston North. New Zealand Grassland Association. 8-21 Feb. New Zealand.
- Cook, B.G. and Franklin, T.G. 1988. Crop management and seed harvesting of *Arachis pintoi* Krap. et Greg. nom. nud. *J Appl Seed Prod*, 6: 26-30.
- Cook, B.G., Williams, R.J. and Wilson, G.P.M. 1990. Register of Australian plant cultivars. B. Legumes. 21. *Arachis* (a) *Arachis pintoi* Krap. & Greg. nom. nud. (pinto peanut) cv. Amarillo. *Aust J Exp Agr*, 30: 445-446.
- Cook, B.G., Jones, R.M. and Williams, R.J. 1994. Regional experience with forage *Arachis* in Australia. In: P.C. Kerridge and B. Hardy (Eds.). Biology and agronomy of forage *Arachis*. CIAT. Cali. Colombia. pp. 158-168.
- Diulgheroff, S., Pizarro, E.A., Ferguson, J.E. and Argel, P.J. 1990. Seed multiplication of tropical forage species in Costa Rica. *Past Trop*, 12: 15-23.
- French, E.C., Prine, G.M., Ocumpaugh, W.R. and Rice, R.W. 1994. Regional experience with forage *Arachis* in the United States. In: P.C. Kerridge and B. Hardy (Eds.). Biology and agronomy of forage *Arachis*. CIAT. Cali. Colombia. pp. 169-186.
- Gallaher, R.N., Weldon, C.O. and Futral, J.G. 1975. An aluminum block digester for plant and soil analysis. *Soil Sci Soc Am Proc*, 39: 803-806.
- Hambleton, L.G. 1977. Semiautomated method for simultaneous determination of phosphorous, calcium and crude protein in animal feeds. *J.A.O.A.C.*, 60: 845-852.
- Kretschmer, A.E. and Wilson, T.C. 1988. A new seed production *Arachis* sp. with potential as a forage in Florida. *Soil Crop Sci Soc Fla Proc*, 47: 229-233.
- Mislevy, P. and Martin, E.G. 2001. Harvest management of *Aeschynomene evenia*. Proc. Soil and Crop Science Society of Florida. Stuart. Florida. pp. 35-37.
- Moore, J.E. and Mott, G.O. 1974. Recovery of residual organic matter from *in vitro* digestion of forages. *J Dairy Sci*, 57: 1258-1259.
- Pizarro, E.A. and Rincón, A.C. 1994. Regional experience with forage *Arachis* in South America. In: P.C. Kerridge and B. Hardy (ed.). Biology and agronomy of forage *Arachis*. CIAT. Cali. Colombia. pp. 144-157.
- Pizarro, E.A., Ramos, A.K.B. and Carvalho, M.A. 1996. Forage potential and seed production of accessions of selected *Calopogonium mucunoides* in the Brazilian Cerrados. *Past Trop*, 18: 25-31.
- Pizarro, E.A., Carvalho, M.A., Valls, J.F.M. y Maciel, D. 1992. *Arachis* spp.: Evaluación agronómica en áreas bajas del Cerrado. En: Proc. Reunión Sabanas. Red Internacional de Evaluación de Pastos Tropicales (RIEPT). CIAT. Brasília. Brazil. pp. 353-356.
- Purcino, H.M.A. and Viana, M.C.M. 1994. Preliminary evaluations of *Arachis pintoi* in lowland areas. Proc. da Reunião Anual da Soc. Bras. Zootec. 31<sup>st</sup>. Maringá, PR. EDUEM. Brazil.
- Rincón, A.C., Cuesta, M.P.A., Pérez, B.R., Lascano, C.E. y Ferguson, J. 1992. Maní forrajero perenne (*Arachis pintoi* Krapovickas et Gregory): Una alternativa para ganaderos y agricultores. Boletín Técnico 219. IAC/CIAT. Cali. Colombia.
- SAS. 1989. SAS/STAT User's Guide, Version 6. 4<sup>th</sup> ed. SAS Inst. Inc. Cary. NC.
- Sousa, M.A., Pizarro, E.A., Carvalho, M.A., Grof, B. and Schulz, A.L. 1992. Agronomic evaluation of forage grasses and legumes in Planaltina, Distrito Federal, Brasília. In: Proc. Reunión Sabanas. Red Internacional de Evaluación de Pastos Tropicales (RIEPT). Brasília. CIAT. Brazil.
- Williams, M.J., Chambliss, C.G. and Brolmann, J.B. 1993. Potential of Savanna stylo as a stockpiled forage for the subtropical USA. *J Prod Agric*, 6: 553-556.