



Journal of Pharmacy & Pharmacognosy
Research

E-ISSN: 0719-4250

editor@jppres.com

Asociación de Académicos de Ciencias
Farmacéuticas de Antofagasta
Chile

Chaudhary, Pankaj H.; Rai, Pallavi D.; Deore, Sharada L.; Khadabadi, Somshekhar S.
Pharmacognostical and phytochemical studies on roots of *Bombax ceiba* Linn.
Journal of Pharmacy & Pharmacognosy Research, vol. 2, núm. 6, noviembre-diciembre,
2014, pp. 172-182
Asociación de Académicos de Ciencias Farmacéuticas de Antofagasta
Antofagasta, Chile

Available in: <http://www.redalyc.org/articulo.oa?id=496050271002>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative



Pharmacognostical and phytochemical studies on roots of *Bombax ceiba* Linn.

[Estudios farmacognóstico y fitoquímico de las raíces de *Bombax ceiba* Linn.]

Pankaj H. Chaudhary, Pallavi D. Rai, Sharada L. Deore, Somshekhar S. Khadabadi

Department of Pharmacognosy, Government College of Pharmacy, Kathora Naka, Amravati – 444604, Maharashtra, India.

*E-mail: pankaj3_chaudhary@rediffmail.com

Abstract

Context: *Bombax ceiba* Linn. (Bombacaceae) is a well-known plant for its antihypertensive, antioxidant, antidiabetic, aphrodisiac and uterine tonic properties.

Aims: To study pharmacognostical, physicochemical and phytochemically the roots of this plant.

Methods: Pharmacognostical study included the macroscopic characters like size, color, surface characteristics, texture, fracture characteristics and odor of the roots. The intact root as well as powdered drug were studied under a microscope to analyze the cellular characteristics of the drug. Physicochemical parameter like extractive values, loss on drying (LOD), total ash, water-soluble and acid insoluble ash, foaming index and hemolytic index of *Bombax ceiba* root powder were determined as per WHO guidelines. Preliminary phytochemical screening and qualitative chemical examination studies have been carried out for the various phytoconstituents. HPTLC have also carried out using cyclohexane: diethyl ether: ethyl acetate as mobile phase.

Results: Chemical evaluation and TLC studies shown presence of alkaloids, glycosides, flavonoids, steroids, saponins and tannins. The microscopic characters have shown presence of cork, cambium, xylem vessels, stone cells, starch grains, calcium oxalate crystals and phloem fibers. Microscopy analysis of the powder included the cork cells, fibers, calcium oxalate crystals and vessel. The presence of steroids was confirmed in HPTLC fingerprinting studies.

Conclusions: Pharmacognostical and preliminary phytochemical screening of *Bombax ceiba* roots will be useful in order to authenticate, standardize and avoid any adulteration in the raw material. The diagnostic microscopic characters and physicochemical data will be helpful in the development of a monograph. The chromatographic fingerprinting profile can be used to standardize extracts and formulations containing *Bombax ceiba* roots.

Keywords: *Bombax ceiba*; HPTLC; microscopy; steroids; thin layer chromatography.

Resumen

Contexto: *Bombax ceiba* Linn. (Bombacaceae) es una planta conocida por sus propiedades anti-hipertensiva, antioxidante, antidiabética, afrodisíaca y tónica uterina.

Objetivos: Estudiar farmacognóstica, fisicoquímica y fitoquímicamente las raíces de esta planta.

Métodos: El estudio farmacognóstico incluyó los caracteres macroscópicos como tamaño, color, características de la superficie, textura, características de fractura y el olor de las raíces. La raíz intacta y la droga en polvo se estudiaron bajo el microscopio para analizar las características celulares de la droga. Los parámetros fisicoquímicos como valores extractivos, pérdida por desecación (LOD), cenizas totales, índice de formación de espuma soluble en agua y cenizas insolubles en ácido, y el índice hemolítico del polvo de raíz de *Bombax ceiba* se determinaron según directrices de la OMS. Se realizaron estudios de tamizaje fitoquímico preliminar y examen químico cualitativo para los distintos fitoconstituyentes. También se realizó el estudio por HPTLC usando como fase móvil ciclohexano: éter dietílico: acetato de etilo.

Resultados: La evaluación química y los estudios de TLC mostraron presencia de alcaloides, glucósidos, flavonoides, esteroides, saponinas y taninos. Los caracteres microscópicos han demostrado la presencia de corcho, cambium, vasos del xilema, células pétreas, granos de almidón, cristales de oxalato de calcio y fibras del floema. El análisis microscópico del polvo incluyó las células de corcho, fibras, cristales de oxalato de calcio y vaso. La presencia de esteroides se confirmó en estudios de huellas por HPTLC.

Conclusiones: El tamizaje farmacognóstico y fitoquímico preliminar de las raíces de *Bombax ceiba* será útil para autenticar, estandarizar y evitar cualquier adulteración en la materia prima. Los caracteres microscópicos de diagnóstico y los datos fisicoquímicos serán útiles en el desarrollo de una monografía. El perfil cromatográfico de huellas (*fingerprinting*) puede utilizarse para estandarizar extractos y formulaciones que contengan raíces de *Bombax ceiba*.

Palabras Clave: *Bombax ceiba*; cromatografía capa delgada; esteroides; HPTLC; microscopía.

ARTICLE INFO

Received | Recibido: July 27, 2014.

Received in revised form | Recibido en forma corregida: December 5, 2014.

Accepted | Aceptado: December 8, 2014.

Available Online | Publicado en Línea: December 10, 2014.

Declaration of interests | Declaración de Intereses: The authors declare no conflict of interest.

Funding | Financiación: The authors confirm that the project have not funding or grants.



INTRODUCTION

Ayurvedic system of medicine is a traditional system of medicine, native to the Indian subcontinent and practiced in other parts of the world as a form of alternative medicine. In Sanskrit, the word *Ayurveda* consists of the words *ayus*, meaning "life", and *Veda*, meaning "related to knowledge" or "science". Evolving throughout its history, Ayurveda remains an influential system of medicine in South Asia. Over the past centuries, ayurvedic practitioners, called "Ayurvedacharyas," have identified a number of medicinal preparations and surgical procedures for curing various ailments and diseases (CRPA, 2002).

However, there is a need to transform Ayurveda into a dynamic, scientifically validated and evidence-based system which takes its roots from rich knowledge base of oral tradition and scriptures. The major hurdle in the wider acceptability of Ayurveda and its products is the lack of proper standardization techniques and its unpreparedness to accept global challenges (Wagner and Farnsworth, 1994). The World Health Organization (WHO) also has emphasized the need to ensure the quality of the medicinal plant products (Hamberger et al., 1991).

Standardization of herbal drugs is very much important to establish their products in the global markets. Standardization starting from production of quality materials, analysis of raw materials for authentication, presence of foreign matter, organoleptic evaluation, microscopic examination, extractive values, chromatographic profiles, pesticides residue, presence of heavy metals, etc., is necessary for standardization of drugs. Similarly, the standardization methods of medicinal plants and its extracts have great importance in the fields of cosmetics and nutraceuticals, which are emerging as the two most important segments in the global markets (Masood, 1997).


The present study has been performed on *Bombax ceiba* Linn or *Bombax malabaricum* D.C. or *Salmalia malabarica* (DC.) Schott & Endl is belonging to family Bombacaceae (The Wealth of India, 2003). The therapeutic effects have been

reported in roots, gums, stem bark, flowers, seeds, prickles and young fruits. The family Bombacaceae consists of 22 genera and 150 species. Genera *Bombax* consists of 60 species, *Ceiba* 15 species, *Durio* 15 species, *Salmalia* 10 species and *Adansonia* 10 species (Rajendra, 2007). This tropical tree has a straight tall trunk and its leaves are deciduous in winter. Red flowers with five petals appear in the spring before the new foliage. It produces a capsule which, when ripe, contains white fibers like cotton. Its trunk bears spikes to deter attacks by animals (Kirtikar and Basu, 1994).


Reports have shown the presence of glycosides and tannins in roots, stem and leaf. In the stem some alkaloids and in roots, proteins are identified. The stem bark and root contains mangiferin, lupeol and β -sitosterol (Fig. 1). The root bark has 3 naphthalene derivatives related to gossypol (toxic principle of cotton seed) and called as 'semigossypol'. Flowers contain β -sitosterol, traces of essential oil, kaempferol and quercetin. On hydrolysis gums yields arabinose, galactose, galacturonic acid and rhamnose (NISC, 2001).

It has been claimed in Ayurveda that *Bombax ceiba* possesses proven medicinal properties and is the ingredient of many formulations. The roots are sweet, cooling, stimulant, restorative, astringent, alternative, aphrodisiac, demulcent, emetic and tonic. It is used in the treatment of diarrhea, dysentery, menorrhagia, styptic and for wounds. The gum is cooling, astringent, stimulant, aphrodisiac, tonic and demulcent in nature. It is useful in dysentery, hemoptysis, and pulmonary tuberculosis, influenza, burning sensation, menorrhagia and enteritis. Bark is mucilaginous, demulcent, emetic and tonic. Flowers are astringent and good for skin troubles and hemorrhoids. Seeds are useful in treating gonorrhea and chronic cystitis. A paste made out of prickles is good for restoring skin color especially on the face. Young fruits are useful in calculus affections, chronic inflammations and ulceration of bladder and kidney (Kirtikar and Basu, 1994; NISC, 2001).

A



B



C




Figure 1. Chemical structure of (A) lupeol, (B) β -sitosterol, (C) mangiferin present in *B. ceiba*.

Numbers of reports are available on pharmacognostic and phytochemical studies of stem (Rai Chaudhuri, 1971; Shantha et al., 2009) and leaf (Mehra and Karnick, 1968; Gopal and Gupta, 1972; Faizi and Ali, 1999; Puckhaber and Stipanovic, 2001; Versiani, 2007) of *Bombax ceiba* Linn. but no work has been reported till date on the pharmacognostic studies of *Bombax ceiba* roots. Although the presence of certain secondary metabolites in *Bombax ceiba* has been reported, no study depicting the complete phytochemical screening has been reported yet. Thus, in the present study an attempt has been made to perform the pharmacognostic studies and proximate analysis of *Bombax ceiba* roots. The preliminary phytochemical screening of the roots was also be done by extraction in different extracts and then performing chemical tests and TLC studies of the same.

MATERIALS AND METHODS

Plant material

Roots of *Bombax ceiba* were collected from Govt. Vidharbha Institute of Science & Humanities localities, Amravati (Maharashtra). The plant was identified and authenticated by Dr. Vishal R.

Marathe of Department of Botany, Shri Shivaji Science College, Amravati, and a voucher sample (Code: SSC-DOB07062011) was deposited at the Herbarium of the this Department. The vegetable drug was dried in the shade at room temperature. Dried roots were powdered in grinder and powder material was kept in an airtight container for further study.

Macroscopic evaluation

The size, color, surface characteristics, texture, fracture characteristics and odor of the roots were studied (Khandelwal, 2004).

Microscopic evaluation

The intact root, as well as powdered drug, was studied under a microscope to analyze the cellular characteristics of the drug.

Study of transverse section

The roots were taken in a test tube and 5% potassium hydroxide in methanol was added so that the sample remained submerged. The samples were boiled for few minutes. Transverse sections of the drugs were taken in a watch glass containing water with the help of a brush. The

sections were then transferred to a watch glass containing phloroglucinol-hydrochloric acid (1:1) and allowed to stain for 2-3 minutes. The sections were again transferred to watch glasses containing water, so as the excess stain was washed away. The sections were then placed on clean glass micro-slides, with the help of a brush. Few drops of water were added, and a clean cover-slip was placed on the slide. The slides were mounted for study on the microscope (Khandelwal, 2004). The transverse sections were also stained using iodine, methylene blue, Sudan red to identify the starch, mucilage, and fats and fixed oils.

Study of powder characteristics

The microscopic structures of powdered drugs were also studied using the slides prepared by above method using powdered drugs in place of sections (Khandelwal, 2004). The fiber length of the root powder was also determined using the reported methods (Khandelwal, 2004). The representative photos of sections were taken with the help of Digital Olympus microscope.

Physicochemical evaluation

Physicochemical parameter like extractive values, loss on drying (LOD), total ash, water-soluble and acid insoluble ash, foaming index and hemolytic index of *Bombax ceiba* root powder were determined as per WHO guidelines (WHO, 1999).

Qualitative phytochemical evaluation

Extraction

The coarse powder of shade-dried plant were successively extracted with petroleum ether, chloroform, methanol, water:ethanol (30:70) and water. The extracts obtained were concentrated and dried under vacuum.

Chemical tests

All plant extracts were subjected to chemical tests for the presence of following phytochemical classes like carbohydrates, alkaloids, anthraquinones, saponin glycosides, phytosterols, pheno-

lics, tannins, flavonoids, proteins and amino acids using reported methods (WHO, 1999).

Thin layer chromatographic studies

The hydro-alcoholic extract obtained was subjected to thin layer chromatographic studies using reported methods to determine the presence of various phyto-constituents (Stahl, 2000). The results were compared with the results obtained in the qualitative tests. The mobile phases and detecting reagents of different classes of compounds were used (Wagner and Bladt, 2003).

High-performance thin layer chromatography

A Camag microlitre sample (Hamilton, Bonaduz, Switzerland) syringe was used for a sample application on pre-coated silica gel aluminium plate 60F-254 (5 cm x 10 cm with 0.2 mm thickness (E. Merck, Darmstadt, Germany) using a Camag Linomat-V (Switzerland). Samples were then separated using solvent systems cyclohexane:diethyl ether:ethyl acetate of different compositions. Densitometric scanning was performed on Camag TLC scanner III in the reflectance-absorbance mode for all measurements and operated by CATS software (V1.4.3 Camag). The plate was scanned at 254 and 366 nm before and after sparging. The detecting reagent used was anisaldehyde sulphuric acid reagent (Wagner and Bladt, 2003). Reference standard lupeol was also applied on pre-coated HPTLC plates, and its presence in *Bombax ceiba* root extracts was confirmed.

The samples were prepared by dissolving 500 mg hydro-alcoholic extract of *Bombax ceiba* root powder in 5 mL distilled water.

All the studies were performed three times.

RESULTS AND DISCUSSION

The morphological studies of roots of *Bombax ceiba* show the shape of the root is more or less cylindrical, slightly tapering. Color of the unpeeled root is yellowish brown to dark brown while a peeled root is pale yellow with a rough surface. All the organoleptic features studied have been summarized in Table 1.

Table 1. Organoleptic features of *Bombax ceiba* roots.

Features	Observations
Shape	Cylindrical
Width	1-5 cm
Length	20-50 cm
Color	Peeled-pale yellow, unpeeled-yellowish brown to dark brown.
Odor	Faint and characteristic
Taste	Characteristic, free from bitterness.

In transverse section (Fig. 2), roots are characterized by the presence of thick brownish continuous eight to ten layers of tubular cells. Outer layers contain reddish brown amorphous cells, and inner layers show thick-walled cells. Single flattened layer of endodermis were converted with distinct medullary rays of multiseriate parenchymatous cells. The phloem was presence of thick walled oval shape cells of lignified parenchymatous sheath. Cambium was observed by double walled polygonal cells of xylem in horizontal tracks. Crystals of calcium oxalate also present with simple or compound starch grains. Small circular pith was seen with dark brown colored outer covering with intercellular spaces at center. Table 2 shows the results of micro-chemical tests on transverse sections of *Bombax ceiba*.

Powder analysis of *Bombax ceiba* root observed that polygonal cells of parenchyma pitted type of vessels, unicellular, bunches of fibers with yellow thick mass of tissues along with mesh-like structure. Fig. 3 shows the presence of the cork cells, fibers, calcium oxalate crystals and vascular bundles.

The fiber length was also determined and found to be $2.7 \times 33.75 \mu\text{m}$. This value can be used as an identifying characteristic for *Bombax ceiba* roots.

The proximate analysis revealed that extraction values in petroleum ether 9.6%, alcohol 19.2% and water 16.8% were observed. Loss on drying of *Bombax ceiba* roots powder was found to be 6.8%. Percentages of total ash, water-

soluble ash and acid insoluble ash were found to be 4.8, 4.4 and 0.5%, respectively. The phytochemical evaluation showed the presence of saponins, thus foaming index and hemolytic index were determined and found to be 142.9 and 500.0, respectively.

Successive solvent extraction values in various organic solvents were observed as with petroleum ether 1.1%, chloroform 2.2%, methanol 5.3% and water 15.0% in percentage yield.

The qualitative phytochemical evaluation by chemical tests of all the extracts obtained after successive extraction showed the presence of alkaloids, glycosides, flavonoids, saponins and tannins. Result is summarized in Table 3. TLC studies of the crude hydroalcoholic extract also showed the presence of similar constituents as shown in Table 4.

HPTLC fingerprinting studies was performed on the hydro-alcoholic extract of *Bombax ceiba* roots. The results obtained are depicted in Fig. 4. The presence of steroids was confirmed by use of detecting agent, anisaldehyde-sulphuric acid. Table 5 shows the comparative R_f values observed. The presence of lupeol in the hydro-alcoholic extracts was confirmed by HPTLC at R_f value ~ 0.26 (Fig. 5).

Through screening of the literature on *B. ceiba* shows that it is a traditional plant by ethnobotanical way. The crude extract and aqueous extracts of stem bark, root and leaf of *B. ceiba* have be screened for some pharmacological activities like hepatoprotective, antiangiogenic, analgesic and antioxidant, hypoglycemic, antimicrobial activity and cholinesterase activity (Versiani, 2007). Other parts of plants such as gum, seed and seed oil which are well documents to possess valuable medicinal properties are not explored for their biological potential (Versiani, 2007). In future study the isolated principle from *B. ceiba* needs to be evaluated in animal model and clinic trial to understand the molecular mechanism of action, in search of lead molecules from natural resources.




Figure 2: Transverse section of *Bombax ceiba* roots. *Cork* with eight to ten layers of tabular cells. *Phelloderm*: One to three layers of radially arranged parenchymatous cells containing stone cells. *Phloem fibers*: Thickened walls, Lignified parenchymatous sheath.

Medullary rays: Distinct, multiseriate parenchymatous cells. *Xylem fibers*: Thickened walls, parenchymatous sheath containing starch grains and calcium oxalate crystals. *Pith*: Parenchymatous cells with intercellular spaces.

Table 2. Micro-chemical tests performed on transverse section (TS) of *Bombax ceiba* roots

Test	Observation	Inference
TS + phloroglucinol + conc. HCl	Pink colour observed	Fibres present
TS + iodine	Blue coloured granules	Starch granules present
TS + methylene blue	Blue colored bundle of cells absent	Mucilage absent
TS + methyl red	Distinguished cork cells and cork cambium	Differentiate Cork, cork cambium
Sudan red III	Red coloured droplet absent	Fixed oil, fats absent
Without staining	Identified cells	Stone cells, calcium oxalate crystal present




Figure 3: Microscopy of powdered *Bombax ceiba* Linn. roots. (A) *Cork cell*: Brownish, thin-walled, wavy cells are present. (B) *Fibres*: Occur in groups and be associated with the vessels. (C) *Calcium Oxalate crystal*: Minute prism shaped crystal present as cell content and scattered. (D) *Vessels*: Fragments of lignified reticulately thickened, annual and spiral vessels.

Table 3. Preliminary phytochemical screening of *Bombax ceiba* root extracts.

Name of chemical test	Petroleum ether extract	Chloroform extract	Methanol extract	Water extract
Carbohydrates	-	-	+	+
Protein and amino acid	-	-	+	+
Fat and oils	+	-	-	-
Steroid	+	+	-	-
Glycosides				
Cardiac glycosides	-	-	-	-
Anthraquinone glycosides	-	-	+	+
Saponin glycosides	-	-	+	+
Flavonoids glycosides	-	-	+	+
Alkaloids	-	-	+	+
Tannins and phenolic compounds	-	-	+	+

Table 4. TLC profile for hydro-alcoholic extract of *Bombax ceiba* roots powder.

Groups	Mobile Phase	Detection	Rf Value
Anthraglycosides	Ethyl acetate: methanol: water (100:13.5:10)	Bornträger reagent	Absent
Bitter Principles	Ethyl acetate: methanol: water (77:15:8)	Vanillin-Sulphuric acid reagent	0.62
Alkaloids	Toluene: ethyl acetate: diethylamine (70:20:10)	Dragendorff reagent	0.78
Flavonoids	Ethyl acetate: formic acid: glacial acetic acid: water (100:11:11:26)	Aluminium chloride reagent	0.91
Saponins	Chloroform: glacial acetic acid: methanol: water (64:32:12:8)	Vanillin-sulphuric acid reagent	0.42
Essential oils	Toluene: ethyl acetate (93:7)	Vanillin-sulphuric acid reagent	Absent
Coumarins	Diethyl ether: toluene (1:1) saturated with 10% acetic acid	UV 365 nm	Absent
Steroid	Cyclohexane: diethyl ether: ethyl acetate (4:6:2.5)	Anisaldehyde-sulphuric acid	0.3, 0.85, 0.88



Figure 4: HPTLC fingerprinting profile of hydroalcoholic extracts of *Bombax ceiba* root powder.

MP 1 (Mobile Phase 1): Cyclohexane: diethyl ether: ethyl acetate (4:3:2.5)

MP 2 (Mobile Phase 2): Cyclohexane: diethyl ether: ethyl acetate (4:4:2.5)

MP 3 (Mobile Phase 3): Cyclohexane: diethyl ether: ethyl acetate (4:6:2.5)

Table 5. HPTLC fingerprinting results of hydro-alcoholic extract of *Bombax ceiba* root powder.

Spot No.	Rf values at 254 nm before spraying			Rf values at 366 nm before spraying		
	MP 1	MP 2	MP 3	MP 1	MP 2	MP 3
01	0.05	0.05	0.05	0.05	0.05	0.05
02	0.09	0.09	0.24	0.36	0.12	0.10
03	0.34	0.21	0.36	0.60	0.35	0.24
04	0.48	0.38	0.67	0.62	0.39	0.37
05	0.60	0.54	0.76	0.70	0.97	0.67
06	0.62	--	0.86	0.75	--	0.87
07	0.72	--	--	0.82	--	0.94
08	0.75	--	--	0.85	--	--
09	0.82	--	--	0.93	--	--
Spot No.	Rf values at 254 nm after spraying			Rf values at 366 nm after spraying		
	MP 1	MP 2	MP 3	MP 1	MP 2	MP 3
01	0.12	0.21	0.10	0.11	0.10	0.10
02	0.20	0.36	0.18	0.34	0.15	0.14
03	0.27	0.42	0.26	0.45	0.20	0.16
04	0.30	0.50	0.31	0.50	0.27	0.19
05	0.36	0.67	0.36	0.59	0.30	0.22
06	0.40	0.70	0.40	0.61	0.35	0.25
07	0.45	0.80	0.44	0.63	0.40	0.29
08	0.53	0.86	0.52	0.67	0.43	0.31
09	0.61	0.90	0.57	0.70	0.59	0.34
10	0.66	0.96	0.62	0.75	0.70	0.37
11	0.75	--	0.68	0.82	0.93	0.67
12	0.85	--	0.78	0.85	--	0.84
13	0.92	--	0.85	0.87	--	0.97

Mobile Phase 1 (MP 1): Cyclohexane: diethyl ether: ethyl acetate (4:3:2.5)

Mobile Phase 2 (MP 2): Cyclohexane: diethyl ether: ethyl acetate (4:4:2.5)

Mobile Phase 3 (MP 3): Cyclohexane: diethyl ether: ethyl acetate (4:6:2.5)




Figure 5. (A) HPTLC chromatograms of lupeol standard (Rf value ~ 0.26) and (B) Hydroalcoholic extracts of *Bombax ceiba* root.

CONCLUSIONS

The present study gives an account on the preliminary pharmacognostic and phytochemical screening of *Bombax ceiba* roots which may be useful in order to authenticate, standardize and avoid any adulteration. The diagnostic microscopical characters and physicochemical data reported in this paper will be helpful in the development of a monograph. TLC & HPTLC studies help to verify adulteration in quality control of crude extract. Further due to the presence of various phytochemicals which may be therapeutically active, pharmacological screening of roots of *Bombax ceiba* extracts can also be performed.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENT

This article is a part of Pankaj Chaudhary Master of Pharmacy thesis.

REFERENCES

- CRPA (2002) Demand Study for Selected Medicinal Plants, Volume II. New Delhi: Center for Research & Planning and Action (CRPA) for Ministry of Health & Family Welfare, GOI, Department of ISM&H & WHO.
- Faizi S, Ali M (1999) Shamimin: a new flavonol C-glycoside from leaves of *Bombax ceiba*. *Plant Medica* 65(4): 383-385.
- Gopal H, Gupta RK (1972) Chemical constituents of *Salmalia malabarica* Schott and Endl. flowers. *J Pharm Sci* 61(5): 807-808.
- Hamberger M, Marston A, Hostettmann K (1991) *Advances in Drug Research*. 20th ed. London: Academic Press.
- Khandelwal KR (2004) *Practical Pharmacognosy: Techniques and Experiments*. 11th edition. Pune: Nirali Prakashan. pp 220
- Kirtikar K, Basu B (1994) *Indian Medicinal Plants*. 2nd ed. Vol I. New Delhi. pp. 1401.
- Masood E (1997) Medicinal plants threatened by over-use. *Nature* 385 (6617):570.
- Mehra PN, Karnick CR (1968) Pharmacognostical studies on *Bombax ceiba* Linn. *Indian J Pharm* 30(12): 284.
- NISC (National Institute of Science Communication) (2001) *The Useful Plants of India*. New Delhi: CSIR (Center for Science& Industrial Research).
- Puckhaber LS, Stipanovic RD (2001) Revised structure for a sesquiterpene lactone from *Bombax malbaricum*. *J Nat Prod* 64(2): 260-261.
- Rai Chaudhuri HN (1971) Pharmacognostic studies on the stem bark of *Bombax ceiba* L. *Bull Bot Surv Bengal* 15: 37-38.
- Rajendra KC (2007) A brief introduction to Semal (*Bombax ceiba* Linn). http://www.forestrynepal.org/images/sim-al_present_report3.pdf [Consulted Aug 01, 2014].
- Shantha TR, Venkateshwarlu G, Indira A, Gopakumar K, Sridhar BN (2009) Pharmacognostical and preliminary phytochemical studies on the thorns of *Bombax ceiba* Linn. *Aryavaidyan* 22(2): 74-81.
- Stahl E (2000) *TLC-A Laboratory Handbook*. 2nd Edition. New York: International Student Edition.
- The Wealth of India (2003) *A dictionary of Indian raw materials & industrial products*. Vol II-B. New Delhi: CSIR.
- Versiani MA (2007) Studies on chemical constituents of *Bombax ceiba*. Ph D Thesis, H.E.J. Research Institute of Chemistry, University of Karachi, Pakistan.

Wagner H, Bladt S (2003) Plant drug analysis- A thin layer chromatography. 2nd edition. New York: Springer.
Wagner H, Farnsworth NR (1994) Economic and medicinal plants research, Vol. 6. London: Academic Press.

WHO (1999) Quality control methods for medicinal plant materials. Geneva.
