



Journal of Pharmacy & Pharmacognosy
Research

E-ISSN: 0719-4250

editor@jppres.com

Asociación de Académicos de Ciencias
Farmacéuticas de Antofagasta
Chile

Abubakar, Umar S.; Abdullahi, Sani; Ayuba, Victoria; Kaigama, Shettima; Halidu, Usman
S.; Ayuba, Muhammad K.

Medicinal plants used for the management of diabetes mellitus in Zaria, Kaduna state,
Nigeria

Journal of Pharmacy & Pharmacognosy Research, vol. 5, núm. 3, mayo-junio, 2017, pp.
156-164

Asociación de Académicos de Ciencias Farmacéuticas de Antofagasta
Antofagasta, Chile

Available in: <http://www.redalyc.org/articulo.oa?id=496053943002>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Medicinal plants used for the management of diabetes mellitus in Zaria, Kaduna state, Nigeria

[Plantas medicinales utilizadas para el manejo de la diabetes mellitus en Zaria, estado de Kaduna, Nigeria]

Umar S. Abubakar*, Sani Abdullahi, Victoria Ayuba, Shettima Kaigama, Usman S. Halidu, Muhammad K. Ayuba

Bioresources Development Centre, Kano. National Biotechnology Development Agency (NABDA), Abuja, Nigeria.

*E-mail: umarfarouk2003@yahoo.com

Abstract

Context: The knowledge of traditional uses of medicinal plants in the management of diabetes mellitus in Zaria is still intact with the traditional medicine practitioners.

Aims: To collect, identify and document medicinal plants used traditionally for the management of diabetes mellitus in Zaria city.

Methods: Open-ended informal interviews and general conversation were administered to twenty-two (22) traditional medicine practitioners (TMPs) and herb sellers. The data collected included the local names of plants, parts used, method of preparations and mode of administration.

Results: Twenty-six (26) species of plants belonging to eighteen (18) families were identified. Herbal remedies were mostly prepared from freshly collected plants, while decoction was the main method of preparation. Leaves and stem bark formed the major parts of plants for herbal preparations.

Conclusions: The study has documented different traditional practices used for the management of diabetes mellitus in the study area. Further studies should be conducted to evaluate the antidiabetic activity of plant species that have not yet been studied and to identify the bioactive compounds responsible for the antidiabetic activity as claimed by the traditional healers.

Keywords: decoction; diabetes mellitus; medicinal plants; traditional healers.

Resumen

Contexto: El conocimiento de los usos tradicionales de las plantas medicinales en el manejo de la diabetes mellitus en Zaria sigue intacto con los médicos tradicionales.

Objetivos: Colectar, identificar y documentar las plantas medicinales utilizadas tradicionalmente para el manejo de la diabetes mellitus en la ciudad de Zaria.

Métodos: Se realizaron entrevistas informales abiertas y conversaciones generales a veintidós (22) practicantes de medicina tradicional (TMPs) y vendedores de hierbas. Los datos recogidos incluyeron los nombres locales de plantas, partes utilizadas, método de preparación y modo de administración.

Resultados: Se identificaron veintiséis (26) especies de plantas pertenecientes a dieciocho (18) familias. Los remedios herbarios se prepararon sobre todo de las plantas recién recogidas, mientras que la decocción era el método principal de la preparación. Las hojas y la corteza del tallo formaron las partes principales de las plantas para las preparaciones herbarias.

Conclusiones: El estudio ha documentado diferentes prácticas tradicionales utilizadas para el manejo de la diabetes mellitus en el área de estudio. Se deben realizar estudios adicionales para evaluar la actividad antidiabética de las especies de plantas que aún no han sido estudiadas y para identificar los compuestos bioactivos responsables de la actividad antidiabética según lo reivindicado por los curanderos tradicionales.

Palabras Clave: curanderos tradicionales; decocción; diabetes mellitus; plantas medicinales.

ARTICLE INFO

Received | Recibido: November 3, 2016.

Received in revised form | Recibido en forma corregida: January 28, 2017.

Accepted | Aceptado: January 29, 2017.

Available Online | Publicado en Línea: February 5, 2017.

Declaration of interests | Declaración de Intereses: The authors declare no conflict of interest.

Funding | Financiación: The authors confirm that the project has no funding or grants.

Academic Editor | Editor Académico: Gabino Garrido.

INTRODUCTION

Diabetes mellitus has been identified as the most common endocrine disorder that currently affects 200 million people of the world's population (Wais et al., 2012). It is projected to rise to over 366 million in the year 2030 (Wild et al., 2004). Diabetes mellitus is characterized by hyperglycemia with alteration of carbohydrates, protein and fat metabolism, resulting from defects in insulin secretion or sensitivity of insulin to body cells or both (Hovens et al., 2005). Such alterations result in elevated blood glucose concentration, which causes acute complications like hyperglycemia and hypoglycemia and long-term complications in many organs, which may lead to increase the risk of atherosclerosis, renal failure, nerve damage, coronary heart disease and blindness resulting in increasing disability (ADA, 2009).

Diabetes mellitus can be classified on the basis of clinical presentation of the disorder, and this can be grouped into three major types; Type I diabetes, Type II diabetes and gestational diabetes mellitus (Velho and Froguel, 2002). Diabetes mellitus is associated with reduced quality of life and increased risk factors for mortality and morbidity (Upendra-Rao et al., 2000).

The search for natural products to cure diseases has received considerable attentions in which medicinal plants have been the most important source (Okwu, 2001). They are believed to be an essential source of new chemical substances with potential therapeutic effects (Kuhn and Winston, 2000), and due to the crucial role that plant-derived compounds have played in drug discovery, and development for the treatment of several diseases, the isolation of new bioactive compounds from medicinal plants based on ethnomedicinal data appears to be a very promising approach (Newman, 2008).

Traditional medicine products are playing greater roles in the lives of the people across the world in the face of the global upsurge of drug resistance, toxicity, adverse effects and increasing costs of synthetic products (Mbi and Bilikha, 1998).

In Nigeria, several thousands of plants have been claimed to possess medicinal properties and are employed in the treatment of many ailments (Iweala and Oludare, 2011). Many of these indigenous medicinal plants are used as spices and food plants and for medicinal purposes (Nwaogu et al., 2007). Presently, medicinal plants continue to play a significant role in the management of diabetes mellitus, especially in developing countries where many people do not have access to modern antidiabetic drugs (Acharya and Shrivastava, 2008). However, the knowledge of medicinal uses of plants in the management of diabetes mellitus is still intact with the traditional medicine practitioners, and this knowledge is either lost or passed to their children by the word of mouth. Therefore, there is need to collect and document this knowledge before such rich heritages are lost. Thus, the present work was aimed to collect and document medicinal plants used for the treatment diabetes mellitus in Zaria local government area, Kaduna state, Nigeria.

MATERIAL AND METHODS

Study area

Zaria is the second biggest city in Kaduna State, Nigeria. It is one of the seven original Hausa city-states and also the capital of the Hausa Kingdom of Zazzau emirate. According to the 2006 Nigerian census, Zaria has a population of 406, 990 people, total land area of about 563 km² and located on 11.09°N and 7.72°E coordinates. It is the home of many educational institutions like the prestigious Ahmadu Bello University, Barewa College, College of Aviation Technology, Nigerian Military School among others (Fig. 1) (Maiwada and Renne, 2007).

Ethnobotanical survey

The main data sources consisted of a series of informal interviews and general conversation administered on the traditional medicine practitioners and herb sellers. The interviews were done in Hausa language. The data collected included the local names of plants, parts of the plants used and method of preparations. Diabetes mellitus is gene-

rally known as “Ciwon Siga” in Hausa language, and that was the term used in this study. The plants were collected in their flowering and fruiting seasons. The voucher specimens were identified and authenticated by a Taxonomist, Mr. Tijjani A. Gawuna of Ethnobotany Unit of Bioresources Development Centre, Kano, and the scientific names were confirmed in “Hausa names for plants and trees” written by Roger Blench, Mallam Dendo 8, Guest Road Cambridge CB1 2AL, United Kingdom. The specimens were later deposited in the herbarium for preservation.

A traditional medicine practitioner was a person who was recognized by the community in which he lives as competent to provide health care by using plant, animal, mineral substances and certain other methods. On the other hand, a herb seller was a person who sells herbs and can also identify them but have little knowledge of their medicinal values.

Statistical analysis

Descriptive statistics such as percentages, pie chart, informant consensus factor (ICF) and Fidelity level (FL) were used in the analysis of the data.

ICF was calculated using the formula as follows:

$$ICF = \frac{Nur - Nt}{Nur - 1} \quad (1)$$

Where Nur = number of use citation and Nt = number of species used.

The fidelity value (FL) was calculated using the formula as follows:

$$FL = \frac{Ip}{Iu} \quad (2)$$

Where Ip = number of respondents who reported the utilization of medicinal plants for a specific main ailment, and Iu = total number of respondents who mentioned the same plant for any ailment.

RESULTS

Respondents identity

A total number of twenty-two (22) respondents were interviewed, with 25% within the age range of 25-35 years, 55% within the age range of 36-45 years, while 20% were within the age range of 46-60 years. Most of the respondents are married (80%), male (80%) and traditional medicine practitioners (80%). Also, the majority of the respondents were primary school leavers (45%), and most of them claimed that they inherited the knowledge from their forefathers. The result is summarized in Table 1.

Ethnobotanical survey

A total number of twenty-six (26) plants belonging to 18 families were described as being used for the management of diabetes mellitus in Zaria. The families Malvaceae, Amaryllidaceae, Fabaceae, Moraceae, Myrtaceae, Anarcadiaceae, Meliaceae, and Combretaceae were the most frequently mentioned families in this study, while leaves, stem bark, and roots were the most frequently used plant parts (Table 2). The informant consensus factor (ICF) has shown a high degree of consensus (0.87) among the respondents, while the fidelity level (FL) value revealed *Vernonia amygdalina* and *Mangifera indica* with high FL values of 82% and 64% respectively.



Figure 1. Map of Zaria city.

Table 1. Demographic characteristics of the respondents.

Variable	Specification	Percentage (%)
Age	25-35	25
	36-45	55
	46-60	20
Sex	Male	80
	Female	20
Marital status	Single	20
	Married	80
Practice	Traditional practitioners	80
	Herb sellers	20
Years of experience	5-15	25
	16-25	55
	26 and above	20
Method of treatment	Herbal alone	100
Tribe	Hausa	75
	Fulani	20
	Yoruba	5
Religion	Islam	90
	Christianity	7
	Pagan	3
Source of information	Ancestral	65
	Acquired	35
Educational status	Primary	45
	Secondary	35
	Tertiary	20

Table 2. Plants used for the treatment of diabetes mellitus in Zaria.

No.	Plant name	Family	Local name	Common name	Part used	Method of preparation	Mode of application	Voucher name
1	<i>Acacia nilotica</i>	Mimosaceae	Bagaruwa	Egyptian mimosa	Stem bark and leaves	Decoction	Oral	BDCKN/EB/1620
2	<i>Adonsonia digitata</i>	Malvaceae	Kuka	Boabab	Seeds	Infusion	Oral	BDCKN/EB/1626
3	<i>Allium cepa</i>	Amaryllidaceae	Albasa	Onion	Whole plant	Decoction	Oral	BDCKN/EB/1610
4	<i>Allium sativum</i>	Amaryllidaceae	Tafarnuwa	Garlic	Whole plant	Swallowing	Oral	BDCKN/EB/1619
5	<i>Anacardium occidentale</i>	Anacardiaceae	Kashu	Cashew	Leaves	Decoction	Oral	BDCKN/EB/1601
6	<i>Anogeissus leiocarpus</i>	Combretaceae	Marke	Chew stick tree	Stem bark	Decoction	Oral	BDCKN/EB/1607
7	<i>Azadirachta indica</i>	Meliaceae	Dogon yaro	Neem	Stem bark and leaves	Maceration	Oral	BDCKN/EB/1618
8	<i>Balanites aegyptiaca</i>	Balanitiaceae	Aduwa	Desert date tree	Stem bark and leaves	Maceration	Oral	BDCKN/EB/1602
9	<i>Brassica oleracea</i>	Brassicaceae	Kabeji	Cabbage	Leaves	Infusion	Oral	BDCKN/EB/1611
10	<i>Carica papaya</i>	Caricaceae	Gwanda	Pawpaw	Leaves	Decoction	Oral	BDCKN/EB/1605
11	<i>Citrus medica</i>	Rutaceae	Lemon tsami	Lemon	Fruit	Infusion	Oral	BDCKN/EB/1622
12	<i>Eugenia caryophyllus</i>	Myrtaceae	Kanunfari	Clove	Seeds	Decoction	Oral	BDCKN/EB/1613
13	<i>Ficus sycomorus</i>	Moraceae	Baure	Fig	Leaves and Stem bark	Decoction	Oral	BDCKN/EB/1621
14	<i>Ficus thonningii</i>	Moraceae	Chediya	Loin cloth fig	Leaves	Decoction	Oral	BDCKN/EB/1617
15	<i>Gossypium herbaceum</i>	Malvaceae	Auduga	Cotton	Leaves	Decoction	Oral	BDCKN/EB/1609
16	<i>Guiera senegalensis</i>	Combretaceae	Barbata	Sabara	Leaves and root	Decoction	Oral	BDCKN/EB/1616
17	<i>Khaya senegalensis</i>	Meliaceae	Madaci	Mahogany	Stem bark	Infusion	Oral	BDCKN/EB/1612

Table 2 (continued...)

No.	Plant name	Family	Local name	Common name	Part used	Method of preparation	Mode of application	Voucher name
18	<i>Lawsonia inermis</i>	Lythraceae	Lalle	Egyptian priest	Stem bark	Decoction	Oral	BDCKN/EB/1604
19	<i>Mangifera indica</i>	Anacardiaceae	Mangwaro	Mango	Leaves and stem bark	Decoction	Oral	BDCKN/EB/1625
20	<i>Moringa oleifera</i>	Moringaceae	Zogale	Drumstick	Leaves	Squeezing	Oral	BDCKN/EB/1623
21	<i>Parkia biglobosa</i>	Fabaceae	Dorawa	Locust bean	Fruits	Grinding	Oral	BDCKN/EB/1608
22	<i>Piliostigma reticulatum</i>	Fabaceae	Kalgo	Camel foot tree	Stem bark	Infusion	Oral	BDCKN/EB/1615
23	<i>Psidium guajava</i>	Myrtaceae	Goba	Guava	Leaves	Infusion	Oral	BDCKN/EB/1624
24	<i>Solanum incanum</i>	Solanaceae	Gauta	Bitter apple	Fruits	Eating fresh fruits	Oral	BDCKN/EB/1603
25	<i>Vernonia amygdalinia</i>	Asteraceae	Shiwaka	Bitter leaf	Leaves	Soup	Oral	BDCKN/EB/1614
26	<i>Ziziphus mucronata</i>	Rhamnaceae	Magaryar Kura	Buffalo thorn	Leaves	Decoction	Oral	BDCKN/EB/1606

N=22 Respondents

Characteristics of recipes used

Recipes were made mostly from combination of different parts of two plant species (Table 3), while some were made from a single plant part. Decoction is the most preferred method of

preparation, while oral administration was the only mode of administration. Leaves are the most used plant parts in the management of Diabetes mellitus in this study; this is followed by stem bark and fruits (Fig. 2).




Figure 2. Percentage occurrence of plant parts used for the management of diabetes mellitus in Zaria.

Table 3. Some antidiabetic recipes and mode of preparation

No.	Recipe	Method of preparation	Mode of administration
1	<i>Acacia nilotica</i> (leaves) + <i>Allium sativum</i> + Honey	Decoction	Oral
2	<i>Adonsonia digitata</i> (Powdered fruits) + Cow milk	Infusion	Oral
3	<i>Allium cepa</i> + <i>Brassica oleracea</i> (leaves)	Decoction	Oral
4	<i>Anarcadium occidentale</i> (leaves) + <i>Carica papaya</i> (brown leaves) + Red potash	Decoction	Oral
5	<i>Anogeissus leiocarpus</i> (Stem bark) + <i>Guiera senegalensis</i> (Leaves)	Decoction	Oral
6	<i>Azadirachta indica</i> (leaves) + <i>Vernonia amygdalina</i> (leaves)	Maceration	Oral
7	<i>Balanites aegyptiaca</i> + <i>Mangifera indica</i> (leaves)	Maceration	Oral
8	<i>Carica papaya</i> (leaves) + Red potash	Decoction	Oral
9	<i>Citrus medica</i> (fruits) + Honey	Decoction	Oral
10	<i>Eugenia caryophyllus</i> + <i>Citrus medica</i>	Decoction	Oral
11	<i>Ficus symorus</i> (leaves) + Red potash	Decoction	Oral
12	<i>Gossypium herbaceum</i> (leaves) + <i>Citrus medica</i> (fruits)	Decoction	Oral

N=12 Respondents

DISCUSSION

The present study has documented the ethnomedicinal management of diabetes mellitus in Zaria local government area of Kaduna State, Nigeria. The indigenous population still relies on medicinal plants to meet their healthcare needs because of the perceived effectiveness, presumed safety, moderate side effects, accessibility and affordability (Vliathan, 1998).

The claim made by most of the respondents that they inherited the knowledge of ethnomedicinal uses of plants from their forefathers further corroborates the fact that the knowledge of traditional medicine is passed from parents to their children and spouses (Enwere, 2009).

Some of the plant species identified in this study were also reported to be used by the indigenous people of Southwestern Nigeria, India, North Centre of Morocco and South Eastern Morocco for the same purpose. The plants include *Azadirachta indica*, *Carica papaya*, *Parkia biglobosa* and *Vernonia amygdalina* (Abo et al., 2008). Therefore, these plant species could be considered as promising candidates for further scientific validation in the search for new, effective and affordable anti-diabetic drugs.

The antidiabetic activity of some plant species reported in this study has been demonstrated using different diabetic models (*in vitro* and *in vivo*). The plants include *Acacia nilotica* (Mwangi et al., 2015), *Azadirachta indica* (Perez-Gutierrez and Domain-Guzman, 2012), *Carica papaya* (Oke, 1998), *Ficus sycomorus* (Mousa, 1994), *Khaya senegalensis* (Ibrahim and Islam, 2014) and *Psidium guajava* (Sabjan and Vinoji, 2012).

The leaves formed the most frequently used plant parts in the management of diabetes mellitus in the present study. The use of leaves could be attributed to easy availability and also due to the presence of high amount of chemicals compounds, which could be easily extracted and used in different forms (Imran et al., 2014). Leaves are the most commonly used plant parts by traditional healers (Ngbolua et al., 2014).

The combination of different plants and parts in the preparation of antidiabetic herbal remedy is common among the respondents, and they believed that some plants enhance the action of other herbs. Polyherbal therapies have the synergistic and agonistic/antagonistic pharmacological agents within themselves that work together in a dynamic way to produce therapeutic efficacy with minimum side effects (Tiwari and Rao, 2002; Alexandros, 2007; Ebong et al., 2008).

Most of the respondents preferred to use fresh plant parts. Previous studies had shown that there were quantitative and qualitative differences in the essential oil components of fresh and dry plant materials (Fateme et al., 2006; Okoh et al., 2008).

CONCLUSIONS

The study has documented different traditional practices used for the management of diabetes mellitus in Zaria. Further studies should be carried out to investigate the antidiabetic activity of other plant species that have not yet been studied, and also the bioactive compounds responsible for the antidiabetic activity need to be evaluated.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENT

The authors are indebted to the respondents who made this study possible. We also acknowledged the contribution of Mr. Abdulganiyu Abdu Yusuf, Head of ICT Unit, Bioresources Development Centre, Kano.

REFERENCES

- Abo KA, Fred-Jaiyesimi AA, Jaiyesimi AEA (2008) Ethnobotanical studies of medicinal plants used in the management of diabetes mellitus in south western Nigeria. *J Ethnopharmacol* 115: 67-71.
- ADA (2009) Standards of medical care in diabetes. *Diabetes Care* 39(suppl. 1): S13-S61.
- Acharya D, Shrivastava K (2008) Indigenous herbal medicines: tribal formulations and traditional herbal practices, 1st edn. Jaipur, India: Aavishkar Publishers, Distributors.
- Alexandros SB (2007) Plants used traditionally to treat malaria in Brazil: The archives of flora medicinal. *J Ethnobiol Ethnomed* 3: 18.
- Ebong PE, Atangwho JJ, Eyong EU, Egbung GE (2008) The antidiabetic efficacy of combined extracts from two continental plants: *Azadirachta indica* (A. Juss) (Neem) and *Vernonia amygdalina* (Del.) (African Bitter Leaf). *Am J Biochem Biotechnol* 4(3): 239-244.
- Enwere OO (2009) Herbs in orthodox practice: A view by medical students. *Afr J Tradit Complement Altern Med* 6: 203-206.
- Fateme S, Khadijeh A, Gholamreza BK (2006) Influence of drying and extraction methods on yield and chemical composition of the essential oil of *Satureja hortensis*. *Food Chem* 99: 19-23.
- Hovens MMC, Van de Laar FA, Cannegieter SC, Vandenbroucke JP (2005) Acetylsalicylic acid (aspirin) for primary prevention of cardiovascular disease in type 2 diabetes mellitus; metabolic and endocrine disorders group, In: The Cochrane library, Chichester, UK: John Wiley and Sons, Limited.
- Ibrahim MA, Islam MS (2014) Butanol fraction of *Khaya senegalensis* root modulates β -Cell function and ameliorates diabetes-related biochemical parameters in a type 2 diabetes rat model. *J Ethnopharmacol* 154 (3): 832-838.
- Imran K, Naser MA, Hassan F, Akash T, Riaz U, Muhammad A (2014) Application of ethnobotanical indices on the use of traditional medicines against common diseases. *Evid Based Complement Alternat Med* 2014(2014): 635371.
- Iweala EEJ, Oludare FD (2011) Hypoglycemic effect, biochemical and histological changes of *Spondias mombin* Linn and *Parinari polyandra* Benth. seed ethanolic extracts in alloxan- induced diabetic rats. *J Pharmacol Toxicol* 6: 101-112.
- Kuhn M, Winston D (2000) Herbal therapy and supplements: a scientific and traditional approach. New York: Lippincott and Wilkins, 347-350.
- Maiwada S, Renne EP (2007) New technologies of embroidered robe production and changing gender roles in Zaria, Nigeria, 1950-2005. *Textile History* 38(1): 25-28.
- Mbi CW, Bilikha JB (1998) Conventional drug production from medicinal plants with contribution from the Cameroon pharmacopoeia. Proceedings of the 1st International Workshop on Herbal Medicinal Products, Department of Pharmacognosy, University of Ibadan, Nigeria, November 22-24, pp 13-14.
- Mousa O, Vuorela P, Kiviranta I, AbdelWahab S, Hiltunen R, Vuorela H (1994) Bioactivity of certain Egyptian *Ficus species*. *J Ethnopharmacol* 41(1-2): 71-76.
- Mwangi JM, Ngugi MP, Njagi ENM, Njagi JM, Aguirio SD, Gathumbi KP, Muchugi NA (2015) Antidiabetic effects of aqueous leaf extracts of *Acacia nilotica* in alloxan induced diabetic mice. *J Diabetes Metab* 6: 568.
- Newman DJ (2008) Natural products as leads to potential drugs: An old process or the new hope for drug discovery? *J Med Chem* 51: 2589-2599.
- Ngbolua KN, Mpiana PT, Mudogo V, Ngombe NK, Tshibangu DST, Ekutsu E, Kabena ON, Gbolo BZ, Muanyishay CL (2014) Ethno-pharmacological survey and floristical study of some medicinal plants traditionally used to treat infectious and parasitic pathologies in the Democratic Republic of Congo. *Int J Med Plant* 106: 454-467.
- Nwaogu LA, Alisi CS, Ibegbulem CO, Igwe CU (2007) Phytochemical and antimicrobial activity of ethanolic extract of *Landoiphia owariensis* leaf. *Afr J Biotechnol* 6 (7): 890-893.
- Oke JM (1998) Antidiabetic potency of pawpaw. *Afr J Biomed Res* 1: 31-34.
- Okoh OO, Sadimenko AP, Asekun OT, Afolayan AJ (2008) The effects of drying on the chemical components of essential oils of *Calendula officinalis* L. *Afr J Biotechnol* 7(10): 1500-1502.
- Okwu DE (2001) Evaluation of the chemical composition of indigenous spices and flavouring agents. *Global J Pure Appl Sci* 7(3): 455-459.

- Perez-Gutierrez RM, Domain-Guzman M (2012) Meliacinolin: A potent α -glucosidase and α -amylase inhibitor isolated from *Azadirachta indica* leaves and *in vivo* antidiabetic property in streptozotocin-nicotinamide-induced type 2 diabetes in mice. *Biol Pharm Bull* 35: 1516-1524.
- Sabjan KB, Vinoji SK (2012) *In vitro* antidiabetic activity of *Psidium guajava* leaves extracts. *Asian Pac J Trop Dis* 2: 98-100.
- Tiwari AK, Rao JM (2002) Diabetes mellitus and multiple therapeutic approaches of phytochemicals: present status and future prospects. *Curr Sci* 83(1): 30-37.
- Upendra Rao M, Sreenivasulu M, Chengaiah B, Jaganmohan RK, Vats RK, Kumar V, Kothari A, Mital A, Uma R (2000) Emerging targets for diabetes. *Curr Sci* 88: 241-247.
- Velho G, Froguel P (2002) Maturity-onset diabetes of the young (MODY), MODY genes and non-insulin-dependent diabetes mellitus. *Diabetes* 23: 34-43.
- Vliathan MS (1998) Healing plants. *Curr Sci* 75: 1122-1126.
- Wais M, Nazish I, Samad A, Beg S, Abusufyan S, Ajaj SA, Aqil M (2012) Herbal drugs for diabetic treatment: an updated review of patents. *Recent Pat Antiinfect Drug Discov* 17: 53-59.
- Wild S, Roglic G, Green A, Scree R, King H (2004) Global prevalence of diabetes. Estimates for the year 2000 and projections for 2030. *Diabetes Care* 27(5): 1047-1053.

Author contribution:

Contribution	Abubakar US	Abdullahi S	Victoria A	Shettima K	Usman SH	Ayuba MK
Concepts or Ideas	X	X	X			
Design	X	X	X	X	X	X
Definition of intellectual content	X	X	X	X	X	X
Literature search	X		X	X		
Data acquisition	X	X			X	X
Data analysis	X	X	X	X	X	X
Statistical analysis	X	X	X	X	X	X
Manuscript preparation	X	X	X			
Manuscript editing					X	X
Manuscript review				X		

Citation Format: Abubakar US, Abdullahi S, Victoria A, Shettima K, Usman SH, Ayuba MK (2017) Medicinal plants used for the management of diabetes mellitus in Zaria, Kaduna state, Nigeria. *J Pharm Pharmacogn Res* 5(3): 156-164.