



Revista de Pesquisa Cuidado é
Fundamental Online

E-ISSN: 2175-5361

rev.fundamental@gmail.com

Universidade Federal do Estado do Rio
de Janeiro
Brasil

Pereira Borges, José Wicto; Magalhães Moreira, Thereza Maria; Pacheco Rodrigues,
Malvina Thaís; Saturnino de Oliveira, Andressa Suely; Braz da Silva, Daniele; Matias de
Santiago, Lindelvânia

HYPERTENSIVE PATIENTS WITH COMPLICATIONS REGISTERED AT HIPERDIA IN
FORTALEZA, CEARA: IMPLICATIONS FOR NURSING CARE

Revista de Pesquisa Cuidado é Fundamental Online, vol. 5, núm. 4, outubro-diciembre,
2013, pp. 556-565

Universidade Federal do Estado do Rio de Janeiro
Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=505750942015>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



RESEARCH

HYPERTENSIVE PATIENTS WITH COMPLICATIONS REGISTERED AT HIPERDIA IN FORTALEZA, CEARA: IMPLICATIONS FOR NURSING CARE
HIPERTENSOS COM COMPLICAÇÕES CADASTRADOS NO HIPERDIA DE FORTALEZA, CEARÁ: IMPLICAÇÕES PARA A ASSISTÊNCIA DE ENFERMAGEM
PACIENTES HIPERTENSOS CON COMPLICACIONES REGISTRADOS EN HIPERDIA, EN FORTALEZA, CEARÁ: IMPLICACIONES PARA LA ATENCIÓN DE ENFERMERÍA

José Wicto Pereira Borges¹, Thereza Maria Magalhães Moreira², Malvina Thaís Pacheco Rodrigues³, Andressa Suelly Saturnino de Oliveira⁴, Daniele Braz da Silva⁵, Lindelvânia Matias de Santiago⁶

ABSTRACT

Objective: To describe the sociodemographic and clinical characteristics of hypertensive complications associated with HIPERDIA registered in Fortaleza-Ceará, glimpsing possibilities of nursing care. **Method:** Cross-sectional quantitative runs from November 2008 to November 2010, with 187 people with hypertension and associated complications HIPERDIA registered in the municipality. The data were analyzed by software Predictive Analytics Software for Windows version 17.0. **Results:** In the results, 61% of women, elderly (66.3%), nonwhite (62.6%), low educational level (59.9%). Regarding clinical characteristics 47.6% and 71.6% had diastolic and systolic changed; 35.3% overweight, 38% stroke, 19.8% AML. **Conclusion:** Nurses should take ownership of this data source and assume his role as one of the professionals responsible for monitoring individual and group those users seeking a closer relationship of care with a view to minimize the complications of hypertension. **Descriptors:** Hypertension, Complications, Nursing.

RESUMO

Objetivo: Descrever as características sociodemográficas e clínicas dos hipertensos com complicações associadas cadastrados no HIPERDIA de Fortaleza-Ceará, vislumbrando possibilidades de cuidados de enfermagem. **Método:** Estudo transversal quantitativo, executado de novembro de 2008 a novembro de 2010, com 187 pessoas com HAS e complicações associadas cadastradas no HIPERDIA do município. Os dados foram analisados pelo software Predictive Analytics Software for Windows versão 17.0. **Resultados:** Nos resultados, 61% de mulheres, idosas (66,3%), não brancas (62,6%), com baixo nível escolar (59,9%). Em relação às características clínicas 47,6% e 71,6% tinham pressão diastólica e sistólica alteradas; 35,3% sobrepeso; 38% AVC; 19,8% IAM. **Conclusão:** O enfermeiro deve se apropriar dessa fonte de dados e assumir seu papel como um dos profissionais responsáveis pelo acompanhamento individual e coletivo desses usuários, buscando o estreitamento da relação de cuidado com vistas à minimizar as complicações da hipertensão arterial. **Descritores:** Hipertensão, Complicações, Cuidados de enfermagem.

RESUMEN

Objetivo: Describir las características sociodemográficas y clínicas de las complicaciones hipertensivas asociadas con HIPERDIA registrado en Fortaleza-Ceará, vislumbrando las posibilidades de atención de enfermería. **Método:** Estudio transversal carreras cuantitativos de noviembre 2008 a noviembre de 2010, con 187 personas con hipertensión y las complicaciones asociadas HIPERDIA empadronados en el municipio. Los datos fueron analizados mediante el software Predictive Analytics Software para Windows versión 17.0. **Resultados:** En los resultados, el 61% de las mujeres, los ancianos (66,3%), no blancos (62,6%), el bajo nivel educativo (59,9%). En cuanto a las características clínicas 47,6% y 71,6% tenían diastólica y sistólica cambiado; 35,3% trazo sobrepeso, el 38%, el 19,8% del AML. **Conclusión:** Las enfermeras deben tomar posesión de esta fuente de datos y asumir su papel como uno de los profesionales encargados del control individual y de grupo a aquellos usuarios que buscan una relación más estrecha de la atención con el fin de minimizar las complicaciones de la hipertensión. **Descriptor:** Hipertensión, Complicaciones, Enfermería.

¹Nurse, Master in Clinical Care in Health, State University of Ceará (UECE). Professor at the Federal University of Piauí - CAFS - Floriano, PI. Address: BR 343, km 3.5 - Neighborhood Meladão-Floriano/PI. CEP: 64800-000. E-mail: wictoborges@yahoo.com.br. ²Nurse, Post-Doctorate in Public Health. Deputy State University of Ceará. E-mail: tmmoreira@yahoo.com. ³Nurse, PhD in Public Health UECE / UFC / UNIFOR. Professor at the Federal University of Piauí. E-mail: malvinat@gmail.com. ⁴Nurse, Master in Clinical Health Care (UECE). Professor at the Federal University of Piauí. E-mail: andressasuely@hotmail.com. ⁵Nurse, Ph.D. in Public Health UECE / UFC / UNIFOR. Family Health Strategy Fortaleza - CE. E-mail: danibraz18@hotmail.com. ⁶Nurse, MS in Nutrition and Health (UECE). Family Health Strategy in Fortaleza, Ceará. E-mail: lm-santiago@bol.com.br. Project funded by CNPq.

INTRODUCTION

High blood pressure (HBP) is a serious public health problem because of its high prevalence in the world population and high mortality rates. It is a risk factor for cardiovascular disease, along with diabetes mellitus, smoking and dyslipidemia. Also, is primarily responsible for the high prevalence of cerebrovascular accidents (CVA) is associated with 40% of cases and 25% of cases of ischemic heart disease, the two leading causes of death in the isolated country.^{1, 2}

This scenario tends to accentuate with current demographic changes that happen in the world, coupled with increasingly intense spread of Western lifestyle, especially for an unhealthy diet and decreased physical activity.³

In this context, discusses the onus is generated by this chronic disease on the lives of affected subjects. The presence of this disease generates transformations in people's lives, including those related to self, justified by the possibility of aggravation in the long run, causing changes that require effort to accept and adapt to the new condition.⁴

Therefore, this condition requires establishment of a pharmacological and non-pharmacological treatment throughout his life, in order to obtain not only the control of hypertension, but also reduce cardiovascular morbidity and mortality and to improve quality of life.^{5, 6} Therefore, the attention to people with hypertension is aligned to strategies that aim to mobilize the meaning assigned to risky behavior, as well as the accession process to the proposed treatment, with their own needs, possibilities and expectations as reference.⁷

In this sense, the prevention and treatment of hypertension involve guidelines for introducing new lifestyle. A professional involved with this theme is the nurse who develops his actions by J. res.: fundam. care. online 2013. out./dez. 5(4):556-65

nursing consultation (EC) or in groups of health education. The relationship between the subjects strongly influences the continuity of care and is associated with greater adherence through a systematic approach, culminating in the prevention and / or control of blood pressure (BP).⁸

For nurses, the knowledge of their clientele carrier HAS enable the design of safe actions for the care and treatment adherence. Based on the foregoing, the study developed aimed to describe the sociodemographic and clinical characteristics of hypertensive complications associated with HIPERDIA registered at Fortaleza - Ceará, glimpsing possibilities of nursing care.

METHODOLOGY

It is cross-sectional, descriptive, quantitative, which integrates design umbrella Analysis of adherence to treatment of people with hypertension and associated complications, in Fortaleza, Ceará, funded by the National Council of Technological and Scientific Development (CNPq) and runs from November 2008 to November 2010.

The universe of the study corresponded to 14,200 registers users in HIPERDIA all Family Health Centers (CSF) from Fortaleza, whose information was tabulated in database software for Windows Predictive Analytics Software (PASW) version 17.0. Of this total, 1315 were tokens of complications associated with hypertension and were distributed in 71 of the 92 municipal CSF. Through formula for calculating finite sample population, there was obtained a sample equivalent to 415 people registered.

For the selection of the CSF that composed the study, the 75th percentile was traced, leaving 23 centers, located throughout the extension of the municipal territory and covering the six Regional Executive Secretariats. A sample of

Borges JWP, Moreira TMM, Rodrigues MTP et al.

Hypertensive patients with...

hypertensive patients with associated complications was drawn by random number table. With the collaboration of local health staff and community health workers were identified in the community, people raffled.

After identification, they organized workshops for sensitizing users to participate in the study. Almost all users invited did not attend the workshops, and only about 5.0% of them (20 people). Those users who did not attend the workshops and received home visits could also be included in the study. It is noteworthy that 118 people were excluded for the reasons described below: areas discovered by family health teams (37), addresses entered into the database were not found (81). With home visits and conducting workshops, other 110 users were excluded from the study, as reported not owning / ignore the complications listed in the register or any other complication associated with SH (56), worked all day and could not attend the interview (20), were unable to answer the questions due to dysarthria / mental disorder (10) were the death or after enrollment (24). Thus, in the end, 187 people with hypertension and associated complications registered in the county were located HIPERDIA, met the inclusion criteria of the study and agreed to participate.

Data were collected from June to September 2010, upon request of the users' consent to participate by signing an informed consent. The data collection instrument contained the following variables: sociodemographic (age, race, education and marital status), behavioral (sedentary, overweight / obesity and smoking), associated complications (left ventricular hypertrophy - LVH, heart failure - heart failure, acute myocardial infarction - MI, coronary artery disease - CAD, kidney disease and stroke).

We performed a descriptive analysis of absolute and relative frequencies concerning

sociodemographic, behavioral and associated complications. The study was approved by the Ethics in Research UECE with No. 08622921-4/2009 opinion. Ethical principles were followed in all phases of the study, in line with what advocates Resolution No. 196/96.

RESULTS AND DISCUSSION

The analysis of the demographic data of the 187 registered users who participated in the study showed that most were female (61.0%) of the corresponding age group for the elderly (66.3%), with non-white skin color (62.6%), with low education levels (one to eight years) (59.9%) and lived with a partner (60.9%), as shown in Table 1.

Table 1. Sociodemographic characteristics of users with hypertension and associated complications, registered in HIPERDIA (n = 187). Fortaleza, 2010.

Variables	F	%
Gender		
Male	73	39,0
Female	114	61,0
Age group		
< 60 years old	62	33,2
≥ 60	124	66,3
Not filled	1	0,5
Race/skin color		
White	68	36,4
Non white	117	62,6
Not filled	2	1,1
Schooling		
Cannot read/write	36	19,3
Up to 8 years of study	112	59,9
> 8 years of study	35	18,7
Not filled	4	2,1
Marital/family situation		
Single/divorced/widowed	73	39,1
Married/consensual union	114	60,9

Regarding the clinical characteristics of the users, it was possible to observe in Table 2 that, in relation to systolic blood pressure, the higher frequency was observed among patients with stage 1 hypertension (33.2%). Systolic pressure classified as normal was present in 16.6% of respondents. With regard to blood pressure,

Borges JWP, Moreira TMM, Rodrigues MTP et al.

Hypertensive patients with...

diastolic BP (DBP), 38.5% of users showed pressure cipher classified as normal while only 16.6% had systolic BP (SBP) normal. As for waist circumference (WC), we found that in males the most frequently corresponded with CA Normal users (19.3%), while nearly half of women (46.0%) had AC equal to or greater than 88 cm. The assessment of body mass index (BMI) indicated that 35.3% of users were overweight.

Table 2. Clinical characteristics of users with hypertension and associated complications, registered in HIPERDIA (n = 187). Fortaleza, 2010.

Variables	F	%
Systolic blood pressure *		
Great (< 120 mmHg)	22	11,8
Normal (< 130 mmHg)	31	16,6
Borderline (130-139 mmHg)	38	20,3
Stage 1 hypertension (140-159 mmHg)	62	33,2
Stage 2 hypertension (160-179 mmHg)	16	8,6
Stage 3 hypertension (≥ 180 mmHg)	16	8,6
Not filled	2	1,1
Diastolic blood pressure *		
Great (< 80 mmHg)	26	13,9
Normal (< 85 mmHg)	72	38,5
Borderline (85-89 mmHg)	7	3,7
Stage 1 hypertension (90-99 mmHg)	45	24,1
Stage 2 (100-109 mmHg)	22	11,8
Stage 3 hypertension (≥ 110 mm Hg)	13	7,0
Not filled	2	1,1
Abdominal circumference		
Male		
< 102 cm	36	19,3
≥ 102 cm	34	18,2
Female		
< 88 cm	26	13,9
≥ 88 cm	86	46,0
Not filled	5	2,7
Body mass index		
Normal (< 25 kg/m²)	51	27,3
Overweight (25 30-kg/m²)	66	35,3
Obesity (≥ 30 kg/m²)	46	24,6
Not filled	24	12,8

*Classification according to the VI Brazilian Guidelines about Hypertension (2010)

Concerning the complications associated with hypertension, the most frequent users was that the stroke (38.0%), followed by AMI (19.8%) and IC (9.6%). Other complications were reported by 19.8% of users.




Figure 1. Complications associated with hypertension reported by users interviewed (n = 187). Fortaleza, CE, 2010.

The higher frequency relative to female users was observed in other studies.⁹⁻¹³ This fact can be explained by the higher number of women enrolled in HIPERDIA certainly due to the greater women seek medical attention, especially because they usually have a higher perception of the disease and are more likely to self-care, increasing the likelihood of having hypertension diagnosed.¹⁴ However, the VI Brazilian Guidelines on Hypertension show the equivalence of global prevalence of hypertension among men and women, being higher in men up to 50 years, reversing from the fifth decade.¹

From the perspective of gender, nurses can use the potential demand for health services by women, using it as a mobilizing the male category for the care of SAH, sensitizing their peers for continuity of care.

Regarding the predominant age group corresponding to the elderly, shows confirmation that SAH is a disease that most commonly affects people aged 60 or more. The relationship between age and BP is direct and linear, with the prevalence of hypertension than 60.0% over 65 years.¹ In this matter, the professional must be aware of the continuity of care of the elderly, noting which supports care used by viewing the family as the focus of discussions and mobilization for the care of these individuals. The participation

Borges JWP, Moreira TMM, Rodrigues MTP et al.

Hypertensive patients with...

of family and other social systems (religious group, working) in treatment adherence is paramount.¹⁵

When viewing the family as the focus of nursing care is essential to establish this partnership with the healthcare team so that together form a unit of health care and that by taking part in actions with the possibility of transformation, family nursing can help to detect problems and needs, discuss the diagnosis, participate in determining the goals and collaborate on the planning application and evaluation.¹⁶

Thus, treatment of chronic disease should be a collaborative process between the patient, family, nurses and other health professionals. The collaboration is not limited to hospital settings it is important in all environments and throughout the illness trajectory.¹⁷ Skip to understand the care to hypertensive patients in a systemic context of relationships, is to glimpse the process of reciprocity and thus bilateral or dialogical care.⁸

As for the emphasis on non-white race, corroborates a study in Salvador-BA with 1,439 adults of which 74.3% were black or mixed race. The prevalence of hypertension in these individuals was 66.4% .¹⁸ It has been found that some ethnic groups are more exposed to complications of hypertension than others, for reasons not fully understood.¹⁹

Regarding the prevalence of low educational level, rescues that this is one of the determinants of adherence, given the low education could hinder the assimilation of guidelines dispensed by health professionals for people with hypertension, necessary to maintain or improve their welfare.²⁰

In this sense, the guidance to the patient and family is one of the most significant aspects of nursing care and can make a difference in the

ability of patients and their families to adapt to chronic health conditions. Educated and well-informed people in general are more concerned with their health and do what is necessary to keep it.¹⁷

Thus, the health education sessions developed by the nurse should possess educational methodologies that enable better understanding of such subjects as the use of recreational resources, cultural and group formation.

The process of care facing people with hypertension, especially with regard to the educational process, carries great burden with regard to communication in nurse-patient binomial, one of the crucial points for greater adherence to treatment of hypertension. Communication with all its nuances allows a dialectical exchange of knowledge, which denotes commands that when understood and mutually constructed by the transceiver allows a bilateral exchange ratio, indispensable for the building of a therapeutic plan.²¹

With a practical approach to therapeutic communication, nurses can use it in queries to couples in order to enhance the therapeutic relationship between the user and the professional, caring and bonding between the couple. The onset of a chronic disease requires significant decisions for changes in lifestyle, which should occur in concert with her partner and family.

Thus, marital status submitted by users of this study showed that most lived with a partner, an important factor for adherence to antihypertensive treatment, as the spouse, in previous research, was the person most mentioned as supportive to the treatment.²²

Having a spouse is often an indication of high level of social support available because caregivers often spouses living with the individual

Borges JWP, Moreira TMM, Rodrigues MTP et al.

Hypertensive patients with...

and provide commitment and closeness. Additionally, the spouse is usually the main support system of the person and although it is not known exactly what type of support provided (emotional, instrumental or functional), the companion plays crucial role in adherence to treatment, helping to reduce the risk hospitalization and mortality.²³

Concerning the value of the PA, it was observed that a significant percentage of people with hypertension and associated complications were with SBP and DBP changed. These data are quite relevant, since it is people monitoring of CSF in Fortaleza and they already have associated complications, which worries even more. It is known that the higher the value of BP, the greater the chance of onset of MI, heart failure, stroke and kidney disease.²⁴⁻²⁶

This result can be problematical to change our therapeutic scheme and thus, the nurse can trigger a multidisciplinary network for the design of a new approach that reduces these blood pressure levels.

With respect to BMI, classification, according to the World Health Organization (WHO) showed that most users were overweight / obese. With respect to AC, it is known that, according to National Cholesterol Education Program's Adult Treatment Panel III (NCEP-ATP), measuring the AC greater than 102 cm in men and greater than 88 cm in women corresponds to an increased risk of affection by cardiovascular diseases (CVD)²⁷. This study showed that women had higher frequency values above the recommended parameter CA. Therefore, the trader may think the search for devices in the community who work with physical activity and / or dietary guidelines. The formation of community groups to develop regular physical activity can be a focus of nursing care within primary care.

The VI Brazilian Guidelines on Hypertension, identified overweight as a risk factor for hypertension and both AC and BMI were strongly associated with the occurrence of hypertension, BMI being more influential in males.¹ It is recommended to maintain the ideal weight with a BMI less than 25 kg/m², increased physical activity and reduced calorie diet. Excess weight associated with fat accumulation in the mesenteric region, called central-type obesity, visceral or androgen is associated with increased risk of atherosclerotic disease²⁸⁻²⁹.

Regarding complications, there was a high frequency of cases of stroke among hypertensive evaluated. Stroke is a neurological syndrome common in adults and is a major cause of morbidity and mortality worldwide³⁰. This high frequency reveals a serious public health problem, because in Brazil, data from the National Health System (SUS) show that stroke (ischemic and hemorrhagic) represent a major cause of death with about 90 000 cases / year. Besides the high mortality, stroke is the leading cause of disability in adults, generating high costs to SUS.³¹

These issues demand the nurse inserted in CSF tracking these people through home visits, observing the demands of care they need, which will depend on the degree of disability acquired after stroke. These visits are important mapping and stratification of these subjects regarding the nutritional demands, hygiene, ambulation, communication, commitment and social support dermatological already involved in that care.

It is worth mentioning the use of community mapping methodologies which support nursing care. The choice of instruments like genogramming enables visualizing each family in its complexity and dynamism. In a study conducted mapping of families with patients with hypertension and diabetes through these instruments, the nuclear family observed the

Borges JWP, Moreira TMM, Rodrigues MTP et al.

Hypertensive patients with...

presence of comorbidities such as heart disease, stroke, AML, among others.³²

It is estimated that about 18 million people have a stroke in 2015 and approximately one third of these cases result in death. The analyzes indicate that, in the next decade, due to the progressive aging of the population and not the control of risk factors, mortality from stroke will increase 20.0% in developing countries and 10.0% in developed countries.³³

As for AML appeared to 19.8% with CAD and 7.4%, for a total of 27.2% involvement of the coronary arteries, which agrees with the literature that shows that hypertension is present in 30% of deaths from CHD. The follow-up care to those individuals who have been affected by this condition is essential because hypertension plays an important role in disease progression in the subject after AML, contributing to ventricular remodeling, heart failure and acceleration of atherosclerosis, and its strict control, thus the target of interest.³⁴

Based on the foregoing and considering that the nurse inserted in the CSF may prescribe medicines in the programs of the Ministry of Health, know about the pharmacology must permeate nursing actions aimed at prescribing and transcription of medicines in order to more accurately on the appropriateness therapy, always seeking multidisciplinary partnerships for the establishment of a safe treatment. One should also pay attention to aspects of nonpharmacological treatment, since these also interfere with BP control.

Thus, it shows the narrowing of the basic relationship of care between health professionals and the user HAS to minimize the complications and costs to the NHS and hypertensive provide a better quality of life. One should act, permanently, with actions aimed at acquiring knowledge that guarantees users a good treatment

J. res.: fundam. care. online 2013. out./dez. 5(4):556-65

adherence, this fundamental question in minimizing cardiovascular events.³

CONCLUSION

The analysis of the sociodemographic characteristics of hypertensive patients with associated complications, registered in HIPERDIA of Fortaleza, has shown that these are predominantly elderly women, non-white, low level of education, married or living with a partner in a consensual union.

In general computation, registered users, had hypertension with blood pressure changes, abdominal circumference above the range considered adequate and overweight. Regarding the complications associated with hypertension, the most frequent were stroke, AML and HF.

It was found that the HIPERDIA is a valuable tool to track users with SAH, being an important source to support nursing care in developing treatment plans that seek to minimize and reduce the complications resulting from not controlling blood pressure.

The nurse must take ownership of that data source and assume his role as one of the professionals responsible for monitoring these individual and collective users with hypertension and should make use of a greater number of educational strategies aimed at increasing knowledge of individuals affected with morbidity, and promote self-care and greater adherence to treatment.

In caring for these people, the nurse can use strategies such as the mobilization of the male category through sensitization of female users, since these are the most people registered. Considering the age and level of education of users, health education strategies, such as individual counseling or group proves essential for treatment compliance and therefore control the impact of complication associated with SH already acquired and prevention involvement by other (s).

Borges JWP, Moreira TMM, Rodrigues MTP et al.

Hypertensive patients with...

These strategies should include the family, because it is device direct support to people with chronic disease, due to the familiarity of everyday life and to share the difficulties of living with a disease.

REFERENCES

01. VI Diretrizes Brasileiras de Hipertensão. Sociedade Brasileira de Cardiologia. Arq Bras Cardiol. 2010;95(1 supl.1):1-51.
02. Soares GP, Brum JD, Oliveira GMM, Klein CH, Silva NAS. Mortalidade por todas as causas e por doenças cardiovasculares em três estados do Brasil, 1980 a 2006. Rev Panam Salud Publica [serial on the Internet]. 2010 Oct [cited 2011 Feb 06] ; 28(4): 258-266. Available from: http://www.scielo.org/scielo.php?pid=S1020-49892010001000004&script=sci_abstract&tlng=pt
03. Benseñor IM, Lotufo PA. A hipertensão arterial no contexto nacional e internacional. In.: Pierin AMG. Hipertensão Arterial: uma proposta para o cuidar. Barueri, SP: Manole, 2004.
04. Pierin AMG. Adesão ao tratamento: conceitos. In.: Nobre F, Pierin AMG, Mion JR D. Adesão ao tratamento: o grande desafio da hipertensão. São Paulo: Lemos Editorial, 2001.
05. Gravina CF, Rosa RF, Franken RA *et al.* Sociedade Brasileira de Cardiologia. II Diretrizes Brasileiras em cardiogeriatría. Arq Bras Cardiol. 2010;95(3, supl 2):1-112.
06. Souza ACC, Moreira TMM, Borges JWP, Andrade AM, Andrade MM, Almeida PC. Caracterização clínico-epidemiológica da clientela com crise hipertensiva atendida em um serviço de emergência de um hospital municipal de Fortaleza-CE. Reme - Rev. Min. Enferm. 2009; 13(1):13-18.
07. Ide CA. Atenção ao doente crônico: um sistema teórico-instrumental em ressignificação.

In.: Pierin AMG. Hipertensão Arterial: uma proposta para o cuidar. Barueri, SP: Manole, 2004.

08. Borges JWP. Hipertensão comunicada e hipertensão compreendida: estudo da comunicação de saberes e práticas de enfermagem em uma Unidade Básica de Saúde da Família; 2008. 118p. Monografia (graduação) - Curso de Graduação em Enfermagem. Faculdade Integrada da Grande Fortaleza; 2008.
09. Ferreira SRG, Moura EC, Malta DC, Sarno F. Frequência de hipertensão arterial e fatores associados: Brasil, 2006. Rev Saúde Pública. 2009;43(supl.2):98-106.
10. Castro RAA, Moncau JEC, Marcopito LF. Prevalência de hipertensão arterial sistêmica na cidade de Formiga. Arq Bras Cardiol. 2007;88(3):334-9.
11. Tavares DMS, Pereira GA, Iwamoto HH, Miranzzzi SSC, Rodrigues LR, Machado ARM. Incapacidade funcional entre idosos residentes em um município do interior de Minas Gerais. Texto Contexto Enferm. 2007;16(1): 32-9.
12. Ramos JN, Damacena LC, Stringhini MLF, Fornés NS. Perfil socioeconômico, antropométrico, bioquímico e estilo de vida de pacientes atendidos no programa “controle de peso”. Comun Ciênc Saúde. 2006;17(3):185-192
13. Teodósio MR, Freitas CLC, Santos NTV, Oliveira ECM. Hipertensão na mulher: estudo em mães de escolares de Jaboatão dos Guararapes - Pernambuco - Brasil. Rev Assoc Med Bras. 2004;50(2):158-62.
14. Zaitune MPA, Barros MBA, César CLG, Carandina L, Goldbaum M. Hipertensão arterial em idosos: prevalência, fatores associados e práticas de controle no Município de Campinas, São Paulo, Brasil. Cad. Saúde Pública. 2006;22(2):285-294.
15. Moreira TMM. Cuidando de pessoas com hipertensão arterial. In: Kalinowski CE (Org.). PROENF- Programas de Atualização em

Borges JWP, Moreira TMM, Rodrigues MTP et al.

Hypertensive patients with...

Enfermagem. Porto Alegre: Artmed/Panamericana, 2007; 2 (Mod 2):77-108.

16. Moreira TMM, Araújo TL, Pagliuca LMF. Alcance da Teoria de King junto a Famílias de pessoas portadoras de hipertensão arterial sistêmica. Rev gaúcha Enferm. 2001;22(1):74-89.

17. Smeltzer S, Bare BG. Brunner & Suddarth, Tratado de enfermagem médico-cirúrgica. 11ª Ed. Rio de Janeiro: Guanabara Koogan, 2009.

18. Lessa Í, Magalhães L, Araújo MJ, Almeida Filho N, Aquino E, Oliveira MMC. Hipertensão Arterial na População Adulta de Salvador (BA) - Brasil. Arq Bras Cardiol 2006;87(6):747-756.

19. Brandão AP, Brandão AA, Magalhães MEC, Pozzan R. Epidemiologia da hipertensão arterial. Rev Soc Cardiol Estado de São Paulo. 2003; 13 (1): 7-19.

20. Guedes NG et al. Crises hipertensivas em portadores de hipertensão arterial em tratamento ambulatorial. Rev. Esc. Enf. USP. 2005;39(2).

21. Borges JWP, Andrade AM, Souza ACC, Pinheiro NMG. Pragmática da Comunicação Humana: uma contribuição para a Consulta de Enfermagem frente à hipertensão arterial. Enfermagem Brasil. 2010; 9(3):56-62.

22. Moreira TMM, Araújo TL. Sistema interpessoal de Imogene King: as relações entre pacientes hipertensos não aderentes ao tratamento e profissionais de saúde. Acta Paul Enf. 2002;15(3):35-73.

23. Chung ML, Lennie TA, Riegel B, Wu JR, Dekker RL, Moser DK. Marital status as an independent predictor of event-free survival of patients with heart failure. Am J Crit Care 2009;18(6):562-70.

24. Luna RL. Hipertensão arterial: diagnóstico e tratamento. Rio de Janeiro: Revinter, 2009.

25. Fuchs SC. Fatores de risco para hipertensão arterial. In: Brandão AA, Amodeo C, Nobre F,

Fuchs FD. Hipertensão. Rio de Janeiro: Elsevier, 2006.

26. Ribeiro RC, Lotufo PA. Hipertensão Arterial: diagnóstico e tratamento. São Paulo: Ed. Sarvier, 2005. 117p.

27. World Health Organization. Obesity: Preventing and managing the global epidemic. Report of a WHO Consultation on Obesity. Geneva:WHO; 1998.

28. Sarno F, Monteiro CA. Importância relativa do Índice de Massa Corporal e da circunferência abdominal na predição da hipertensão arterial. Rev. Saúde Pública. 2007;41(5): 788-796.

29. Sociedade Brasileira de Cardiologia. IV Diretriz Brasileira sobre dislipidemias e prevenção da aterosclerose: Departamento de Aterosclerose da Sociedade Brasileira de Cardiologia. Arq. Bras. Cardiol. 2007;88(S1):2-19.

30. Giles MF, Rothwell PM. Measuring the prevalence of stroke. Neuroepidemiology 2008; 30:205-6.

31. André C, Curioni CC, Cunha CB, Veras R. Progressive decline in stroke mortality in Brazil from 1980 to 1982, 1990 to 1992, and 2000 to 2002. Stroke. 2006; 37(11):2784-789.

32. Pereira APS, Tixeira GM, Bressan CAB, Martini JG. O genograma e o ecomapa no cuidado de enfermagem em saúde da família. Rev Bras Enferm. 2009;62(3):407-16.

33. Strong K, Mathers C, Bonita R. Preventing stroke: saving lives around the world. Lancet Neurol. 2007;6(2):182-87.

34. III Diretrizes sobre o Tratamento do Infarto Agudo do Miocárdio. Arquivos Brasileiros de Cardiologia - Volume 83, Suplemento IV, Setembro 2004.

35. Florêncio RS. Análise do tratamento farmacológico anti-hipertensivo de idosos com hipertensão arterial e complicações associadas cadastrados no HIPERDIA de Fortaleza-Ceará.

J. res.: fundam. care. online 2013. out./dez. 5(4):556-65

Borges JWP, Moreira TMM, Rodrigues MTP et al.

Hypertensive patients with...

2011. 70p. Monografia (Graduação em Enfermagem) - Centro de Ciências da Saúde, Universidade Estadual do Ceará, Fortaleza; 2011.

Received on: 27/05/2012

Required for review: no

Approved on: 10/01/2013

Published on: 01/10/2013