



UNED Research Journal / Cuadernos de
Investigación UNED

ISSN: 1659-4266

cuadernosuned@gmail.com

Universidad Estatal a Distancia
Costa Rica

Iqbal Mir, Javaid; Kumar Sarkar, Uttam; Prakash Gusain, Om; Kumar Dwivedi, Arvind
Length-weight relationships of the Indian Major Carp, *Labeo rohita* (Cypriniformes:
Cyprinidae) from six drainages of Ganga basin, India

UNED Research Journal / Cuadernos de Investigación UNED, vol. 7, núm. 1, julio-
diciembre, 2015, pp. 71-77

Universidad Estatal a Distancia
San José, Costa Rica

Available in: <http://www.redalyc.org/articulo.oa?id=515651508009>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in [redalyc.org](http://www.redalyc.org)

[redalyc.org](http://www.redalyc.org)

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Length-weight relationships of the Indian Major Carp, *Labeo rohita* (Cypriniformes: Cyprinidae) from six drainages of Ganga basin, India

Javaid Iqbal Mir¹, Uttam Kumar Sarkar^{*2}, Om Prakash Gusain¹ & Arvind Kumar Dwivedi²

1. Department of Zoology and Biotechnology, Hemwati Nandan Bahuguna Garhwal University, Srinagar Garhwal, Uttarakhand, India, r.javaid@rediffmail.com, omprakashgusain@gmail.com
2. National Bureau of Fish Genetic Resources, Canal Ring Road, Dilkusha, Lucknow- 226002, India, usarkar1@rediffmail.com

* Correspondence

Received 12-VI-2014 • Corrected 02-IX-2014 • Accepted 22-X-2014

ABSTRACT: The Indian Major Carp, *Labeo rohita*, is a geographically widespread and economically important food fish species in tropical freshwater of India and adjacent countries. We studied the length-weight relationships of 1033 specimens collected from the main channel of Ganga river and its five major drainages from March 2009 to July 2012. The length of males ranged from 16 to 92cm (females: 16 to 94cm). The growth is allometric positive ($b>3$) for males, females and pooled sexes. The coefficient of determination (r^2) in males ranged from 0,978 to 0,989 and for females from 0,958 to 0,985. Data from field populations are scarce and our results will be useful in the management and conservation of *L. rohita* populations in its natural range.

Key words: LWR; Ganga basin; growth; wild; India.

RESUMEN: La carpa mayor de la India, *Labeo rohita* es una especie de distribución amplia y de importancia económica en las aguas dulces tropicales de India y países cercanos. El presente estudio trata las relaciones talla-peso de 1033 especímenes que fueron recolectados del canal principal del río Ganges y sus cinco principales drenajes, durante marzo 2009 y junio 2012. La longitud de los machos en diferentes ríos oscilaba entre 16 a 92cm y para las hembras entre 16 a 94cm. El crecimiento de esta especie fue alométrico positivo ($b>3$) para machos, hembras y sexos combinados. Además, la gama de coeficiente de determinación (r^2) en los hombres se registró desde 0,978 a 0,989 y para las mujeres que iba desde 0,958 a 0,985. Los datos generados en la presente investigación serán útiles para el biólogo pesquero y agencias de conservación en el manejo y conservación de las poblaciones silvestres de esta especie.

Palabras clave: relación talla-peso; río Ganges; crecimiento; silvestre; India.

The length-weight relationships (LWR) of fishes are important in fisheries biology because they allow the estimation of the average weight of the fish of a given length group by establishing a mathematical relation between the two (Mir, Mir, Patiyal & Kumar, 2014). Like any other morphometric characters, the LWR can be used as a character for the differentiation of taxonomic units and the relationship changes with the various developmental events in life such as metamorphosis, growth and onset of maturity (Thomas et al., 2003). Besides this, the LWR can also be used in setting yield equations for estimating the number of fish landed and comparing the population in space and time (Beverton & Holt, 1957). LWR parameters (a and b) are useful in fisheries science in many ways: to estimate weight of individual fish from its length, to calculate condition indices, to compare life

history and morphology of populations belonging to different regions (Petrakis & Stergiou, 1995) and to study ontogenetic allometric changes (Teixeira de Mello et al., 2006). Furthermore the empirical relationship between the length and weight of the fish enhances the knowledge of the natural history of commercially important fish species, thus making the conservation possible.

The Indian major carp, *Labeo rohita* (Hamilton, 1822), commonly known as 'Roho labeo' is a geographically widespread species in tropical freshwater of India and adjacent countries with considerable variation in growth parameters (Chondar, 1999). The existence of interpopulation differences may indicate local adaptations, either through phenotypic plasticity or genetic variability (Mir, Sarkar, Dwivedi, Gusain & Jena, 2014). Part of this interpopulation variation results from differences in local

environmental conditions (Begg, Cappo, Cameron, Boyle & Sellin, 1998). This species thrives well in lakes, ponds, rivers (Talwar & Jhingran, 1991), and is reported from brackish-water system also (Riede, 2004). A column feeder herbivore showing rapid growth in terms of flesh, Roho labeo is the most choice and prestigious culturable fishery in India and constitutes a main capture fishery of the Ganga, especially in the upper and lower stretches and tributaries. India is by far the largest producer of Roho labeo and the total global aquaculture production peaked in 2006, at nearly 16.5×10^5 tonnes (FAO, 2006-2012). The major source of Roho labeo seed in India is contributed via the Ganga basin (Chondar, 1999). The fish grows to a maximum total length of 200cm (Frimodt, 1995). Recent studies have shown fall in the catches of Roho labeo across Ganga basin in India (Mir, J. I., et al., 2014). To thwart this fall the generation of data on the growth condition of this species is an important task.

A number of reports are available on biological aspects of *L. rohita* from different water bodies (Varghese, 1973; Salam & Janjua, 1991; Sarkar, Negi & Deepak, 2006; Mir, Sarkar, Dwivedi, Gusain & Jena, 2013). The LWR studies on *L. rohita* are mostly restricted to cultured environments (Khan, 1972; Kamal, 1971; Sarkar, Medda, Ganguly & Basu, 1999). However, the information on LWR of *L.*

rohita is scanty from its wild habitat, especially from the rivers under this study. For the successful management and conservation of wild populations of Roho labeo, it is important to understand the relationship between length and weight of this fish species in its natural environment. Therefore, aim of the present investigation was to provide information on this relationship in Roho labeo from its wild habitat across the Ganga basin in India.

MATERIAL & METHODS

Study area: The Ganges river basin is the fifth largest basin in the world and is located $70-88^{\circ}30'E$ and $22^{\circ}-31^{\circ}N$. The length of main channel from the traditional source of the Gangotri Glacier in India is 2550Km. Annual volume of water discharged by the Ganges is the fifth highest in the world, with a mean discharge rate of $18.7 \times 10^3 m^3 sec^{-1}$ (Welcomme, 1985). The main sources of water in the basin are direct seasonal rainfall, mainly from the south west and glacial and snowmelt during the summer (Chapman, 1995). The main channel of the Ganges begins at the confluence of Bhagirathi and Alaknanda, which descend from upper Himalayas to Devprayag (520m above sea level) and receives a number of major tributaries (Fig. 1). The northern tributaries principally include




Fig. 1. Collection sites of *L. rohita* from three Indian rivers (Source Khan et al. (2013) with slight modifications).

Ghagra, Gomti, Buri Gandak and Kosi and southern tributaries include Yamuna (Ken, Betwa), Son, Chambal, Tons, Kalisindh, Sharda and Damodar. Most of these tributaries are controlled by irrigation barrages and there are two major barrages across the main channel, one at Hardwar which abstracts much of the water at this point to irrigate the Doab region and one at Farakka which diverts water down to Calcutta (Mir et al., 2013). All of these structures modify the flow of the river and may considerably influence fish distribution. Fish and fisheries are both important resource and activity in their own right but also provide indicators of the overall impact of anthropogenic changes over the basin.

Sample collection: A total of 1 033 specimens of *L. rohita* were collected during March 2009 to July 2012 from six rivers of the Ganga basin in India (Fig. 1). The sites on selected rivers, GPS coordinates, land use pattern of the selected sites on these rivers are shown in Table 1. The samples were collected from the rivers (upstream, midstream and downstream) in the morning before the sunrise, by using cast nets (9m length, 9m breadth and 1/2cm mesh size) and drag nets (100m length, 20m breadth, 1/2cm mesh size), and also from the 18 landing centers present on these six rivers. Measurements on fish length and weight were taken; besides, fish were dissected and sexes identified at the respective collection sites. Total length of each fish was taken from the tip of snout (mouth closed) to the extended tip of the caudal fin to the nearest 0,01mm by digital caliper and ruler and weighed to the nearest 0,01g (total weight) (by digital weighing machine, ACCULAB Sartorius Group).

Length-weight relationships: The relationships between length and weight of fish was analyzed by measuring length and weight of fish specimens collected from the study areas. The statistical relationship between

these parameters of fishes was established by using the parabolic equation by Froese (2006).

$$W = aL^b$$

where, W=weight of fish in grams, L=length of fish in mm, *a*=constant and *b*=an exponential expressing relationship between length-weight.

The relationship ($W = aL^b$) when converted into the logarithmic form gives a straight line relationship graphically.

$$\text{Log } W = \text{Log } a + b \text{ Log } L$$

where *b* represents the slope of the line, Log *a* is a constant. Additionally, 95% confidence limits of *b* and the coefficient of determination r^2 were estimated.

RESULTS

The regression parameters for length-weight relationships, coefficient of determination (r^2), 95% confidence interval of *a* and *b*, total length and body weight range in males, females and combined sexes is shown in table 2. The linear regressions of different populations were highly significant ($P < 0,001$). The regression coefficient (*b*) of LWRs of six rivers for this species was within the normal range of 2,5-3,5, as suggested by Froese (2006). A positive allometric growth ($b > 3$) was observed.

The overall value of r^2 ranged from 0,958 in females of river Betwa to 0,989 in males of river Ghagra. All the populations exhibited positively allometric growth irrespective of sex ($b > 3$; Fig. 2).

TABLE 1
GPS coordinates and land use pattern of different collection sites of *Labeo rohita* across Ganga river basin, India

Rivers (Sites)	Latitude (°N)	Longitude (°E)	Land use pattern
Ganga (Narora)	28°12'12"	78°23'33"	Atomic power plant, Dams, temples, Semi-urban, agriculture, domestic sewage
Ghagra (Faizabad)	25°45'55"	84°38'14"	Semi-urban, agriculture, domestic sewage
Betwa (Bhojpur)	24°08'54"	76°30'35"	Small dams, water lifting pumps, new road construction activities, industrial discharge, temples, rural, agriculture
Sharda (Palia)	22°49'47"	75°45'59"	Rural area, buffer zone (Preserved Area) agriculture activities, Forest
Ken (Patan)	23°17'04"	79°41'27"	Rural area, buffer zone (Preserved Area) agriculture activities, Forest
Gomti (Lucknow)	26°52'24"	80°55'42"	Urban, barrage, domestic sewage, beverage, distillery industry, temple in the river bank

TABLE 2
Total length and weight data, regression parameters and 95% confidence interval for *Labeo rohita*
in different drainages of Ganga river basin in India

Rivers	Sex	N	Total Length (cm)		Total body weight (g)		Regression parameters				
			Min	Max	Min	Max	a	95% CI of a	b	95% CI of b	R ²
Ganga	Male	104	16	90	40	10 250	-5,682	-5,876-5,489	3,288	3,216-3,360	0,983
	Female	85	30	92	300	11 750	-5,777	-5,991-5,562	3,325	3,245-3,405	0,980
	Overall	189	16	92	40	11 750	-5,718	-5,860-5,576	3,302	3,249-3,355	0,982
Ghagra	Male	70	30	80	364	8 600	-4,361	-4,535-4,146	3,145	3,015-3,304	0,989
	Female	77	16	85	50	11 900	-5,364	-5,546-5,224	3,211	3,126-3,353	0,977
	Overall	147	16	85	50	11 900	-5,766	-5,959-5,573	3,348	3,275-3,421	0,982
Betwa	Male	87	30	92	250	11 750	-6,263	-6,578-5,948	3,503	3,389-3,617	0,980
	Female	76	28	87	200	9 600	-6,245	-6,623-5,868	3,481	3,342-3,621	0,958
	Overall	163	28	92	200	11 750	-6,275	-6,502-6,047	3,500	3,417-3,584	0,977
Sharda	Male	89	28	85	250	9 000	-5,290	-5,587-4,992	3,155	3,044-3,266	0,978
	Female	71	29	94	250	11 500	-5,212	-5,424-4,999	3,123	3,044-3,203	0,985
	Overall	160	28	94	250	11 500	-5,305	-5,486-5,123	3,158	3,090-3,226	0,981
Ken	Male	102	31	71	250	4 500	-5,751	-5,988-5,514	3,310	3,221-3,399	0,980
	Female	95	16	87	40	9 600	-5,931	-6,174-5,687	3,386	3,296-3,477	0,985
	Overall	197	16	87	40	9 600	-5,890	-6,056-5,723	3,366	3,304-3,428	0,983
Gomti	Male	97	26	81	210	9 000	-5,473	-5,700-5,245	3,221	3,135-3,307	0,978
	Female	80	38	88	600	10 000	-6,064	-6,352-5,777	3,424	3,319-3,529	0,972
	Overall	177	26	88	210	10 000	-5,437	-5,622-5,252	3,201	3,132-3,270	0,972

DISCUSSION

The growth coefficient (b) of our study was in accordance with the findings of Jhingran (1952), Ahmed and Saha (1996), Sarkar et al. (1999) and Bhat (2011) for *Labeo rohita* from different man-made water bodies. Jhingran (1952) reported value of regression coefficient ($b = 3,0$) for *L. rohita* from a moat and spring; Ahmed and Saha (1996) reported regression coefficient of *L. rohita* as 3,14 from Kaptai lake, Bangladesh; Sarkar et al. (1999) has observed the b value for bundh and hatchery as 3,30 and 3,12 respectively from Calcutta and Bhat (2011) reported growth coefficient (b) equal to 2,97 and coefficient of determination (r^2) to be equal to 0,98 from Phuj reservoir, Jhansi. However, Choudhury, Kolekar and Chandra (1982) reported negative allometric growth ($b = 2.347$) from river Brahmaputra, Assam. According to Le Cren (1951), ecological conditions of the habitats or variation in the physiology of the animals, or both, are responsible for growth rate variations in the same species from different localities. In our study the growth condition is towards a declining trend from river to river. Similar results were obtained by Haniffa, Nagarajan and Gopalakrishnan (2006) in case of *Channa punctatus* from the rivers of Western Ghats. This provides evidence that the ecosystem might be the causative factor for the growth.

Even though the change in the b value depends primarily on the shape and fatness of the species, various factors may be responsible for the differences in the parameters of the length-weight relationships among seasons and years. According to Bagenal and Tesch (1978); Ozaydin, Uckun, Akalin, Leblebici and Tosunoglu (2007) and Mir, F.A. et al. (2014) the parameter b unlikely may vary seasonally, and even daily, and between habitats. Thus, the length-weight relationship in fish is affected by a number of factors including feeding, sex, maturity, specimen number, area, seasonal effects, degree of stomach fullness, habitat, health and general fish condition, differences in the observed length ranges of the specimen caught (Tesch, 1971; Khan, Miyan and Khan, 2013), only some of which were taken into account for the present study. Although plenty of work has been done on length-weight relationship of *L. rohita* from artificial water bodies like, ponds, moats etc., but the information available from the wild populations is scanty, especially for the rivers under this study.

This study is the first attempt to provide information about the growth condition of *L. rohita* from wild population of different geographical locations of river Ganga and its major tributaries. Our study will enlighten the




Fig. 2. Least-squares regression of $\log_{10} \text{W}$ vs $\log_{10} \text{L}$ of pooled samples of *L. rohita* in different drainages of Ganga river basin, India.

biologists about the status and growth of this fish in natural waters and this work will be useful for the fishery biologists and conservation agencies, for successful development, management, production and ultimate conservation of *L. rohita*.

ACKNOWLEDGEMENTS

The authors are grateful to the Director NBFGR, Lucknow, for providing necessary facilities. Thanks are due to the Indian Council of Agricultural Research, New Delhi for financial support to carry out this work.

REFERENCES

- Ahmed, K. K., & Saha, S. B. (1996). Length-weight relationships of major carps in Kapati Lake, Bangladesh. *Naga ICLARM Q*, 19, 22.
- Bagenal, T. B., & Tesch, F. W. (1978). Age and growth. In T. Bagenal (Ed.), *Methods for assessment of fish production in freshwater* (pp. 101–136). Oxford, UK: Blackwell Scientific Publication.
- Begg, G. A., Cappel, M., Cameron, D. S., Boyle, S., & Sellin, M. J. (1998). Stock discrimination of school mackerel (*Scomberomorus queenslandicus*) and spotted mackerel (*S. munroi*) in coastal waters of eastern Australia using analysis of minor and trace elements in whole otoliths.

Fisheries Bulletin, 96, 653-666. Retrieved from <http://research.mblwhoilibrary.org/works/21583>.

- Beverton, M. A., & Holt, S. J. (1957). On the dynamics of exploited fish populations, U.K. Ministry of Agriculture and Fisheries: *Fisheries Investigation Series*, 2, 533.
- Bhat, J. A. (2011). Length-weight relationship and condition factor of *Labeo rohita* (Cyprinidae) in Phuj reservoir, Jhansi, U.P., India. *Journal of Experimental Zoology, India*, 14, 339-344. Retrieved from http://www.connectjournals.com/file_html_pdf/943201H_A66.pdf
- Chapman, G. P. (1995). The Ganges plains. In G.P. Chapman & M. Thompson (Ed.), *Water and the quest for sustainable development in the Ganges Valley* (pp. 113-129). London: Mansfield Publishing.
- Chondar, S. L. (1999). *Biology of finfish and shellfish*. Howrah, India: SCSC Publisher.
- Choudhury, M., Kolekar, V., & Chandra, R. (1982). Length-weight relationship and relative condition factor of four Indian major carps of river Brahmaputra, Assam. *Journal of Inland Fisheries Society of India*, 14, 42-48.
- FAO. (2006-2012). Cultured Aquatic Species Information Programme. *Labeo rohita*. *Cultured Aquatic Species Information Programme*. Text by Jena, J.K. In FAO Fisheries and Aquaculture Department. Rome. Retrieved from http://www.fao.org/fishery/culturedspecies/Labeo_rohita/en.
- Frimodt, C. (1995). *Multilingual illustrated guide to the world's commercial warm-water fish*. Osney Mead, Oxford, England: Fishing News Books.
- Froese, R. (2006). Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22, 241-253. doi: 10.1111/j.1439-0426.2006.00805.x
- Haniffa, M. A., Nagarajan, M., & Gopalakrishnan, A. (2006). Length-weight relationships of *Channa punctata* (Bloch 1793) from Western Ghats Rivers of Tamil Nadu. *Journal of Applied Ichthyology*, 22, 308-309. doi:10.1111/j.1439-0426.2006.00779.x
- Jhingran, V. G. (1952). General length weight relationship of three major carps of India. *Proceedings of National Institute of Science India*, 18, 449-460.
- Kamal, M. Y. (1971). Length weight relationship of *Cirrhina mrigala* (Ham.) from commercial catches at Allahabad. *Proceedings of National Institute of Science India*, 41, 419-21.
- Khan, H. A. (1972). *Studies on the biology of some important major carps*. Thesis, Department of Zoology, Aligarh Muslim University, Aligarh.
- Khan, M. A., Miyan, K., & Khan, S. (2013). Morphometric variation of snakehead fish, *Channa punctatus*, populations from three rivers. *Journal of Applied Ichthyology*, 29, 637-642. doi: 10.1111/j.1439-0426.2012.02058.x
- LeCren, E. D. (1951). The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *Animal Ecology*, 20, 201-219. Retrieved from <http://www.jstor.org/stable/1540>.
- Mir, F. A., Mir, J. I., Patiyal, R. S., & Kumar, P. (2014). Length-weight relationships of four snowtrout species from the Kashmir Valley in India. *Journal of Applied Ichthyology*, 30(5), 1103-1104. doi:10.1111/jai.12482.
- Mir, J. I., Sarkar, U. K., Dwivedi, A. K., Gusain, O. P., & Jena, J. K. (2013). Age and growth in the Indian major carp *Labeo rohita* (Cypriniformes: Cyprinidae) from tropical rivers of Ganga basin, India. *Revista de Biologia Tropical*, 61, 1955-1966. Retrieved from <http://revistas.ucr.ac.cr/index.php/rbt/article/viewFile/12871/12128>.
- Mir, J. I., Sarkar, U. K., Dwivedi, A. K., Gusain, O. P., & Jena, J. K. (2014). Comparative Pattern of Reproductive Traits in *Labeo rohita* (Hamilton 1822) from Six Tropical Rivers of Ganges Basin: A New Insight. *Proceedings of National Academy of Sciences, India, Section B Biological Sciences*, 84, 91-103. doi: 10.1007/s40011-013-0181-y.
- Ozaydin, O., Uckun, D., Akalin, S., Leblebici, S., & Tosunoglu, Z. (2007). Length-weight relationships of fishes captured from Izmir Bay, Central Aegean Sea. *Journal of Applied Ichthyology*, 23, 695-696. doi: 10.1111/j.1439-0426.2007.00853.x
- Petrakis, G., & Stergiou, K. I. (1995). Weight-length relationships for 33 fish species in Greek waters. *Fisheries Research*, 21, 465-469. doi: 10.1016/0165-7836(94)00294-7
- Riede, K. (2004). *Global register of migratory species from global to regional scales*. Final Report of the R&D-Project 80805081. Federal Agency for Nature Conservation, Bonn, Germany.
- Salam, A., & Janjua, M. Y. (1991). Morphological studies in relation to body size of farmed Rohu, *Labeo rohita*. A cultureable major carp from Multan. *Journal of Research (Science)*, 3, 59-63.
- Sarkar, S. K., Medda, C., Ganguly, S., & Basu, T. K. (1999). Length-weight relationship and relative condition of Bundh and Hatchery-bred *Labeo rohita* (Ham.) during the early period of development. *Asian Fisheries Science*, 12, 289-296.
- Sarkar, U. K., Negi, R. S., & Deepak, P. K. (2006). Age Structure of Indian Carp *Labeo rohita* (Hamilton-Buchanan) from Different Wild Populations. *Environment & Ecology*, 24, 803-808.
- Talwar, P. K., & Jhingran, A. G. (1991). *Inland Fisheries of India and adjacent countries*. New Delhi, India: Oxford and IBH publishing co. Pvt. Ltd.

- Teixeira de Mello, F., Iglesias, C., Borthagaray, A. I., Mazzco, N., Vilches, J., Larrea, D., & Ballabio, R. (2006). Ontogenic allometric coefficient changes. Implications of diet shift and morphometric attributes in *Hoplias malabaricus* (Bloch) (Characiforme, Erythrinidae). *Journal of Fish Biology*, 69, 1770-1778.
- Tesch, F. W. (1971). Age and growth. In: *Methods for assessment of fish production in fresh waters*. In W.E. Ricker (Ed.), (pp. 98-130). Oxford, UK: Blackwell Scientific Publications.
- Thomas, J., Venus, S., & Kurup, B. M. (2003). Length- weight relationship of some deep-sea fish inhabiting continental slope beyond 250m depth along West coast of India. Naga. *World Fish Center Quarterly*, 26, 17-21.
- Varghese, T. J. (1973). The fecundity of the rohu, *Labeo rohita* (Hamilton). *Proceedings of Indian Academy of Science-Section B*, 77, 214-229.
- Welcomme, R. L. (1985). River Fisheries. *Fish Technical Paper 202* (330 pp). Rome: FAO.