

**Revista Mexicana de
Astronomía y Astrofísica**

Revista Mexicana de Astronomía y Astrofísica

ISSN: 0185-1101

rmaa@astroscu.unam.mx

Instituto de Astronomía

México

Medina, S.; Peña, M.

Expansion velocities from different ions of planetary nebulae with [wc]-type central stars

Revista Mexicana de Astronomía y Astrofísica, vol. 12, noviembre, 2002, p. 169

Instituto de Astronomía

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=57112066>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

EXPANSION VELOCITIES FROM DIFFERENT IONS OF PLANETARY NEBULAE WITH [WC]-TYPE CENTRAL STARS

S. Medina and M. Peña

Instituto de Astronomía, Universidad Nacional Autónoma de México, Apdo. Postal 70-264,
04510 México, D. F., México (selene, miriam@astroscu.unam.mx)

About 50 PNe in our Galaxy are known to have central stars of the carbon sequence Wolf-Rayet ([WC]) type stars and their atmospheres are almost pure helium and carbon. It has been suggested that the spectral sequence of [WC] stars and the “weak emission line stars” (WELS, as defined by Tytenda et al. 1993) corresponds to the following evolutionary sequence: late-[WC] \rightarrow early-[WC] \rightarrow WELS \rightarrow PG 1159 type stars. This suggestion is based upon both the analysis of the stellar atmospheres and the nebular properties of the planetary nebulae around [WC] stars (WRPNe) (e.g., Parthasarathy et al. 1998). This suggestion is controversial (Gorny & Tytenda 2000; Peña et al. 2001) and in order to study the nature and evolution of WRPNe and WELS, we have obtained a homogeneous high spectral resolution (echelle) data set of 30 planetary nebulae and their nuclei using the 2.1-m telescope at the Observatorio Astronómico Nacional, San Pedro Mártir, México.

High resolution is essential to estimate the expansion velocities of the nebular envelopes and in the present work we present preliminary results of the kinematic analysis of these objects from several nebular lines of different ionization degrees in order to map the kinematics along the ionization structure of the nebula.

We have analyzed the expansion velocities (V_{exp}) of different ionization species and we have found no systematic trends. In each object, high and low ionization species show the same V_{exp} . Although $V_{\text{exp}}(\text{H I})$ vs. $V_{\text{exp}}(\text{low ionization stages})$ diagrams show larger spreading than $V_{\text{exp}}(\text{H I})$ vs. $V(\text{high ionization stages})$ diagrams.

In general, expansion velocities show no correlation with nebular physical conditions (T_e , N_e , excitation class) nor with stellar [WC]-type. This latter result seems to un-favor the stellar evolutionary sequence previously proposed.

Another result is that no correlation has been found between the $[\text{O II}]/[\text{O III}]$ expansion velocity ratios and the nebular parameters.

A few objects, BD+30 3639, K2-16 and IC1747, show an anomalous behavior. In the cases of BD+30 3639 and K2-16, they show V_{exp} in [O III] higher than in H I by a factor of two, while IC1747 shows a higher V_{exp} in H I than in [O II]. A thorough study is needed to explain such anomalies.

REFERENCES

- Gorny, S. K., & Tytenda R. 2000, *A&A*, 362, 1008
- Parthasarathy, M., Acker A., & Stenholm, B. 1998, *A&A*, 329, L9
- Peña, M., Stasińska, G., & Medina, S. 2001, *A&A*, 367, 983
- Tytenda, R., Acker, A., & Stenholm, B. 1993, *A&AS*, 102, 595