**Revista Mexicana de
Astronomía y Astrofísica**

Revista Mexicana de Astronomía y Astrofísica

ISSN: 0185-1101

rmaa@astroscu.unam.mx

Instituto de Astronomía

México

Luridiana, V.; Peimbert, M.

The structure and star-formation history of ngc 5461

Revista Mexicana de Astronomía y Astrofísica, vol. 12, noviembre, 2002, p. 262

Instituto de Astronomía

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=57112107>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

THE STRUCTURE AND STAR-FORMATION HISTORY OF NGC 5461

V. Luridiana¹ and M. Peimbert¹

We compute photoionization models for the giant extragalactic H II region NGC 5461, and compare their predictions to several observational constraints. The models are constrained to reproduce the observed density profile, and our analysis takes into consideration the bias introduced by the shapes and sizes of the slits used by different observers.

The region can be satisfactorily described as radiation bounded, with no substantial leakage of photons taking place. We estimate a total stellar mass of about $M_*^{\text{tot}} = 3 \times 10^6 M_\odot$ in the $(0.01\text{--}80) M_\odot$ range, and an ionized-gas mass $M_{\text{gas}}^{\text{tot}} \geq 1.6 \times 10^6 M_\odot$. Accounting for the gas in neutral and molecular form, we find that the star-formation efficiency lies in the 3–5% range.

This work is discussed more fully in Luridiana & Peimbert (2001).

Fig. 1. Selected properties of our best-fit model. Upper panel: Ionization fractions of S^+ , S^{++} , and S^{3+} . Middle panel: $[S II] \lambda\lambda 6717, 6731$ emissivities as a function of radius. Lower panel: Density and filling factor as a function of radius.

Our results strongly depend on the assumed density law, since constant density models overestimate the hardness of the ionizing field, affecting the deduced properties of the stellar cluster. We find that an asymmetric nebula with a gaussian density distribution, powered by a young burst of 3.1 Myr, satisfactorily reproduces most of the constraints. Fig. 1 shows some of the features of this model, while Fig. 2 compares the predicted $H\alpha$ and $I(\lambda 6717)/I(\lambda 6731)$ profiles to their observational counterparts.

¹Instituto de Astronomía, UNAM, Apdo. Postal 70–264, 04510 México D. F., México (vale,peimbert@astroscu.unam.mx).

Fig. 2. Upper panel: $H\alpha$ brightness profile of the best-fit model (solid line), compared to the observed profile by Castañeda et al. (1992) (dot-dashed line). Lower panel: $I(\lambda 6717)/I(\lambda 6731)$ ratio profile of the best-fit model, superposed to the observational data by Castañeda et al. (1992) (asterisks).

REFERENCES

- Castañeda, H. O., Vilchez, J. M., & Copetti, M. V. F. 1992, A&A, 260, 370
Luridiana, V. & Peimbert, M. 2001, ApJ, 553, 633