

**Revista Mexicana de  
Astronomía y Astrofísica**

Revista Mexicana de Astronomía y Astrofísica

ISSN: 0185-1101

rmaa@astroscu.unam.mx

Instituto de Astronomía

México

Martínez-García, E. E.; González-Lópezlira, R. A.; Bruzual, G.  
TRIGGERED STAR FORMATION IN SPIRAL ARMS  
Revista Mexicana de Astronomía y Astrofísica, vol. 35, junio, 2009, pp. 231-232  
Instituto de Astronomía  
Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=57115758088>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

## TRIGGERED STAR FORMATION IN SPIRAL ARMS

E. E. Martínez-García,<sup>1</sup> R. A. González-Lópezlira,<sup>1</sup> and G. Bruzual<sup>2</sup>

### RESUMEN

Presentamos el estado actual de nuestra investigación, referente al desencadenamiento de formación estelar por choques galácticos a gran escala asociados a ondas de densidad espirales. Alrededor de una tercera parte de las galaxias de nuestra muestra parecen no ser adecuadas para esta clase de estudio, porque presentan un efecto, probablemente de opacidad, que no está bien entendido. Los objetos restantes parecen mostrar formación estelar inducida por ondas de densidad en los brazos. La comparación con modelos de síntesis de población estelar y las posiciones de las resonancias orbitales para estas galaxias (derivadas a partir de las velocidades angulares del patrón espiral), corroboran esta hipótesis.

### ABSTRACT

We present the status of our research, relative to the triggering of star formation by large-scale galactic shocks associated with spiral density waves. Around a third of the galaxies in our sample do not seem suitable for this kind of study, because they present an effect, probably due to opacity, that is not well understood. The remaining objects seem to favor the idea of density wave triggering of star formation in the arms. The comparison with stellar population synthesis models, and the orbital resonance positions for these galaxies (derived by means of spiral pattern angular speeds) corroborate this hypothesis.

*Key Words:* galaxies: spiral — stars: formation

### 1. GENERAL

After the density wave theory was established, the idea that spiral density waves were able to trigger star formation in the spiral arms of disk galaxies was proposed (Roberts 1969). Although this scenario may take place in real galaxies, the role and importance of this process is still an open question. González & Graham (1996) developed a method sensitive to the light of red and blue supergiants, using the reddening-free photometric index

$$Q(rJgi) = (r - J) - \frac{E(r - J)}{E(g - i)}(g - i). \quad (1)$$

Using this index, they were able to detect spiral density wave triggering of star formation in M99. Stellar population synthesis models (e.g. Bruzual & Charlot 2003) predict that the behavior of  $Q$  for a burst of star formation is well defined and can be observed for near face-on galaxies. This behavior is manifested in real galaxies as an azimuthal color gradient across spiral arms.

A sample of 31 galaxies with deep photometry was obtained during 1994–1996, in the optical filters

$g, r, i$ , and the infrared passbands  $J, K$ . A preliminary analysis (Martínez-García et al. 2007) of the data shows that azimuthal age gradients can be detected in other disk galaxies besides M99. A more careful inspection of the behavior of the  $Q$  index for the disks of this sample shows that in around a third of them the two halves of the disk present different mean values of  $Q$ . We believe that this effect may be due to the position of the dust lanes and stars with respect to the observer. More research needs to be done to understand the origin of this effect. So, in order to proceed with the original project, the galaxies with this “ $Q$  opacity effect” have been excluded. These galaxies were NGC 864, NGC 1703, NGC 3001, NGC 3059, NGC 3513, NGC 3338, NGC 4254, NGC 4593, NGC 4603, NGC 7083, and NGC 7217. The remaining galaxies were divided in two groups (see Table 1). The first one includes non barred galaxies types A and AB (de Vaucouleurs 1959), and the second B type barred galaxies. Preliminary results for the first group of galaxies indicate that some of them present azimuthal color gradients due to spiral density wave triggering of star formation. The candidate zones were compared with stellar population synthesis models (Bruzual & Charlot 2003). The relative positions of dust lanes (traced by the color  $g - J$ ), the density waves (indicated by the  $K$ -band data), and the age gradients (inferred from  $Q$ ) seem

<sup>1</sup>Centro de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México, Apdo. Postal 3-72, 58090, Morelia, Michoacán, México (e.martinez, r.gonzalez@astrosmo.unam.mx).

<sup>2</sup>Centro de Investigaciones de Astronomía, Apdo. Postal 264, Mérida 5101-A, Venezuela (bruzual@cida.ve).

TABLE 1  
GALAXY SAMPLE

	Name	Type
Group 1	NGC 578	SAB(rs)c
	NGC 918	SAB(rs)c
	NGC 7125	SAB(rs)c
	NGC 1421	SAB(rs)bc
	NGC 3162	SAB(rs)bc
	NGC 5371	SAB(rs)bc
	NGC 6951	SAB(rs)bc
	NGC 7753	SAB(rs)bc
	NGC 1417	SAB(rs)b
	NGC 718	SAB(s)a
	NGC 3938	SA(s)c
	NGC 7126	SA(rs)c
	NGC 4939	SA(s)bc
Group 2	NGC 7479	SB(s)c
	NGC 6907	SB(s)bc
	NGC 5383	SB(rs)b
	NGC 7496	SB(s)b
	NGC 266	SB(rs)ab
	NGC 986	SB(rs)ab
	NGC 4314	SB(rs)a

to be in accordance with theory.

By a simple approximation,

$$\Omega_p \cong \frac{1}{R} \left( \frac{v_{\text{app}}}{\cos(I.A.)} - v_{\text{rel}} \right) \quad (2)$$

(where  $R$  is the mean radius of the zone of study,  $v_{\text{app}}$  is the apparent velocity of rotation,  $I.A.$  is the

inclination angle, and  $v_{\text{rel}}$  is obtained from the fit of the model to the data) the pattern speed of the spiral wave can be obtained. The pattern speeds acquired this way must be consistent with the positions of major resonances (e.g., corotation, the outer and inner Lindblad resonances – OLR & ILR), if we are indeed observing an azimuthal age gradient as expected from theoretical considerations. The preliminary results for NGC 7125 show that the age gradient found in one of the arms of this galaxy yields a pattern speed  $\Omega_p \sim 20 \text{ km s}^{-1} \text{ kpc}^{-1}$ . With this value, the spiral pattern starts at the ILR and extends all the way to the OLR. Ongoing work is taking place to obtain global statistics for the A, AB & B galaxies in our sample. The color gradients will be compared with the latest stellar population synthesis models of Bruzual & Charlot (2003).

So far, the age gradients found in some of the galaxies of our sample confirm that such gradients are mainly observed in sites devoid of bright HII regions, as already observed by González & Graham (1996). HII regions may mask the gradients. The observed gradients indicate that spiral density wave triggering of star formation takes place for real galaxies and plays an important role for the cycling of the ISM in disk galaxies.

## REFERENCES

- Bruzual, G., & Charlot, S. 2003, MNRAS, 344, 1000  
 de Vaucouleurs, G. 1959, Handb. Phys., 53, 275  
 González, R. A., & Graham, J. R. 1996, ApJ, 460, 651  
 Martínez-García, E. E., González-Lópezlira, R. A., & Bruzual, G. 2007, IAU Symp. 237, Triggered Star Formation in a Turbulent ISM, ed. B. G. Elmegreen & J. Palous (Cambridge: Cambridge Univ. Press), 445  
 Roberts, W. W. 1969, ApJ, 158, 123