

**Revista Mexicana de
Astronomía y Astrofísica**

Revista Mexicana de Astronomía y Astrofísica

ISSN: 0185-1101

rmaa@astroscu.unam.mx

Instituto de Astronomía

México

Balucinska Church, M.; Church, M. J.; Halai, G.; Szostek, A.
Revealing the physical changes taking place in z-track Motion in low mass x-ray binaries
Revista Mexicana de Astronomía y Astrofísica, vol. 20, julio, 2004, p. 206
Instituto de Astronomía
Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=57120087>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

REVEALING THE PHYSICAL CHANGES TAKING PLACE IN Z-TRACK MOTION IN LOW MASS X-RAY BINARIES

M. Bałucińska-Church,^{1,2} M. J. Church,^{1,2} G. Halai,¹ and A. Szostek³

The explanation of the strong physical changes clearly taking place in the Z-track class of Low Mass X-ray Binaries has so far not been obtained, and this remains a significant astrophysical problem, without which we cannot claim to understand accretion in LMXB. We have for the first time applied the Birmingham emission model (2,3) to this problem to attempt to obtain a solution from the spectral evolution along the Z-track in the source GX 340+0 observed with *Rossi-XTE*. In this model, X-ray emission consists of blackbody from the neutron star, plus Comptonized emission from an extended ADC.




Fig. 1. Neutron star blackbody temperature and radius: evolution along the Z-track in GX 340+0

The results immediately demonstrate that strong changes take place on the neutron star and inner disk during Z-track movement. Starting from the one extreme end of the Z-track on the Horizontal Branch, the neutron star blackbody temperature kT_{BB} is initially high at the Eddington temperature, and the emitting area very small, indicating that radiation pressure effects are strong as we have found in other sources (1). However, along the Normal Branch, the source clearly recovers from these effects with kT_{BB} falling and the blackbody radius increasing R_{BB} to

the full radius of the neutron star. In flaring, the radius increases further due to the release of energy on the surface of the neutron star.




Fig. 2. Comparison of GX 340+0 with several Z-sources and the dipping, flaring source X 1624-490

In Fig. 2, we compare these results for R_{BB} in GX 340+0 with previous results for other Z-track sources and with the flaring in the dipping source X 1624-490, showing that flaring follows the same pattern as in Sco X-1, except that in Sco X-1, the maximum value of R_{BB} is 9.6 km consistent with the neutron star radius, whereas in GX 340+0, the radius increases to more than 20 km, as seen in a very small number of Galactic sources, implying that the emitter is a cloud of hot material enveloping the neutron star. The most interesting question is what drives the above recovery process. A clear possibility is that the mass accretion rate \dot{M} to the neutron star decreases on the Normal Branch; however, this contrasts with the assumption widely made that \dot{M} increases monotonically from the Horizontal to the Flaring Branch.

Acknowledgments: This work was supported in part by the Polish KBN grants PBZ-KBN-054/P03/2001 and KBN-1528/P03/2003/25

REFERENCES

1. Bałucińska-Church, M., Barnard, R., Church, M. J. & Smale, A. P. 2001, A&A, 378, 847
2. Church, M. J. & Bałucińska-Church, M. 2001, A&A, 369, 915
3. Church, M. J. & Bałucińska-Church, M. 2004, MN-RAS, 348, 955

¹School of Physics & Astronomy, University of Birmingham, Birmingham B15 2TT, UK (mbc@star.sr.bham.ac.uk).

²Astronomical Observatory, Jagiellonian University, ul. Orla 171, 30-244 Cracow, Poland.

³Copernicus Astronomical Centre, ul. Bartycka 18, 00-716 Warsaw, Poland.