

**Revista Mexicana de
Astronomía y Astrofísica**

Revista Mexicana de Astronomía y Astrofísica

ISSN: 0185-1101

rmaa@astroscu.unam.mx

Instituto de Astronomía

México

Gelino, Dawn M.

Measuring the masses of compact objects in low-mass X-ray binaries

Revista Mexicana de Astronomía y Astrofísica, vol. 20, julio, 2004, p. 214

Instituto de Astronomía

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=57120095>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

MEASURING THE MASSES OF COMPACT OBJECTS IN LOW-MASS X-RAY BINARIES

Dawn M. Gelino¹

Low-mass X-ray binaries (LMXBs) contain compact, black hole (BH) or neutron star (NS) primaries, and cool, low-mass secondary stars. We measure the orbital inclination of the system in quiescence by modeling infrared (IR) ellipsoidal variations from the secondary star in order to determine the compact object mass. I present our results for a few LMXBs, including the first BH that appears to conclusively fall in the $3 - 5 M_{\odot}$ range.

Compact object masses have significant consequences for the physics of supernova explosions, BH formation mechanisms, and even the expected event rate for gravitational radiation detectors. Even with these limited statistics (19 BH, 17 NS masses), there is a lack of BHs with masses between 1.4 and $5 M_{\odot}$. This observational finding does not agree with the detailed prediction of a continuous distribution with the largest number of BHs just above the NS mass limit, and decreasing for higher masses (Fryer & Kalogera 2001, ApJ, 554, 548).

Ellipsoidal variations result from the tidal and rotational distortions of the Roche lobe-filling secondary star, as well as its non-uniform surface brightness distribution. The amplitude of the ellipsoidal variations is a strong function of i , and can therefore be modeled to find the inclination of the system. IR variations are much less contaminated/diluted by accretion disk light than optical variations. It is important to accurately account for this extra light since dilution of the variations results in an underestimate of i , and an overestimate of M_1 . Fig. 1 shows KPNO 4 m IR light curves and WD98 models for the *faint* NS system, XTE J2123–058.

Table 1 presents some orbital inclination angles, compact object masses, and distances derived with our IR observing and modeling technique, while Table 2 presents the secondary star masses, orbital separations, and secondary star Roche lobe volume radii. XTE J2123–058 appears to harbor an overmassive neutron star, however, higher precision photometry and radial velocity measurements are needed to further constrain the mass of this faint




Fig. 1. XTE J2123–058 J -, H -, & K -band light curves (points) and WD98 models (lines). Error bars are 1σ . The data of this *very faint* ($J=20.12$ mag) source show no obvious signs of IR contamination. With $q = 0.36$ (Tomsick et al. 2002, ApJ, 581, 570), a χ^2 minimization finds the best fit model to have $i = 65 \pm 4^\circ$.

TABLE 1

IR DERIVED LMXB PARAMETERS

Object	i ($^\circ$)	M_1 (M_{\odot})	d (kpc)
XTE J2123–058 ^a	65 ± 4	1.71 ± 0.73	7.92 ± 2.21
GRO J0422+32 ^b	42 ± 2	3.97 ± 0.95	2.49 ± 0.30
GRS 1009–45 ^c	67 ± 3	5.24 ± 0.60	3.82 ± 0.27
GRS 1124–68 ^d	54 ± 1.5	7.24 ± 0.70	5.89 ± 0.26
GS 2000+25 ^e	64 ± 1.5	7.70 ± 0.66	2.29 ± 0.15
A0620–00 ^f	41 ± 3	11.0 ± 1.9	1.16 ± 0.11

^aThis work

^bGelino & Harrison 2003, ApJ, in Press

^cGelino & Harrison 2002, BAAS, 34, 654

^dGelino, Harrison, & McNamara 2001, AJ, 122, 971

^eGelino 2001, PhD Thesis, NMSU

^fGelino, Harrison, & Orosz 2001, AJ, 122, 2668

TABLE 2

IR DERIVED LMXB PARAMETERS^a

Object	M_2 (M_{\odot})	a (R_{\odot})	R_{L2} (R_{\odot})
XTE J2123–058	0.62 ± 0.59	2.20 ± 0.37	0.65 ± 0.18
GRO J0422+32	0.46 ± 0.31	2.45 ± 0.24	0.53 ± 0.16
GRS 1009–45	0.72 ± 0.16	3.30 ± 0.13	0.75 ± 0.06
GRS 1124–68	0.94 ± 0.20	4.85 ± 0.15	1.08 ± 0.05
GS 2000+25	0.32 ± 0.11	4.14 ± 0.13	0.66 ± 0.08
A0620–00	0.68 ± 0.18	4.47 ± 0.27	0.80 ± 0.07

^aReferences are the same as in Table 1.

system. GRO J0422+32 contains the lowest mass black hole measured, and the first to fall in the $3 - 5 M_{\odot}$ range. More accurate masses are needed to test the true distribution of BH masses.

¹Center for Astrophysics and Space Sciences, University of California San Diego, MSC 0424, 9500 Gilman Drive, La Jolla, CA 92093-0424 (dgelino@ucsd.edu).