

**Revista Mexicana de
Astronomía y Astrofísica**

Revista Mexicana de Astronomía y Astrofísica

ISSN: 0185-1101

rmaa@astroscu.unam.mx

Instituto de Astronomía

México

Dobrotka, A.; Hric, L.; Petřík, K.

The new view on the long-term flickering in t crb

Revista Mexicana de Astronomía y Astrofísica, vol. 20, julio, 2004, p. 235

Instituto de Astronomía

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=57120115>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

THE NEW VIEW ON THE LONG-THERM FLICKERING IN T CrB

A. Dobrotka,¹ L. Hric,² and K. Petřík³

The big scatter of data on the light curve of T CrB is identified with the flickering activity of the system. The data performed during April 1996 have a falling trend and they can be a part of a downward branch of a long-term and energetically powerful flare. Estimated energy $2 \cdot 10^{35}$ J and the duration of this event were compared with the theoretical assumptions based on three typical physical scenarios, which can be the source of flickering. The dissipation of magnetic loops and the existence of the turbulent eddies are energetically deficient, but these scenarios are real in the case of less powerful flares. The real explanation could be the instability of the secondary and variable mass transfer rate through the Lagrangian point, therefore through whole disc.

Our own material was performed at the observatory at Skalnaté pleso with 60 cm Cassegrain telescope with single channel photoelectric photometer using U, B, V filters (Hric et al. 1998).

The estimate of energy ($\sim 10^{35}$ J) of one probable long lasting flare was compared with the theoretical models of few flickering scenarios. We used the same algorithm as Bruch (1992), but in the case of turbulences we did not use constant diameter of turbulent eddies. In such a solution we obtained strong flares but with small duration and vice versa. We assumed the diameter of turbulent structures to be the high scale of the disc, therefore we obtained higher energies and durations of events (Fig. 1).

Another estimated and compared parameter was the time scale for the studied physical scenarios. We were searching for very long events (hours to days).

It is very difficult to work with energies because of more reasons. We do not know exactly what the continuum is, hence we can not identify precisely the start, the end and the amplitude of a flare. Our calculated energy is only in the observed inter-




Fig. 1. Energy of two physical scenarios as a function of distance from the white dwarf, mass transfer rate and for the magnetic loops as a function of magnetic flux density. Both are not able to produce energies $\sim 10^{35}$ J.

val UBV. Most of the radiation can be emitted in other wavelengths. The inner part of the disc is radiating in the UV and the boundary layer in the X-ray. The spectral distribution of emitted energy is a function of the distance from the central body, therefore we can not simply compare observationally estimated and theoretically calculated energy. The only one reliable parameter is the time scale. In this case the long-term brightness fluctuations can be identified as dissipation of magnetic loops, turbulences or variations in the mass transfer rate through the disc. If we take into consideration our estimated energy, the first two scenarios are deficient.

For identification of physical mechanism producing flickering we need observation in very wide spectral range. Combination of UBV, far UV and X data is required. The simple study of time scales is not inadequate.

We suggest, that similar algorithms could be used in the study of short-term variations in active galactic nuclei.

REFERENCES

- Bruch, A., 1992, A&A 266, 237
Hric, L., Petřík, K., Urban, Z., Niarchos, P., Anupama, G.C., 1998 A&A, 339, 449

¹Departement of Physics, Faculty of Materials Science and Technology, Slovak University of Technology in Bratislava, Paulínska 16, Trnava 917 24, The Slovak Republic(dobrotka@mtf.stuba.sk).

²Astronomical Institute of the Slovak Academy of Sciences, SK-059 60 Tatranská Lomnica, The Slovak Republic.

³Department of Physics, Pedagogical Faculty, University of Trnava, SK-918 43 Trnava, The Slovak Republic.