

**Revista Mexicana de
Astronomía y Astrofísica**

Revista Mexicana de Astronomía y Astrofísica

ISSN: 0185-1101

rmaa@astroscu.unam.mx

Instituto de Astronomía

México

Neill, James D.; Shara, Michael M.; Halbedel, Elaine; Malnushenko, Viktor

A comprehensive ha nova survey of m81

Revista Mexicana de Astronomía y Astrofísica, vol. 20, julio, 2004, p. 240

Instituto de Astronomía

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=57120120>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

A COMPREHENSIVE H α NOVA SURVEY OF M81

James D. Neill,^{1,2} Michael M. Shara,² Elaine Halbedel,¹ and Viktor Malnushenko¹

A spatially complete H α survey of M81 for novae was conducted continuously over a 5 month interval using the Calypso Telescope at Kitt Peak, AZ. A raw nova rate for M81 gives 23 yr^{-1} which is a lower limit. Monte Carlo simulations using nova light curves and survey frame limits yield a nova rate of $33_{-8}^{+13} \text{ yr}^{-1}$. Using this value and the K-band photometry for M81 from the 2MASS Large Galaxy Atlas of Jarret et al. (2003) gives a luminosity specific nova rate of $\rho_k = 3.96_{-1.1}^{+1.8} \text{ yr}^{-1} [10^{10} L_{\odot, K}]^{-1}$. The spatial distribution of the novae follows the bulge light much better than the disk or total light according to KS tests of their radial distribution. The asymmetric nova distribution across the major axis line of M81 implies a bulge-to-disk nova ratio of > 9 and supports the idea that novae originate primarily in older stellar populations.

The high nova rate we find for M81 implies that nova rates for other galaxies derived from surveys with incomplete spatial coverage and widely spaced epochs may be underestimates. Given the spatial distribution of novae in M81, most of the incompleteness arises in the central regions of the galaxies surveyed.

The asymmetry of the nova distribution apparent in Figure 1 (see Hatano et al., 1997) and the radial distributions shown in Figure 2 imply the predominance of bulge novae in M81. This adds weight to the idea that novae are associated with older stellar populations and calls for comprehensive surveys, like this one, of other galaxies to further test this idea. If verified, it would imply that novae don't appear in young stellar populations and would greatly constrain theories of nova formation and evolution.

REFERENCES

- Hatano, K., Branch, D., Fisher, A., & Starrfield, S. 1997, *ApJ*, 487, L45
 Jarrett, T. H., Chester, T., Cutri, R., Schneider, S. E., & Huchra, J. P. 2003, *AJ*, 125, 525
 Simien, F. & de Vaucouleurs, G. 1986, *ApJ*, 302, 564

¹Columbia University, 538 W. 120th St. MC C5247, New York, NY 10027, USA (neill@astro.columbia.edu).

²American Museum of Natural History.




Fig. 1. Spatial distribution of novae in M81




Fig. 2. Nova radial distribution and the bulge/disk decomposition of Simien & de Vaucouleurs (1986)