

**Revista Mexicana de
Astronomía y Astrofísica**

Revista Mexicana de Astronomía y Astrofísica

ISSN: 0185-1101

rmaa@astroscu.unam.mx

Instituto de Astronomía

México

Froning, C. S.; Long, K. S.; Szkody, P.; Gänsicke, B. T.
A fuse survey of disk-accreting cataclysmic variables
Revista Mexicana de Astronomía y Astrofísica, vol. 20, julio, 2004, p. 251
Instituto de Astronomía
Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=57120131>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

A FUSE SURVEY OF DISK-ACCRETING CATAclysmic VARIABLES

C. S. Froning,¹ K. S. Long,² P. Szkody,³ and B. T. Gänsicke⁴

We present initial results of a survey of the FUV spectra of disk-accreting cataclysmic variables obtained with the Far Ultraviolet Spectroscopic Explorer (FUSE).

FUSE covers the 905 – 1188 Å range at spectral resolutions $\simeq 0.1$ Å. To date, FUSE has observed more than 65 cataclysmic variables (CVs). Publicly-available data include observations of 11 dwarf novae (DN), 15 non-magnetic novalikes (NLs), 7 intermediate polars and DQ Her stars, at least 15 polars, and 4 super-soft X-ray binaries. The FUV spectra of CVs are highly complex, with emission from multiple continuum and spectral line source regions. We have instituted an FUV observational survey of CVs with FUSE. Combining the survey observations with data from targeted programs, we have created time-averaged spectra of all publicly-available FUSE spectra of disk-accreting CVs. The survey is on-going; here, we give an initial presentation of the characteristic FUV spectra of disk-accreting CVs.

Figure 1 shows example FUV spectra of two quiescent DN and two NLs. The spectrum of U Gem is populated by absorption lines of H I, He II, and ionized metals and is dominated by the WD. Several other quiescent DN, including SS Aur, Z Cam, VW Hyi, and WZ Sge, also show WD spectra (e.g., Froning et al. 2001, Long et al. 2003, Godon et al. 2003). This is not true of all DN, however, as the spectrum of EK TrA demonstrates. EK TrA, along with EM Cyg, SS Cyg, RU Peg, and SW UMa, have FUV spectra with relatively flat continua and strong, broad emission lines of OVI, CIII, and NIII. The underlying WD is not visible, and the source of the emission spectrum remains poorly understood.

Also shown in Figure 1 are typical FUV spectra of non-eclipsing NLs. RZ Gru has strong, broad, blueshifted OVI absorption. RX And and SS Cyg in outburst, IX Vel, and RW Sex show similar spec-




Fig. 1. Example FUV spectra of disk-accreting CVs. The two panels on the left show FUSE spectra of DN in quiescence: U Gem in the upper panel and EK TrA in the lower panel. The two panels on the right show two high-state NLs: RZ Gru in the upper panel and BP Lyn in the lower panel.

tra, which are believed to be dominated by continuum emission from a steady-state accretion disk and line absorption in a disk wind (e.g., Prinja et al. 2003). In contrast, numerous high state CVs, including U Gem in outburst, V795 Her, AH Men, LS Peg, V3885 Sgr, and IX Vel, show spectra similar to that of BP Lyn, with narrow, low-velocity absorption lines that are unlikely to originate in the inner accretion disk or in a disk wind (Froning et al. 2001). A vertically-extended chromosphere in the outer accretion disk is one possible site for the line absorption. Not shown here are spectra of eclipsing NLs and IPs. The eclipsing systems show strong, broad emission lines or a complex blend of spectral emission and absorption components (e.g., UX UMa; Froning et al. 2003). The IPs also have emission spectra, often superimposed on a blue continuum.

REFERENCES

- Froning, C. S., et al. 2003, ApJ, 584, 433
 Froning, C. S., et al. 2001, ApJ, 562, 963
 Godon, P., et al. 2003, ApJ, submitted
 Long, K. S., et al. 2003, ApJ, 591, 1172
 Prinja, R. K., et al. 2003, MN, 340, 551

¹CASA, University of Colorado, 389 UCB, Boulder, CO 80309 USA (cfroning@casa.colorado.edu).

²STSci, 3700 San Martin Dr., Baltimore, MD 21218 USA (long@stsci.edu).

³Astronomy Dept., Univ. of Washington, Seattle, WA 98195 USA (szkody@astro.washington.edu).

⁴Dept. of Physics and Astronomy, Univ. of Southampton, Highfield, Southampton SO17 1BJ UK (btg@astro.soton.ac.uk).