

**Revista Mexicana de
Astronomía y Astrofísica**

Revista Mexicana de Astronomía y Astrofísica

ISSN: 0185-1101

rmaa@astroscu.unam.mx

Instituto de Astronomía

México

Nunn, N.; Wickramasinghe, D. T.; Ferrario, L.
Towards self-consistent spectral models for outbursting Discs in am cvn binaries
Revista Mexicana de Astronomía y Astrofísica, vol. 20, julio, 2004, p. 255
Instituto de Astronomía
Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=57120135>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

TOWARDS SELF-CONSISTENT SPECTRAL MODELS FOR OUTBURSTING DISCS IN AM CVn BINARIES

N. Nunn,¹ D. T. Wickramasinghe,¹ and L. Ferrario¹

Models for AM CVn discs are usually constructed under the assumption of a steady state. Nevertheless, many of these systems have unstable discs. We have constructed models for the spectral evolution of such discs using a disc instability code (Cannizzo 1984) and detailed calculations of the vertical disc structure assuming hydrostatic equilibrium following El-Khoury and Wickramasinghe (2000).




Fig. 1. Low state accretion disc




Fig. 2. Transitional state accretion disc




Fig. 3. Outbursting accretion disc

For the present illustration, we consider a binary system of period 10 minutes with a white dwarf primary of mass $M_2 = 0.7M_\odot$ and a mass losing star of mass $M_1 = 0.066M_\odot$ - parameters which are possibly relevant for the short period AM Her system ES Cet. We show in Figures 1 to 3 snap-shots of the time dependent discs at three phases representing respectively low, transitional and high states of the disc. In each figure, the bottom panel gives the disc structure and the upper panel the integrated spectrum. We note the change in vertical disc structure as the heating front propagates through the disc. The spectrum changes from a pure emission line spectrum to an absorption line spectrum as the disc passes through an outburst cycle which is an observed characteristic of the sub-class of these systems which show unstable discs. It is generally not possible to mimic these spectra by constructing equivalent constant \dot{M} discs -that is, despite the compactness of the disc, no single annulus can adequately represent the integrated spectrum.

A detailed account of these calculations will be published in a forthcoming paper.

REFERENCES

- Cannizzo, J. K., 1984, *Nature*, 311, 443
 El Khoury, W. & Wickramasinghe, D. T., 2000, *A & A*, 358, 154

¹Dept. of Mathematics, The Australian National University, Canberra.