

**Revista Mexicana de
Astronomía y Astrofísica**

Revista Mexicana de Astronomía y Astrofísica

ISSN: 0185-1101

rmaa@astroscu.unam.mx

Instituto de Astronomía

México

Bulik, T.; Moderski, R.; Belczyński, K.

Constraints on stellar evolution from observations of Compact object binaries

Revista Mexicana de Astronomía y Astrofísica, vol. 20, julio, 2004, p. 266

Instituto de Astronomía

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=57120145>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

CONSTRAINTS ON STELLAR EVOLUTION FROM OBSERVATIONS OF COMPACT OBJECT BINARIES

T. Bulik,¹ R. Moderski,¹ and K. Belczyński²

The masses of compact object (black hole, neutron star) binaries depend strongly on the parameters describing stellar evolution. Such masses or their functions can be measured using gravitational waves or through microlensing searches. We analyze an example of the varying common envelope efficiency and show the dependence of distributions of the measured chirp masses in gravitational waves mass ratios through microlensing taking into account the relevant selection effects.

We use the StarTrack binary population synthesis code (Belczyński, Kalogera, Bulik, 2002) to calculate mass distributions of compact object binaries: double neutron stars (NSNS), black hole neutron star (BHNS), and double black hole binaries (BHBH). The mass distributions of compact object binaries depend on a number of parameters specified in the description of the StarTrack code. For the illustrative purpose we concentrate here on one parameter: the common envelope efficiency $\alpha_{CE\lambda}$ and discuss only the dependence of the results on this parameter.

Compact object binaries can be detected through gravitational waves they emit during inspiral. A primary observable is the chirp mass. A thorough analysis of the dependence of the observed chirp mass distribution in an Euclidean space has already been published (Bulik & Belczyński, 2003). A detailed

analysis (Bulik, Belczyński, Rudak 2003) shows that the cosmological effects and various star formation rate histories do not play a significant role in determining the shape of the distribution of observed chirp masses.

We also investigate the possibility of detecting compact object binaries through gravitational lensing. The masses of individual objects in a binary lens depend on the unknown position of the lens between the observer and the lensed star. However, the mass ratio can be estimated quite precisely. We use an analytical model of the Galactic potential to analyze the distribution of observed mass ratios in the microlensing events on stars in the galactic bulge (Bulik, Moderski, Belczyński 2003).

We acknowledge the KBN grants 5P03D01120 (TB), 5P03D00221 (RM), and the Lindheimer Fellowship (KB).

REFERENCES

- Belczyński, K., Kalogera, V., & Bulik, T. 2002, *ApJ*, 572, 407
- Bulik, T. & Belczyński, K. 2003, *ApJ*, 589, L37
- Bulik, T., Belczyński, K. & Rudak, B. 2003, *A&A*, in press
- Bulik, T., R. Moderski & Belczyński, K. 2003, in preparation

¹Nicolaus Copernicus Astronomical Center, Warsaw, Poland.

²Northwestern University, Evanston, IL, USA.