



Autopsy and Case Reports

E-ISSN: 2236-1960

autopsy.hu@gmail.com

Hospital Universitário da Universidade de
São Paulo
Brasil

Martins de Oliveira Junior, Edson; Ferreira Alves, Carlos Augusto; Gonzaga Santos-Queiroz, Adalmir; Melhem Elias, Fernando; de Campos, Antônio Carlos

Bilateral lingual lipoma: a case report

Autopsy and Case Reports, vol. 2, núm. 2, abril-junio, 2012, pp. 21-23

Hospital Universitário da Universidade de São Paulo

São Paulo, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=576060814006>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Bilateral lingual lipoma: a case report

Edson Martins de Oliveira Junior^a, Carlos Augusto Ferreira Alves^a,
Adalmir Gonzaga Santos-Queiroz^a, Fernando Melhem Elias^{a,b}, Antônio Carlos de Campos^{a,b}

Oliveira Junior EM, Alves CAF, Santos-Queiroz AG, Elias FM, Campos AC. Bilateral lingual lipoma: a case report. Autopsy Case Rep [Internet]. 2012;2(2):21-23. <http://dx.doi.org/10.4322/acr.2012.012>

ABSTRACT

Lipoma of the tongue is an uncommon benign tumor, being even more rare when there are multiple lesions. Lipoma accounts for 1-5% of all oral neoplasms. Lipoma occurs predominantly in males over the age of forty, although, in rare cases, children can be affected. Here, we report an unusual case of a female patient with bilateral lingual lipomas that were treated through surgical excision.

Keywords: Lipoma; Tongue; Surgical procedures, operative.

INTRODUCTION

A lipoma is a benign, slow-growing, asymptomatic tumor of mesenchymal origin; lipomas consist of adipose tissue and can be sessile or pedunculated, as well as single or lobulated, typically surrounded by a fibrous capsule.^{1,2} Although the etiology of lipoma remains unclear, causal factors include endocrine changes, trauma, and genetic makeup.³

Although lipoma of the oral cavity is uncommon, lipomas of the head and neck account for 15-20% of all head and neck neoplasms.⁴

Lipomas and 1-5% of all neoplasms of the oral cavity.²

The diagnosis of lipoma is predominantly based on clinical findings, histopathological findings being conclusive in all cases. Imaging tests can be useful, especially when lipomas are located in the deeper fascia of the neck or face. Histologically, lipomas consist of well-differentiated adipose tissue surrounded by a connective tissue capsule.⁵ Here, we report a case of bilateral lipoma of the lateral borders of the tongue treated with surgical excision.

CASE REPORT

An 85-year-old female patient sought medical attention complaining of a "lump" in her tongue. The patient reported that the lump had been present for approximately 5 years. Oral examination revealed

painless, endophytic nodules in the right and left lateral borders of the tongue. The lesion on the right was 1.0 cm in diameter, compared with 1.8 cm for the lesion on the left, in their longest axis. Both were

^a Department of Dentistry - Hospital Universitário - Universidade de São Paulo, São Paulo/SP - Brazil.

^b Department of Surgery, Prosthesis and Maxillofacial Trauma - Faculdade de Odontologia - Universidade de São Paulo, São Paulo/SP - Brazil.

soft, yellowish lesions with well-defined borders and no ulcerations (Figure 1).

A presumptive diagnosis of lipoma was made, and the patient underwent surgical excision under local anesthesia. Macroscopic examination revealed yellowish nodules with a lobular surface (Figure 2). Microscopic examination revealed the presence of mature adipose tissue interspersed with bands of dense connective tissue, confirming the diagnosis of lipoma.

In the case reported here, the histological features of the lipomas (i.e., predominance of simple lipomas), their clinical features (asymptomatic, encapsulated, round, yellowish/mucosa-like in color, smooth, and without ulceration), their anatomical location (the dorsum of the tongue), and their presentation (bilateral) were consistent with those reported in the literature, although the age and gender of the patient (an 85-year-old female) were not. At this writing, 2 years after the surgical excision of the lipomas, there were no signs of recurrence.

DISCUSSION

Lipomas rarely affect the oral cavity, accounting for only approximately 5% of all intraoral neoplasms. The etiology of lipoma has yet to be definitively established but includes endocrine changes, hereditary factors, local trauma, and infection.⁶ Histologically, lipomas are indistinguishable from normal adipose tissue.⁵




Figure 1 – Initial clinical appearance of the lesions, which were located in the lateral border of the tongue.



Figure 2 – Surgical excision of the lipomas.

Although some studies have reported that males are more often affected,^{2,7} others have reported that lipomas are more common in females.^{8,9} Individuals in the 30-70 year age bracket are the most affected, lipomas being rare in those under 20 years of age.^{2,10} Regarding the intraoral commitment, the buccal mucosa, the vestibular fornix, the floor of the mouth, the tongue, the inner surface of the lips, the alveolar ridge, the palate, and the salivary glands, are the most commonly affected sites.¹⁰

Lipomas can be double (bilateral) or, in rare cases, multiple. Multiple lipomas are generally related to lipomatosis, neurofibromatosis type 1 (formerly known as von Recklinghausen's disease), Gardner syndrome, and adiposis dolorosa (also known as Dercum's disease).

The differential diagnosis of lipoma includes epidermoid cysts, lymphoepithelial cysts, ranulas, pleomorphic adenomas, and mucoepidermoid carcinomas.¹¹ In suspected cases of deep-seated lipoma, computed tomography can assist in ruling out vascular lesions and in identifying important anatomical structures that are adjacent to the lesions.¹²

The histopathological features of lipomas are typical: they consist of well-differentiated adipose tissue surrounded by a connective tissue capsule. Although morphologically indistinguishable from normal fat cells, tumor cells are metabolically more active. The fibrous capsule may sometimes be absent or ruptured.¹³

Conservative treatment of lipomas involves complete surgical excision by excisional biopsy, especially in cases of infiltrative lipoma.¹⁴ Recurrence is rare when the lesion is completely excised.^{2,3,6,10}

As evidenced by the case reported here, lipoma of the oral cavity can be treated successfully through conservative treatment.

REFERENCES

1. Weiss SW, Goldblum JR. Benign lipomatous tumours. In: Weiss SW, Goldblum JR, editors. *Enzinger and Weiss's soft tissue tumours*. 4th ed. St. Louis: Mosby; 2001. p. 571-639.
2. Fregnani ER, Pires FR, Falzoni R, Lopes MA, Vargas PA. Lipomas of the oral cavity: clinical findings, histological

- classification and proliferative activity of 46 cases. *Int J Oral Maxillofac Surg.* 2003;32(1):49-53. PMID:12653233. <http://dx.doi.org/10.1054/ijom.2002.0317>
3. Tommasi AF. Diagnóstico em patologia bucal. São Paulo: Artes Médicas; 1982. Português.
 4. Gnepp DR. Diagnostic surgical pathology of the head and neck. Philadelphia: WB Saunders; 2001.
 5. Ribeiro Neto N, Marques JAF, Santos MAM, Parra GR, Mota GCC, Barreto AP. Limpoma de Tamanho Incomum em Lábio Inferior. *Rev Cir Traumatol Buco-Maxilo-Fac.* 2010;10(4):9-12. Português.
 6. Marzola C. Fundamentos de cirurgia buco-maxilo-facial. Bauru: Independente; 2005. Português.
 7. Furlong MA, Fanburg-Smith JC, Childers EL. Lipoma of the oral and maxillofacial region: site and subclassification of 125 cases. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2004;98(4):441-50. PMID:15472660. <http://dx.doi.org/10.1016/j.tripleo.2004.02.071>
 8. Neville BW, Damm DD, Allen CM, Bouquot. Patologia oral & maxilofacial. 2a. ed. Rio de Janeiro: Guanabara Koogan; 2008. Português.
 9. Assis GM, Silvam SRP, Moraes PH, Amaral JIQ, Germano AR. Lipoma facial removido por acesso intrabucal: caso clínico. *Rev Bras Cir Traumatol Buco-Maxilo-Fac.* 2010;10(2):89-93.
 10. Prado R, Ribeiro DPB, Fontoura RA, Sampaio RKPL, Moreira LC. A case of sublingual lipoma. *Rev Bras Odontol.* 1998;55(4):226-8.
 11. Hattori H. Atypical lipomatous tumor of the lip with pleomorphic lipoma-like myxoid area, clinically simulating mucocele. *J Oral Pathol Med.* 2002;31(9):561-4. PMID:12269996. <http://dx.doi.org/10.1034/j.1600-0714.2002.00151.x>
 12. Zhong LP, Zhao SF, Chen GF, Ping FY. Ultrasonographic appearance of lipoma in the oral and maxillofacial region. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2004;98(6):738-40. PMID:15583549. <http://dx.doi.org/10.1016/j.tripleo.2004.04.022>
 13. Epivatianos A, Markopoulos AK, Papanayotou P. Benign tumors of adipose tissue of the oral cavity: a clinicopathologic study of 13 cases. *J Oral Maxillofac Surg.* 2000;58(10):1113-7. PMID:11021705. <http://dx.doi.org/10.1053/joms.2000.9568>
 14. Salvatore C, Antonio B, Del Vecchio W, Lanza A, Tartaro G, Giuseppe C. Giant infiltrating lipoma of the face: CT and MR imaging findings. *AJNR Am J Neuroradiol.* 2003;24(2):283-6. PMID:12591650.

Conflict of interest: None

Submitted on: 5th January 2012

Accept on: 16th January 2012

Correspondence: Divisão de Odontologia

Av. Prof. Lineu Prestes, 2565 – Cidade Universitária - São Paulo/SP – Brazil

CEP: 05508-900 – Phone: +55 (11) 3091-9290

E-mail: jr.odonto@gmail.com