



Revista Mexicana de Fitopatología

ISSN: 0185-3309

mrlegarreta@prodigy.net.mx

Sociedad Mexicana de Fitopatología, A.C.

México

García Velasco, Rómulo; Ochoa Martínez, Daniel Leobardo; Zavaleta Mejía, Emma; Mora Aguilera, Gustavo; Gaytán Acuña, Elda Araceli
Características Morfológicas de Basidiosporas de *Puccinia horiana* Henn. en *Dendranthema grandiflora* Tzvelev cv. Polaris
Revista Mexicana de Fitopatología, vol. 20, núm. 2, julio-diciembre, 2002, pp. 213-215
Sociedad Mexicana de Fitopatología, A.C.
Texcoco, México

Available in: <http://www.redalyc.org/articulo.oa?id=61220214>

Abstract

Light brown pustules were collected to characterize basidia of *Puccinia horiana*. Transversal cuts of pustules were made, and the pieces were placed in a drop of blue lactophenol on a glass slide. Fungal structures were observed with a light microscope and measured. Basidia were oval, cream color; the mean length was $9.75 \mu\text{m}$ x $7.20 \mu\text{m}$ width, and an acute form near the union to the sterigma.

Keywords

Basidia, teliospores, chrysanthemum

- ▶ How to cite
- ▶ Complete issue
- ▶ More information about this article
- ▶ Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative