



Madera y Bosques

ISSN: 1405-0471

mabosque@inecol.mx

Instituto de Ecología, A.C.

México

Báez-Pérez, Ana Laura; Lindig-Cisneros, Roberto; Villegas, Javier  
Survival and growth of nursery inoculated *Fraxinus uhdei* in acrisol gullies  
Madera y Bosques, vol. 23, núm. 3, 2017, pp. 7-14  
Instituto de Ecología, A.C.  
Xalapa, México

Available in: <http://www.redalyc.org/articulo.oa?id=61753521001>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



# Survival and growth of nursery inoculated *Fraxinus uhdei* in acrisol gullies

## Supervivencia y crecimiento de *Fraxinus uhdei*, inoculado en vivero, en cárcavas de acrisoles

Ana Laura Báez-Pérez<sup>1</sup>, Roberto Lindig-Cisneros<sup>1\*</sup> and Javier Villegas<sup>2</sup>

<sup>1</sup> Universidad Nacional Autónoma de México. Instituto de Investigaciones en Ecosistemas y Sustentabilidad. Laboratorio de Ecología de Restauración. Morelia, Michoacán, México.

<sup>2</sup> Universidad Michoacana de San Nicolás de Hidalgo. Instituto de Investigaciones Químico-Biológicas. Morelia, Michoacán, México.

\* Corresponding author. rlindig@iies.unam.mx.

### ABSTRACT

Presence of degraded soils is increasing worldwide, but in many scenarios the knowledge to undertake restoration has not been acquired in an efficient, economical and socially acceptable manner. There are estimates placing moderate to severe degradation of agricultural land worldwide at 80%. Severely degraded substrates, such as gullies, sometimes lack the necessary microbiota for appropriate plant growth, under these conditions nursery inoculation can improve plant performance in the field. In order to explore the role of dual mycorrhization in *Fraxinus* growing in poor substrates, and also to contribute information on their usefulness in gully restoration, an experiment was conducted in which individuals of this species were inoculated with an endomycorrhizal fungus, and ectomycorrhizal fungus and with both types of mycorrhizal fungi simultaneously, as well as urea fertilization. The results of this experiment show that establishment of *Fraxinus uhdei* is possible in severely degraded sites, it also provides information on the effects on the plant of multiple inoculation that cause some performance variables to improve as a result of the interaction, as was the case for the improved survival with the *P. tinctorius*, *G. intraradices* dual inoculation; whereas growth variables showed improved response to inoculation with only one of the fungi, height to inoculation with *G. intraradices*, and cover, number of leaves and diameter at base to inoculation with *P. tinctorius*. Given the improvements in growth and survival, our results suggest that in severely degraded soils, nursery inoculation of plants adapted to adverse conditions with mycorrhizal fungi should be implemented.

**KEYWORDS:** erosion, mycorrhiza, plant cover, positive interactions, restoration, revegetation.

### RESUMEN

La presencia de suelos degradados sigue incrementándose a nivel mundial y, para muchos casos, no se cuenta con el conocimiento necesario para llevar a cabo su restauración de manera eficiente. Los sustratos severamente degradados, como las cárcavas, en ocasiones no tienen la microbiota necesaria para un crecimiento adecuado de las plantas y, bajo estas condiciones, la inoculación en vivero puede mejorar el desempeño de las plantas en campo. Para explorar el papel de la inoculación dual de *Fraxinus* creciendo en sustratos pobres, así como para aportar información sobre la restauración de cárcavas, se llevó a cabo un experimento en el cual se inocularon individuos de esta especie con un hongo ectomicorrízico, un hongo endomicorrízico y ambos simultáneamente, así como la fertilización con urea. Los resultados del experimento indican que el establecimiento de *Fraxinus uhdei* es posible en sitios severamente degradados y proporcionan información sobre el efecto de la inoculación múltiple en plantas que causa que algunas variables de desempeño mejoren como consecuencia de la interacción, como fue el caso de mayor supervivencia con la inoculación dual con *P. tinctorius* y *G. intraradices*. Mientras que las variables de crecimiento respondieron a la inoculación con uno de los hongos, altura a la inoculación con *G. intraradices*, y la cobertura, número de hojas y diámetro a la base a *P. tinctorius*. Dada la mejoría en supervivencia y crecimiento, estos resultados sugieren que en suelos severamente degradados la inoculación de plantas capaces de crecer en sitios adversos debe de implementarse.

**PALABRAS CLAVE:** erosión, micorrizas, cobertura vegetal, interacciones positivas, restauración, revegetación.

## INTRODUCTION

The presence of degraded soils is increasing worldwide and their restoration must be part of the strategies to meet the challenges brought about by global change (Lal, 2004; Bigas, Gudbrandsson, Montanarella and Arnalds, 2007). However, even though the importance of restoration of degraded sites has been recognized for some time (Bradshaw and Chadwick, 1980), in many scenarios the requisite knowledge to undertake restoration has not yet been acquired in an efficient, economical and socially acceptable manner. Soil erosion is a severe form of physical degradation. There are estimates placing moderate to severe degradation of agricultural land worldwide at 80% (Lal and Stewart, 1995). It is well known that plants control erosion by acting as a barrier to the impact of rain and the force of winds, while favoring infiltration and reducing runoff, thus preventing soil loss, and, in severe cases, gully formation. Once the soil has degraded and gullies have formed, complex measures are required to stop even further degradation, and recovery can take several decades (Mongil-Manso, Navarro-Hevia, Diaz-Gutierrez, Cruz-Alonso and Ramos-Diez, 2016). For example, Morgan (1986) suggests that a multiple-strata revegetation process is required using grasses, legumes, shrubs and trees around gullies to stop their expansion. Nevertheless, this approach does not solve the problem of establishing plant cover on the slopes of the gullies themselves.

Mycorrhizal fungi are of particular importance for the recovery of soil structure and dynamics. These fungi play a key role in the establishment and development of most plants (Colpaert, Tichele, Assche and Laere, 1999; Allen, Swenson, Querejeta, Egerton-Warburton and Treseder, 2003). Among the multiple benefits attributed to mycorrhization, its effect on plant development stands out, since there is widespread proof that mycorrhizas improve plant growth by increasing the absorption surface of the root system (Smith and Read, 1997), selectively absorbing and storing certain nutrients (Colpaert *et al.*, 1999), solubilizing certain chemicals and making some chemical elements that are normally insoluble, in particu-

lar phosphorus, available for the plant (Casarin, Plassard, Hinsinger and Arvieu, 2004; Liu, Loganathan, Hedley and Skinner, 2004). There are plants capable of forming simultaneous symbiotic associations with both arbuscular mycorrhizal fungi and ectomycorrhizal fungi, which may contribute to improved environmental restoration strategies (Gómez-Romero *et al.*, 2012; Gómez-Romero, Villegas, Sáenz-Romero and Lindig-Cisneros, 2013)

Ashes (*Fraxinus* spp.) are able to establish themselves on degraded sites to the extent that at least one species of the genus has turned into an invasive species by its introduction into areas outside its distribution range (Mulla *et al.*, 2014). Nevertheless, some native species can be a good choice for revegetation of gullies within its natural distribution ranges, since the species of this genus are resistant to stress conditions caused by soils that are poor in nutrients and organic matter (Francis, 1990; Stabler, Martin and Stutz, 2001), and this may be the case of *F. uhdei* (Wenz.) Lingelsh. This is a species found in Mexico and Central America that is capable of forming symbiotic associations with both endo and ectomycorrhizas (Ambriz, Báez-Pérez, Sánchez-Yáñez, Moutoglis and Villegas, 2010), and previous experiments have shown that urea has a significant effect on the development of the external mycelium of the *Pisolithus tinctorius* ectomycorrhizal fungus associated with this host plant (Báez-Pérez, Sánchez and Villegas, 2010). For other species of this genus, it has been suggested that success in the establishment in degraded areas may be result of their capability in forming symbiotic associations with mycorrhizal fungi (Stabler *et al.*, 2001).

In a preliminary controlled experiment in large containers (Báez-Pérez *et al.*, 2015), filled with substrate obtained from gullies, suggested the potential of inoculated *F. uhdei* plants for restoration of gullies formed in Acrisols but the field test was lacking. In fact, there are very few articles that address the contribution of the *Fraxinus*-mycorrhizal fungi system to plant performance in substrata, under field conditions, that are unsuitable for normal plant development due to severe soil erosion processes.



## OBJECTIVES

In order to explore the role of dual mycorrhization and urea fertilization on *Fraxinus* plants survival and growth in poor substrates, and also to contribute information on their usefulness in gully restoration, an experiment was conducted in which individuals of this species were inoculated with an endomycorrhizal fungus, and ectomycorrhizal fungus and with both types of mycorrhizal fungi simultaneously, and urea fertilization.

## MATERIALS AND METHODS

Seeds of *Fraxinus uhdei* were collected in the locality of Huertitas, Ejido de Atécuaro, Morelia, Michoacán (19° 33' 05" and 19° 37' 08" N and 101° 05' 07" W; altitude 2275 m a.s.l.), where the experimental site is located. The area is mostly devoid of vegetation as a result of deforestation and agricultural practices unsuited for the fragile acrisol soils, which have led to the formation of deep gully systems. The area has a subhumid climate with rains in summer (July to October), with an average annual temperature of 14.8 °C and annual mean rainfall of 1000 mm (Duvert *et al.*, 2010).

Plants of *F. uhdei* were propagated in a shade house (20%) in March 2012. The seeds were scarified manually and disinfected with hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) at 10% (v:v) for 15 min. They were then planted in rigid plastic containers (375 cm<sup>3</sup>), using a perlite-peat moss substrate (2:1 v/v) that was autoclaved at 121 °C and 15 PSI during 30 min.

After 9 weeks, the plants were separated into four groups. Plants in the first group were not inoculated, and those in the other three groups were inoculated as follows: a) one group with the mycorrhizal fungus *P. tinctorius* (1 x 10<sup>6</sup> spores/plant) using micronized peat as a vehicle; b) a second one with the arbuscular mycorrhizal fungus *G. intraradices* (50 spores/plant) applied as a suspension in distilled water; and c) the third one with a mixture of both fungal species (dual inoculation), each species applied as mentioned above. The *P. tinctorius* inoculum was acquired from Biosyneterra Solutions Inc. (L'Assomption, Québec, Canada), and *G. intraradices* was obtained from isolated cultures from the Laboratorio de Interacción

Suelo-Planta-Microorganismo of the Instituto de Investigaciones Químico Biológicas de la Universidad Michoacana de San Nicolás de Hidalgo.

After 17 weeks had passed from inoculation, and before being taken to the field, 10 individual ash plants from each inoculation treatment were randomly selected to verify mycorrhization as follows. The arbuscular mycorrhizal fungus was identified using the ink and vinegar staining technique described by Vierhelig, Coughlan, Wyss and Piche (1998) by clarifying the roots with 10% KOH (w/v) for 30 min, to stain afterwards with Shaeffer black ink dissolved to 10% with 25% acetic acid, rinsing excess staining agent with running water. Once the roots had been stained, the characteristic structures of the species (arbuscules, vesicles and/or not-septated mycelium) were identified by taking photographs with a Leica DFC 295 (version 7.0.1.0) camera, using specialized software (Leica Application Suite Version 3.4.1). *P. tinctorius* was detected using a technique developed by Thompson, Grove, Malajczuk y Hardy (1993) modified by Ambriz *et al.* (2010), by staining the mantle and external mycelium of the fungus associated to the fine root segments of the ash with trypan blue at 0.05% for 24 h and then rinsing the dye with distilled water before observation through a microscope to identify the characteristic septated mycelium of this species.

The experimental plot had very little organic matter (0.78%), very low total available nitrogen (19.60 kg/ha) and phosphorus (0.347 mg/kg). The ash plants were distributed randomly, 14 replicas per treatment, on a westward-facing slope of a gully 6 m deep. Once planted, half of the plants of each inoculation treatment were fertilized with urea (NH<sub>2</sub>CONH<sub>2</sub>) dissolved in water at a concentration of 200 mM, each plant receiving 0.384 g of urea dissolved in 250 ml of water, once every 15 days during the rainy season (July to November). No watering took place during the dry season.

Due to the combination of inoculation and urea fertilization factors, the experiment consisted in an orthogonal design with eight treatments: 1) *F. uhdei* plants, 2) *F. uhdei* plants + urea, 3) *F. uhdei* plants + *G. intraradices*,

4) *F. uhdei* plants + *G. intraradices* + urea, 5) *F. uhdei* plants + *P. tinctorius*, 6) *F. uhdei* plants + *P. tinctorius* + urea, 7) *F. uhdei* plants + dual inoculation and 8) *F. uhdei* plants + dual inoculation + urea. The experiment was evaluated during 23 months, registering plant survival and measuring height on a monthly basis, diameter at base (DAB 1 cm from soil surface), cover (measured by the ellipse formula  $r_1 r_2 \pi$  with two radiuses for the crown) and number of leaves every six months.

Statistical analyses were carried out using generalized linear models followed by hypothesis testing with error II. A normal distribution was used for the analysis of all of the variables except the number of leaves, for which a Poisson-type distribution was used. Where one of the factors was not significant for a given variable, the factor was removed from the analysis, which allowed using the full data set in those cases (Crawley, 2007). All of the analyses were carried out using R (R Development Core Team, 2015).

## RESULTS

Root tissue preparations of the plants for the three inoculation treatments showed that all plants formed mycorrhiz-

as with *Glomus intraradices*, *Pisolithus tinctorius* or with both fungus species simultaneously (Fig. 1). There were differences in survival due to the inoculation treatments. Greatest survival was observed in plants with dual inoculation (64%), followed by those inoculated with *P. tinctorius* (57%) and finally by those that were not inoculated (46%), regardless of fertilization. The significant differences among treatments ( $\chi^2 = 11.1$ ; d.f. = 3;  $p = 0.01$ ) were the result of the difference between the plants that received dual inoculation and control plants.

Plant growth started to show differences among treatments at the end of the first growing season (during the fourth and fifth months of evaluation) when mean values became noticeable different for some of the treatments, as reflected in changes in height (Fig. 2). However, differences started to be significant after 12 months (during the second growing season), in particular between the plants inoculated with *G. intraradices*, regardless of fertilization and non-inoculated control plants (Table 1). The data on the last evaluation showed that plants inoculated with *G. intraradices* were the tallest ( $26.6 \text{ cm} \pm 4.6 \text{ cm}$ ), followed by the plants inoculated with




FIGURE 1. In (a) roots of *Fraxinus uhdei* from control plants with no evidence of mycorrhiza, in (b) roots with *Glomus intraradices* vesicles; in (c) roots with *Pisolithus tinctorius* external mycelium (me); in (d) spores of *Glomus intraradices* (e) and external mycelium of *Pisolithus tinctorius* (indicated by the arrows), in (e) septated hyphae of *P. tinctorius*.



*P. tinctorius* ( $26.0 \pm 4.9$ ), then the plants with dual inoculation ( $24.0 \text{ cm} \pm 4.6 \text{ cm}$ ) and finally, non-inoculated plants ( $22.3 \text{ cm} \pm 4.1 \text{ cm}$ ). Differences attributable to this factor are significant ( $F_{(3,98)} = 4.6$ ,  $p = 0.005$ ), since plants inoculated with any of the mycorrhizogenic fungi were taller than those in the control group. The effect of fertilization was marginally significant ( $F_{(1,98)} = 4.6$ ,  $p = 0.03$ ), since fertilized plants were only slightly taller ( $25.9 \text{ cm} \pm 5.0 \text{ cm}$ ) than non-inoculated plants ( $23.8 \text{ cm} \pm 4.5 \text{ cm}$ ), and there being no interaction between fertilization and inoculation.

Cover also responded to inoculation treatments and followed a pattern similar to that for height, but the values for the plants inoculated with only one of the fungal species were inverted, since this variable was greater for plants inoculated with *P. tinctorius* ( $101.3 \text{ cm} \pm 69.8 \text{ cm}$ ), followed by those inoculated with *G. intraradices* ( $69.6 \text{ cm} \pm 45.8 \text{ cm}$ ), then plants with dual inoculation ( $68.6 \text{ cm} \pm 55.1 \text{ cm}$ ) and finally non-inoculated plants ( $46.4 \text{ cm} \pm$

$37.1 \text{ cm}$ ). The difference among inoculation treatments was significant ( $F_{(3,98)} = 4.3$ ,  $p = 0.007$ ), and the effect of fertilization was not ( $F_{(1,98)} = 3.6$ ,  $p = 0.06$ ). In regard to number of leaves, the highest values were observed for plants inoculated with *P. tinctorius* ( $13 \pm 6$ ), followed by plants with dual inoculation ( $10 \pm 5$ ), then those inoculated with *G. intraradices* ( $10 \pm 5$ ), and finally non-inoculated plants ( $9 \pm 4$ ). Differences among treatments were statistically significant ( $\chi^2 = 17.1$ ; d.f. = 3;  $p = 0.0006$ ), as a result of the difference between plants inoculated with *G. intraradices* and control plants. The effect of fertilization on the number of leaves was marginal ( $\chi^2 = 6.2$ ; d.f. = 1;  $p = 0.01$ ). Plants fertilized with urea had more leaves ( $11 \pm 6$ ) than those that were not. Lastly, diameter at base was larger for the plants inoculated with *P. tinctorius* ( $0.94 \pm 0.20$ ), followed by those inoculated with *G. intraradices* ( $0.83 \text{ cm} \pm 0.18 \text{ cm}$ ), then those with dual inoculation ( $0.81 \text{ cm} \pm 0.20 \text{ cm}$ ), and finally non-inoculated plants ( $0.74 \text{ cm} \pm 0.17 \text{ cm}$ ). The difference between inocu-




FIGURE 2. Growth, as height, of *Fraxinus uhdei* during the lapse of the experiment. Growth occurred during the rainy seasons and the differences become evident after one year of planting.

TABLE 1. Performance parameters of *Fraxinus uhdei* plants at the end of the experiment in response to inoculation.

Treatments	All plants	Westward-facing slope plants		
	Survival (%)	Height (cm)	Cover (cm <sup>2</sup> )	Leaves
<i>Fraxinus uhdei</i>	46	22.3 ± 4.1	46.4 ± 37.1	9 ± 4
<i>F. + G. intraradices</i>	62	26.6 ± 4.6	69.6 ± 45.8	10 ± 5
<i>F. uhdei + P. tinctorius</i>	57	26.0 ± 4.9	101.3 ± 69.8	13 ± 6
<i>F. uhdei + G.intraradices + P. tinctorius</i>	64	24.0 ± 4.6	68.6 ± 55.1	10 ± 5

Survival is for all plant in both slopes (see text) and growth parameters are for the plants in the westward-facing slope where enough plants survived allowing statistical analyses. Data are means and standard deviations.

lation treatments was significant ( $F_{(3,98)} = 4.9$ ,  $p = 0.003$ ), and the effect of fertilization was not ( $F_{(1,98)} = 21$ ;  $p = 0.15$ ), and no interaction was detected between both experimental factors.

## DISCUSSION

The results of this experiment show that establishment of *Fraxinus uhdei* is possible in severely degraded sites where the substrate is no longer suitable for natural plant development. Although it has been documented that some of the species of this genus are tolerant to stress in sites that are unsuitable for other tree species (Francis, 1990; Stabler *et al.*, 2001), conditions present at this study site were even more severe than those mentioned in such studies. In addition, this study provides information on the effects on the plant of multiple mycorrhizal interactions in scenarios of this type, in which some performance variables improve as a result of the interaction, as was the case for improved survival with the *P. tinctorius*, *G. intraradices* dual inoculation; whereas growth variables showed improved response to inoculation with only one of the fungi, height to inoculation with *G. intraradices*, and cover, number of leaves and diameter at base to inoculation with *P. tinctorius*.

The *F. uhdei*-*G. intraradices*-*P. tinctorius* interaction has not been studied in depth, and has only been addressed under controlled conditions in growth chambers (Ambriz *et al.*, 2010) or in mesocosms (Báez-Pérez *et al.*, 2015) but never in the field, as in this study. Nevertheless,

our results coincide with those found by Ambriz *et al.* (2010), who first reported the symbiotic association with both species of mycorrhizal fungi and observed the synergic effect of dual inoculation on ash development when compared to single inoculation.

It is interesting to note that nitrogen fertilization had a marginal effect on survival and height of *F. uhdei* only, and had no effect on the remaining variables analyzed. As regards to growth, they were consistent with the results found by Canellas, Finat, Bachiller and Montero (1999); however, with respect to survival they were not, and the difference could be attributed to the conditions to which the plants were exposed in our experiment, given the extremely low level of nutrients in the substrate.

## CONCLUSIONS

The experiment showed that the highest values obtained for the growth variables analyzed were those for inoculation with *P. tinctorius*, but not different from dual inoculation and statistically different when compared to inoculation with *G. intraradices* or the control (no inoculation), and this differs somewhat from the results of other works that show a synergy for growth variables in plants with dual inoculation (Founoune *et al.*, 2001; Misbahuzzaman and Newton, 2006; Ambriz *et al.*, 2010; Báez-Pérez *et al.*, 2015). It is possible for this difference to be the result of drought conditions during the dry season, because it has been shown that ectomycorrhizal fungi increase the ability of plants to resist drought stress through morphophysio-



gical and biochemical mechanisms (Parke, Linerman and Black, 1983; Alvarez *et al.*, 2009), although a more careful study in this regard is required. Finally, and given the improvements in survival of *F. uhdei*, in response to dual inoculation, our results suggest that in severely degraded soils, inoculation of this type may constitute an efficient strategy for environmental restoration.

## ACKNOWLEDGEMENTS

This research was partially funded by a grant by DGAPA-UNAM (IN-203316). A.L.B.P. wishes to thank CONACYT for a postdoctoral grant.

## REFERENCES

- Allen, M. F., Swenson, W., Querejeta, J. I., Egerton-Warburton, L. M. and Treseder, K. K. (2003). Ecology mycorrhizae: A conceptual framework for complex interactions among plants and fungi. *Annual Review in Phytopathology*, 41, 271-303. <https://doi.org/10.1146/annurev.phyto.41.0520.02.095518>
- Alvarez, M., Huygens, D., Fernandez, C., Gacitúa, Y., Olivares, E., Saavedra, I., Alberdi, M. and Valenzuela, E. (2009). Effect of ectomycorrhizal colonization and drought on reactive oxygen species metabolism of *Nothofagus dombeii* roots. *Tree Physiology*, 29(8), 1047-1057. doi: 10.1093/treephys/tpp038
- Ambriz, E., Báez-Pérez, A., Sánchez-Yáñez, J. M., Moutoglou, P. and Villegas, J. (2010). *Fraxinus-Glomus-Pisolithus* symbiosis: Plant growth and soil aggregation effects. *Pedobiologia*, 53, 369-373. doi: 10.1016/j.pedobi.2010.07.001
- Báez-Pérez, A. L., Sánchez, N. J. and Villegas, M. J. (2010). Efecto de la simbiosis *Pisolithus tinctorius-Fraxinus uhdei*, bajo tres fuentes de N (NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup> y urea). *Ciencia Nicolaita*, No. Especial 2010, 1-9.
- Báez-Pérez, A. L., Gómez-Romero, M., Villegas, J., de la Barrera, E., Carreto Montoya, L. and Lindig-Cisneros, R. (2015). Inoculación con hongos micorrízicos y fertilización con urea de plantas de *Fraxinus uhdei* en acrisoles provenientes de sitios degradados. *Botanical Sciences*, 93, 501-508.
- Bigas, H., Gudbrandsson, G. I., Montanarella, L. and Arnalds, A. (Eds.) (2007). Soils, society & global change. Proceedings of the international forum celebrating the centenary of conservation and restoration of soil and vegetation in Iceland. Retrieved from [http://inweh.unu.edu/wp-content/uploads/2013/05/SoilsSocietyGlobalChange\\_final\\_low-res1.pdf](http://inweh.unu.edu/wp-content/uploads/2013/05/SoilsSocietyGlobalChange_final_low-res1.pdf)
- Bradshaw, A. D. and Chadwick M. J. (1980). *The restoration of land: the ecology and reclamation of derelict and degraded land*. Los Angeles, United States: University of California Press.
- Canellas, I., Finat, L., Bachiller, A. and Montero, G. (1999). Comportamiento de planta de *Pinus pinea* en vivero y campo: ensayos de técnicas de cultivo de planta, fertilización y aplicación de herbicidas. *Investigaciones Agrícolas: Producción y Protección Vegetal*, 8, 335-359.
- Casarin, V., Plassard, C., Hinsinger, P. and Arvieu, J. (2004). Quantification of ectomycorrhizal fungal effects on the bioavailability and mobilization of soil P in the rhizosphere of *Pinus pinaster*. *New Phytologist*, 163, 177-185. doi: 10.1111/j.1469-8137.2004.01093.x
- Colpaert, J. V., Tichelen, K. K., Assche, J. A. and Laere, A. (1999). Short-term phosphorus uptake rates in mycorrhizal and non-mycorrhizal roots of intact *Pinus sylvestris* seedlings. *New Phytologist*, 143, 589-597. DOI: 10.1046/j.1469-8137.1999.00471.x
- Crawley, R. (2007). *The R Book*. John Wiley and Sons, Ltd.
- Duvert, C., Gratiot, N., Evrard, O., Navratil, O., Némery, J., Prat, C. and Esteves, M. (2010). Drivers of erosion and suspended sediment transport in three headwater catchments of the Mexican Central Highlands. *Geomorphology*, 123, 243-256. doi: 10.1016/j.geomorph.2010.07.016
- Founoune, H., Duponnois, R., Ba, M. A., Sall, S., Branget, I., Lorquin, J., Neyra, M. and Chotte, L. J. (2001). Mycorrhiza helper bacteria stimulate ectomycorrhizal symbiosis of *Acacia holosericea* with *Pisolithus alba*. *New Phytologist*, 153, 81-89. doi: 10.1046/j.0028-646X.2001.00284.x
- Francis, J. K. (1990). *Fraxinus uhdei* (Wenzig) Lingelsh. Fresno, tropical ash. SO-ITF-SM-28. New Orleans, United States: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station.



- Gómez-Romero, M., Soto-Correa, J. C., Blanco-García, J. A., Sáenz Romero, C., Villegas, J. and Lindig-Cisneros, R. (2012). Estudio de especies de pino para restauración de sitios degradados. *Agrociencia*, 46, 795-807.
- Gómez-Romero, M., Villegas, J., Sáenz-Romero, C. and Lindig-Cisneros, R. (2013). Efecto de la micorrización en el establecimiento de *Pinus pseudostrobus* en cárcavas. *Madera y Bosques*, 19(3), 51-63. doi: 10.21829/myb.2013.193327
- Lal, R. (2004). Soil carbon sequestration to mitigate climate change. *Geoderma*, 123, 1-22.
- Lal, R. and Stewart, B. A. (1995). *Soil management, experimental basis for sustainability and environmental quality*. Inc, Boca Raton, Florida: CRC Press.
- Liu, Q., Loganathan, P., Hedley, M. J. and Skinner, M. F. (2004). The mobilization and fate of soil and rock phosphate in the rhizosphere of ectomycorrhizal *Pinus radiata* seedlings in an Allophanic soil. *Plant and Soil*, 264, 219-229.
- Misbahuzzaman, K. and Newton, A. (2006). Effect of dual arbuscular ectomycorrhizal inoculation on mycorrhiza formation and growth in *E. camaldulensis* Dehnh. seedlings under different nutrient regimes. *International Journal of Agriculture and Biology*, 8(6), 848-854.
- Mongil-Manso, J., Navarro-Hevia, J., Diaz-Gutierrez, V., Cruz-Alonso, V. and Ramos-Diez, I. (2016). Badlands forest restoration in Central Spain after 50 years under a Mediterranean-continental climate. *Ecological Engineering*, 97, 313-326. doi: 10.1016/j.ecoleng.2016.10.020
- Morgan, R. P. C. (1986). *Soil erosion and conservation*. Essex, Inglaterra: Longman, Harlow.
- Parke, J. L., Linerman, R. G. and Black, C. H. (1983) The role of ectomycorrhizas in drought tolerance of douglas fir seedlings. *New Phytologist*, 95, 83-95. doi: 10.1111/j.1469-8137.1983.tb03471.x
- R Development Core Team. (2015). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <http://www.R-project.org>.
- Smith, S. F. and Read, D. J. (1997). *Mycorrhizal symbiosis* (2nd ed.). London, England: Academic Press London.
- Stabler, L. B., Martin, C. A. and Stutz, J. C. (2001). Effect of urban expansion on arbuscular mycorrhizal fungal mediation of landscape tree growth. *Journal of Arboriculture*, 27, 193-202.
- Thomson, B. D., Grove, T. S., Malajczuk, N. y Hardy, G. E. St J. (1993). The effectiveness of ectomycorrhizal fungi in increasing the growth of *Eucalyptus globulus* Labill. in relation to root colonization and hyphal development in soil. *New Phytologist*, 126, 517-524.
- Vierhelig, H., Cougilan, A. P., Wyss, U. and Piche, Y. (1998). Ink and vinegar, a simple staining technique for arbuscular mycorrhizal fungi. *Applied Environmental Microbiology*, 64, 5004-5007.

Received: 6 October 2016.

Accepted: 19 June 2017.

This paper must be cited as:

Báez-Pérez, A. L., Lindig-Cisneros, R. and Villegas, J. (2017). Survival and growth of nursery inoculated *Fraxinus uhdei* in acrisol gullies. *Madera y Bosques*, 23(3), 7-14. doi: 10.21829/myb.2017.2331418