



Revista Mexicana de Ingeniería Química

ISSN: 1665-2738

amidiq@xanum.uam.mx

Universidad Autónoma Metropolitana

Unidad Iztapalapa

México

Cardenas-Sandoval, B.A.; Bravo-Luna, L.; Bermúdez-Torres, K.; Trejo-Espino, J.L.;  
Zamilpa, A.; Trejo-Tapia, G.

ENHANCEMENT OF PHENYLETHANOID GLYCOSIDES BIOSYNTHESIS IN *Castilleja*  
*tenuiflora* Benth. SHOOT CULTURES WITH CELL WALL OLIGOSACCHARIDES FROM  
*Fusarium oxysporum* f. sp. *lycopersici* RACE 3

Revista Mexicana de Ingeniería Química, vol. 14, núm. 3, 2015, pp. 631-639

Universidad Autónoma Metropolitana Unidad Iztapalapa

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=62043088006>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in [redalyc.org](http://www.redalyc.org)

[redalyc.org](http://www.redalyc.org)

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



**ENHANCEMENT OF PHENYLETHANOID GLYCOSIDES BIOSYNTHESIS IN  
*Castilleja tenuiflora* Benth. SHOOT CULTURES WITH CELL WALL  
OLIGOSACCHARIDES FROM *Fusarium oxysporum* f. sp. *lycopersici* RACE 3  
ESTIMULACIÓN DE LA BIOSÍNTESIS DE LOS FENILETANOIDES  
GLICOSILADOS EN CULTIVOS DE BROTES DE *Castilleja tenuiflora* Benth. CON  
OLIGOSACÁRIDOS DE PARED CELULAR DE *Fusarium oxysporum* f. sp. *lycopersici*  
RAZA 3**

B.A. Cardenas-Sandoval<sup>1</sup>, L. Bravo-Luna<sup>1</sup>, K. Bermúdez-Torres<sup>1</sup>, J.L. Trejo-Espino<sup>1</sup>,  
A. Zamilpa<sup>2</sup>, G. Trejo-Tapia<sup>1\*</sup>

<sup>1</sup>Centro de Desarrollo de Productos Bióticos, Instituto Politécnico Nacional, P. O. Box 24, 62730, Yautepec, Morelos, México.

<sup>2</sup>Centro de Investigación Biomédica del Sur, Instituto Mexicano del Seguro Social, Argentina No. 1, Col. Centro, Xochitepec, Morelos, 62790, México.

Received April 8, 2015; Accepted July 14, 2015

**Abstract**

Phenylethanoid glycosides (PhGs) are promising natural products for the treatment of chronic diseases because of their wide range of biological activities. Biotic stress as pathogen attack (fungi) may stimulate the synthesis of PhGs through activation of phenylalanine ammonia-lyase (PAL). *Castilleja tenuiflora* Benth. (Orobanchaceae) *in vitro* cultures are alternative sources of PhGs, however in that conditions, cultures have a diminished synthesis of these compounds. To increase the yields of PhGs, the identification of factors affecting their biosynthesis is required. Here, we show that elicitation with cell wall oligosaccharides (CWOs) from *Fusarium oxysporum* f. sp. *lycopersici* race 3 (Hyphomycetes) stimulates biosynthesis of PhGs by increasing the activity of PAL. Upon elicitation with CWOs (13 µg/mL) the production of PhGs was enhanced by 5-fold compared with untreated control. The maximum PAL activity in shoots cultured under CWOs elicitation were also increased. Elicitation did not affect the shoot growth (length and biomass) but induced chlorosis, and delayed root formation of *C. tenuiflora* shoots. Our results demonstrate that elicitation with CWOs increases PhGs biosynthesis in *C. tenuiflora* shoot culture.

**Keywords:** PAL, verbascoside, isoverbascoside, *Castilleja tenuiflora*, *Fusarium oxysporum* f. sp. *lycopersici* race 3.

**Resumen**

Los feniletanoides glicosilados (FEGs) son sustancias naturales que pueden utilizarse para el tratamiento de enfermedades crónicas debido a su amplia gama de actividades biológicas. Un estrés biótico como el ataque de patógenos (hongos) a plantas, puede estimular la síntesis de FEGs a través de la activación de la enzima fenilalanina amoníaco-lias (PAL). Los brotes de *Castilleja tenuiflora* Benth. (Orobanchaceae) cultivados *in vitro* producen FEGs, sin embargo, en esas condiciones se presenta una disminución en la síntesis de estos compuestos. Para aumentar los rendimientos de los FEGs se requiere la identificación de los factores que afectan a su biosíntesis. En este trabajo se muestra que la elicitación con oligosacáridos de pared celular (OPCs) de *Fusarium oxysporum* f. sp. *lycopersici* raza 3 (Hyphomycetes) estimula la biosíntesis de FEGs aumentando la actividad de la enzima PAL. La elicitación de los brotes de *C. tenuiflora* con OPCs (13 µg/mL), aumento la producción de FEGs 5 veces en comparación con el control y, provocó un aumento en la actividad máxima de la enzima PAL. Por otro lado, la elicitación no afectó el crecimiento de los brotes (longitud y biomasa), pero si indujo clorosis y retardo la formación de raíces. Nuestros resultados demuestran que la elicitación con OPCs incrementa la biosíntesis de FEGs en cultivos de brotes de *C. tenuiflora*.

**Palabras clave:** PAL, verbascósido, *Castilleja tenuiflora*, elicitación fungica, *Fusarium oxysporum* f. sp. *lycopersici* raza 3.

\* Corresponding author. E-mail: gttapia@ipn.mx  
Tel. +52-735-39-420-20

## 1 Introduction

*Castilleja tenuiflora* Benth. is a member of the Orobanchaceae family, a perennial medicinal herb with proven health benefits including anti-inflammatory (Sánchez *et al.*, 2013), antioxidant (López-Laredo *et al.*, 2012), anti-ulcerogenic (Sánchez *et al.*, 2013) and cytotoxic activities (Moreno-Escobar *et al.*, 2011). The main chemical compounds found in this species are: iridoids glycosides such as aucubin, the phenylethanoid glycosides (PhGs) verbascoside and isoverbascoside (Gómez-Aguirre *et al.*, 2012) and flavonoids such as quercetin-type glycosides (López-Laredo *et al.*, 2012) and apigenin (Gómez, 2011). PhGs are water-soluble natural occurring compounds, absorbable by human intestinal cells and with a wide spectrum of biological activities including anti-fungal (Funari *et al.*, 2012), anti-inflammatory (Gyurkovska *et al.*, 2011), antiradical (López-Laredo *et al.*, 2012), cytotoxic (Pettit *et al.*, 1990), and gastroprotective (Singh *et al.*, 2010) among others.

*Castilleja tenuiflora* in vitro cultures exhibit pharmacological activities comparable to that of wild plants and are attractive sources of PhGs (Sánchez *et al.*, 2013) and iridoids (Martínez-Bonfil *et al.*, 2011) constituting a sustainable alternative to produce high-value products. Previously, we have shown that abiotic stress provoked by nitrogen limitation enhances PhGs accumulation and *de novo* biosynthesis of anthocyanins through phenylalanine ammonia-lyase (PAL) activity. However, that condition had a severe negative effect in biomass thus, decreasing possibilities of commercial scale (Medina, 2011). Fungal elicitation of plant cell cultures is a strategy for promoting specialized metabolism having its basis on plant defense responses. Oligosaccharides originating from fungal cell walls are potent inducers of resistance of many plants species against a wide range of pathogens since they act as primary signals in the induction of plant defense reactions. In cell suspension cultures of *Cistanche deserticola* (Orobanchaceae) fungal elicitors derived from *Fusarium solani* (Lu and Mei, 2003), chitosan (Cheng *et al.*, 2006) or yeast (Cheng *et al.*, 2005) presented a positive effect on the production of PhGs. Since PhGs biosynthesis appears to be influenced by both biotic and abiotic (Sánchez-Rangel *et al.*, 2014) factors, we hypothesized that fungal elicitation may enhance biosynthesis of PhGs in *C. tenuiflora*. Here we tested the effects of cell wall oligosaccharides (CWOs) from *Fusarium oxysporum* f. sp. *lycopersici* race 3 in PhGs accumulation by *C.*

*tenuiflora* shoot cultures and the activity of PAL and peroxidase (POD), two defense-related enzymes.

## 2 Materials and methods

### 2.1 Plant culture and experimental design

Shoots culture of *C. tenuiflora* was initiated and propagated in vitro as described previously (Trejo-Tapia *et al.*, 2012). Shoots were subcultured in 300-ml shake flasks containing 50 mL B5 culture medium (Gamborg *et al.*, 1968), 3% (w/v) sucrose and no plant growth regulators. Cultures were maintained under continuous agitation at 110 rpm on an orbital shaker in a growth room at  $25 \pm 2^\circ\text{C}$  under a 16-h light/8-h dark photoperiod with illumination of  $103 \mu\text{mol/m}^2/\text{s}$  provided by 75 W cool-white fluorescent lamps. For elicitation treatments, preliminary assays were made and the elicitation time, 15 days, was defined. *C. tenuiflora* shoot cultures were grown in shake flasks for 15 days, then, were supplemented with CWOs from *Fusarium oxysporum* f. sp. *lycopersici* as elicitor.

### 2.2 Elicitation of shoots culture

The fungal strain used to obtain the CWOs for elicitation was *Fusarium oxysporum* f. sp. *lycopersici* race 3 (the strain was isolated from tomato and identified by molecular biology techniques (Ortega-García 2010)). The elicitor was prepared as described by El Modafar *et al.* (2001) with slight modifications. The pathogen was grown on potato dextrose broth under agitation at 150 rpm. Five-days-old mycelia from fungi was centrifuged at  $13,000 \times g$  for 40 min, the collected mycelium was harvested and rinsed thrice with sterile distilled water, then, dried at  $47^\circ\text{C}$ , grinded with a mortar and pestle and suspended in deionized water (2 mg/mL). This suspension was autoclaved at  $120^\circ\text{C}$  for 40 min and centrifuged at  $1,000 \times g$  for 20 min. The supernatant (CWOs) was used as elicitor and supplied to 15-day-old *C. tenuiflora* shoot cultures at 5 (+CWOs-L; lowest dose) and 13 (+CWOs-H; highest dose)  $\mu\text{g/mL}$  doses, based on the total carbohydrate content of the fungal homogenate, which was determined by the phenol/sulfuric acid method (Dubois *et al.*, 1956) using glucose as the standard. Shoots were collected for analysis at 1-8 days after elicitation.

### 2.3 Chlorophyll content

The chlorophyll extraction was performed with acetone (80%) and the absorbances were measured at 663.2 and 648.8 nm. The total chlorophyll content was calculated using the equations described by Lichtenthaler (1987).

### 2.4 Phenylalanine ammonia-lyase (PAL) activity

PAL was extracted and its activity was determined according to the methods described by Yan *et al.* (2006). Fresh samples (0.2 g) were frozen in liquid nitrogen and ground in a mortar with 40 mg of polyvinyl polypyrrolidone. The powder was mixed with 3 mL of a solution comprising 100 mM sodium phosphate buffer (pH 6.0), 2 mM EDTA, and 4 mM DTT. The mixture was centrifuged at  $13,500 \times g$  at 4°C for 15 min, and the supernatant was employed for the enzymatic assay. The protein concentration was determined using the method of Bradford (1976). The reaction mixture comprising 550  $\mu$ L of 50 mM Tris-HCl (pH 8.8) and 200  $\mu$ L of extract (2-16  $\mu$ g of protein) was preincubated at 40°C for 60 min and the reaction was initiated by addition of 250  $\mu$ L of 20 mM L-phenylalanine (pH 8.8). After 1 h of incubation at 40°C, the reaction was stopped by addition of 50  $\mu$ L of 5 N HCl. The blank reaction for every sample was carried out with Tris-HCl and extract but without phenylalanine. The absorbance change at 290 nm was evaluated after 1 h. The enzymatic activity was reported as  $\mu$ mol cinnamic acid  $\text{h}^{-1} \text{mg}^{-1}$  protein.

### 2.5 Guaiacol peroxidase (POD) activity

POD activity was measured according to the method of Stasolla and Yeung (2007). The reaction mixture (1 mL) contained 100 mM sodium phosphate buffer (pH 6.0), 0.3% hydrogen peroxide, 100 mM guaiacol, 2 mM EDTA, and 4 mM DTT, and enzymatic extract (2-16  $\mu$ g of protein). The change in absorbance at 470 nm caused by guaiacol oxidation was followed at 30-s intervals for 6 min. The POD activity was expressed as mM tetraguaiacol  $\text{min}^{-1} \text{mg}^{-1}$  protein using the extinction coefficient of tetraguaiacol (26.6  $\text{mM}^{-1}\text{cm}^{-1}$ ).

### 2.6 Evaluation of total phenolic compounds (TPC)

Fresh plant material (0.2 g) was ground and extracted with methanol (1 mL) for 24 h with agitation. The extract was filtered through Whatman paper No. 1, and concentrated in a rotary evaporator (Büchi-490; Büchi, Switzerland) under reduced pressure (210 mbar) at 40°C for 50 min. The residue was suspended in 10 mL of water and freeze-dried. Phenolic compounds in the methanol extract were estimated using the Folin-Ciocalteu reagent. Results were expressed as mg of gallic acid equivalents (GAE) per g of dry weight (DW) (Cardenas-Sandoval *et al.*, 2012). The samples were analyzed in triplicate.

### 2.7 Quantification of phenylethanoid glycosides (PhGs) by HPLC

PhGs were quantified by HPLC using a Waters 2695 separation module HPLC system equipped with a Waters 996 photodiode array detector and Empower Pro software (Waters Corporation, USA) as described previously (Gómez-Aguirre *et al.*, 2012). Compounds were separated on a LiChrospher 100 RP-18 column (4 mm  $\times$  250 mm i.d., 5- $\mu$ m particle size) (Merck, Darmstadt, Germany) connected to a guard column. The mobile phase consisted of water (pH 3.5) containing 5% TFA (solvent A) and acetonitrile (solvent B). The gradient system was as follows: 0-2 min, 100-0% B; 2-10 min, 77-23% B; 12-13 min, 70-30% B; 14-15 min, 100% B; 16-17 min, 0% B. The flow rate was maintained at 1  $\text{mL min}^{-1}$  and the injection volume was 20  $\mu$ L. The absorbance was measured at 330 nm. The PhGs peaks were identified by comparison of the retention times and UV spectra with those of reference compounds isolated and purified from the roots of *C. tenuiflora* (Gómez-Aguirre *et al.*, 2012). The retention times for isoverbascoside and verbascoside were 7.5 min ( $\lambda = 218, 247, 292, \text{ and } 331 \text{ nm}$ ) and 7.9 min ( $\lambda = 218, 247, 290, \text{ and } 330 \text{ nm}$ ), respectively. The amounts of PhGs were estimated by interpolation of the peak areas and comparison with a calibration curve constructed for verbascoside. The calibration curve was linear in the range of 62.5-500  $\mu$ g verbascoside  $\text{mL}^{-1}$  in methanol ( $y = 22,144x - 16,1443$ ;  $R^2 = 0.999$ ). All analyses were performed in triplicate and the data were expressed as mean values in  $\text{mg g}^{-1}$  DW.

Table 1. Growth parameters of *C. tenuiflora* shoots under control conditions or elicitor-treated cultures with two doses of CWOs from *F. oxysporum*

	Fresh biomass (g per flask)	Shoot length (cm)	Chlorophyll ( $\mu\text{g/g DW}$ )	Tissue water content (%)
Control	$11.76 \pm 0.01^A$	$3.36 \pm 0.17^B$	$1.05 \pm 0.01^A$	$93.36^A$
+CWOs-L	$11.82 \pm 0.01^A$	$3.49 \pm 0.01^B$	$0.79 \pm 0.01^B$	$93.20^A$
+CWOs-H	$11.70 \pm 0.01^A$	$4.25 \pm 0.22^A$	$0.78 \pm 0.01^B$	$93.08^A$

Elicitor were added after 15 days of growth and shoots were cultivated for further 8 days. Data represent means  $\pm$  standard errors ( $n = 9$ ). Within each column, different letters indicate significant difference according to Tukey's test ( $\alpha = 0.05$ ).

## 2.8 Statistical analysis

Statistical analyses were performed by one-way ANOVA and the all-pairwise multiple comparison procedure of Tukey was employed. The significance level for all statistical analyses was 5%. Differences of  $P < 0.05$  were considered significant. The software SigmaPlot for Windows version 11.0 software (Systat Software Inc., San Jose, CA, USA) was used to perform the statistical analyses.

## 3 Results

### 3.1 Effects of fungal elicitor on growth and enzymes activities

The effects of two doses, 5 and 13  $\mu\text{g/mL}$  of elicitor, on growth and enzyme activities were tested in shoots of *C. tenuiflora*. Neither of the doses tested altered biomass nor tissue water content. Elicitor-treated shoots with the highest dose (+CWOs-H, 13  $\mu\text{g/mL}$ ) were significantly longer than untreated shoots or shoots treated with the lowest dose (+CWOs-L, 5  $\mu\text{g/mL}$ ). Elicited shoots presented yellowish of their leaves (chlorosis) and total chlorophyll content was significantly reduced in these shoots compared with untreated shoots (Table 1). Elicitor application delayed root formation compared to the control. Maximum POD activity was higher in +CWOs-H shoots compared with untreated shoots or +CWOs-L shoots, from days 1 to 8 after elicitation (Fig. 1A).

PAL activity observed in untreated shoots and +FE-L shoots was 1.5-1.7  $\mu\text{mol CA/h/mg protein}$  and decreased at days 2 and 3 (0.83  $\mu\text{mol CA/h/mg protein}$ ) (Fig. 1B). Meanwhile, the maximum PAL activity in +FE-H elicited shoots was reached 4 days after elicitation (2.67  $\mu\text{mol CA/h/mg protein}$ ). PAL activity observed at this point was significantly higher in +FE-H shoots ( $p < 0.05$ ) than in untreated shoots and +FE-L shoots (1.67  $\mu\text{mol CA/h/mg protein}$ ).




Fig. 1. Time course of POD (A) and PAL (B) activity in *C. tenuiflora* control and elicitor-treated cultures with two doses of CWOs from *F. oxysporum*. Elicitor was added after 15 days of growth and shoots were cultivated for further 8 days. Data represents means  $\pm$  standard errors ( $n = 5$ ). Within each time point assayed, (\*) indicates significant difference according to Tukey's test ( $\alpha = 0.05$ ).

### 3.2 Contents of TPC and PhGs

Time-course accumulation of total phenolic content in *C. tenuiflora* after elicitation with the two doses of CWOs is shown in Fig. 2. In this case, the effect turned out to be different depending on the CWOs dose. TPC on untreated shoots ranged from 21.6-49.5 mg GAE/g DW; in shoots elicited with 5  $\mu\text{g CWOs/mL}$  were 30.8-






Fig. 2. Time course of total phenolic compounds content in *C. tenuiflora* control and elicitor-treated cultures with two doses of CWOs from *F. oxysporum*. Elicitor was added after 15 days of growth and shoots were cultivated for further 8 days. Data represents means  $\pm$  standard errors ( $n = 9$ ). Within each time point assayed, (\*) indicates significant difference according to Tukey's test ( $\alpha=0.05$ ).

51.4 mg GAE/g DW. In shoots elicited with the highest dose, the content of phenolic compounds increased from days 1 (60.2 mg GAE/g DW) to 8 of elicitation (91.8 mg GAE/g DW), reaching the maximum at day 4 (97.1 mg GAE/g DW) (Fig. 2).

Out of the two elicitor doses tested, the highest yield in TPC and the changes in enzymes activities were found in shoots treated with 13  $\mu\text{g}$  FE/mL (+FE-H), thus, further experiments were done only with shoots treated with this dose of elicitor. According to HPLC analyses *C. tenuiflora* shoots accumulated the PhGs verbascoside (1,  $t_R = 7.9$  min,  $\lambda_{max} = 200, 330$  nm) and isoverbascoside (2,  $t_R = 7.5$  min,  $\lambda_{max} = 200, 330$  nm) and, a yet unidentified phenylethanoid glycoside (3,  $t_R = 6.3$  min,  $\lambda_{max} = 200, 330$  nm) (Fig. 3). The HPLC profile suggested that verbascoside was the main PhGs in untreated shoots and fungal elicited.




Fig. 3. Chromatographic HPLC profiles of methanol extracts from *C. tenuiflora* control and elicitor-treated cultures with two doses of CWOs from *F. oxysporum*. Elicitor was added after 15 days of growth and shoots were cultivated for further 8 days. Verbascoside (1,  $t_R = 7.9$  min,  $\lambda_{max} = 200, 330$  nm), Isoverbascoside (2,  $t_R = 7.5$  min,  $\lambda_{max} = 200, 330$  nm) and unidentified phenylethanoid glycoside (3,  $t_R = 6.3$  min,  $\lambda_{max} = 200, 330$  nm).

Table 2. Phenylethanoids glycosides concentration in *C. tenuiflora* shoots under control conditions or elicitor-treated cultures with 13  $\mu\text{g/mL}$  (+CWOs-H) from *F. oxysporum*

	Verbascoside (mg/g DW)	Isoverbascoside (mg/g DW)
Control	19.19 $\pm$ 0.2 <sup>B</sup>	14.11 $\pm$ 0.1 <sup>B</sup>
+CWOs-H	95.66 $\pm$ 0.8 <sup>A</sup>	54.29 $\pm$ 0.3 <sup>A</sup>

Elicitor was added after 15 days of growth and shoots were cultivated for further 8 days. Data represents means  $\pm$  standard errors ( $n = 5$ ). Within each column, different letters indicate significant difference according to Tukey's test ( $\alpha=0.05$ ).

Eight days after elicitation, the specific concentration of verbascoside in fungal elicited shoots was 95.66 mg/g DW compared with 19.19 mg/G DW in untreated shoots (Table 2). Isoverbascoside was also enhanced by fungal elicitation, but to a lesser extent (14.11 vs 54.29 mg/g DW).

## 4 Discussion

Elicitation with CWOs from *F. oxysporum* did not modify *C. tenuiflora* growth measured as biomass. However, elicitation provoked that leaves turned yellow (chlorosis), total chlorophylls decreased, shoots elongated and root formation was delayed. This response was similar to reported by (Boari and Vurro, 2004) where, development of shoots from *Orobancha ramosa* (Orobanchaceae) was severely affected by *F. oxysporum* and *F. solani*. Similarly, *F. oxysporum* has been used in growth control of *Striga hermonthica* (Del). (Orobanchaceae) since it can cause plant death (Ndambi *et al.*, 2011). In contrast, growth of cell suspension cultures of *Cistanche deserticola* (Orobanchaceae) was not affected by elicitation with *F. solani* (carbohydrate equivalents)(40 mg/L) (Lu and Mei, 2003). Oligosaccharides (galactoglucomannans) inhibited growth, morphology and adventitious root formation of *Karwinskia humboldtiana* (Rhamnaceae) (Kollárová *et al.*, 2007). That effect depended on oligosaccharides concentration. Similarly, hypocotyl elongation of *Vigna radiata* (L.) Wilczek (Fabaceae) was negatively affected by oligosaccharides (Kollárová *et al.*, 2009). Those changes were explained based on the effect of oligosaccharides on auxins metabolism and thus, influence in plant growth regulation. In other species,

exposure to oligosaccharides provoked oxidative stress condition (Orozco-Cárdenas *et al.*, 2001) and chlorosis (De Wit *et al.*, 1984).

TPC was used as a first evidence of the effect of elicitation on secondary metabolism; changes were observed since the first 24 h with the highest TPC at fourth day. Based on this result, the influence of fungal elicitation on enzymes activities and HPLC profile was analyzed. Fungal elicitation provoked changes in the activities of POD and PAL compared with the control. POD is a pathogen-related enzymes associated with the plant defense pathway that is challenged by elicitors (Dutsadee and Nunta, 2008) and used as an indicative of an oxidative stress condition (Arencibia *et al.*, 2012). Fungal elicitation with CWOs from *F. oxysporum* provoked an evident increase (20-fold) in *C. tenuiflora* POD activity compared with non-elicited shoots. A similar response was recorded in *Lupinus luteus* (Fabaceae) embryos inoculated with *F. oxysporum* spore suspension, where POD activity increased between 0 to 96 h after inoculation (Morkunas and Gmerek, 2007). PAL is a key branch enzyme on the biosynthesis of phenolic compounds such as phenolic acids, flavonoids or anthocyanins. The increase in its activity is an indicator of plant resistance. In *C. tenuiflora*, PAL activity increased in response to elicitor with a differential effect depending of dose. With the highest dose, increase was obvious at 2 days with a maximum at 4 days (1.7-fold compared with the control). While with the lowest dose, the increase was evident at 3 days. PAL activity agrees with the reports on *Taxus chinensis* suspension cultures (Yuan *et al.*, 2002) and *C. deserticola* (Lu and Mei, 2003) elicited with *F. oxysporum* and *F. solani*, respectively. Elicitation with oligosaccharides isolated from *F. oxysporum* induced PAL activity in *Rubus fruticosus* cells (Nita-Lazar *et al.*, 2004). Plant cell response to pathogen attack or other stress condition involves events at biochemical and molecular level occurring at different times (Zhao *et al.*, 2005). Reactive Oxygen Species (ROS) production constitutes one of the earliest events while secondary metabolism accumulation occurs later. This may explain that maximum POD activity preceded maximum PAL activity. Other authors explain that POD activity is link to lignin biosynthesis, which is also associated to PAL activity. Thus, POD activation indirectly participates in PAL activation as one of the control points for synthesis of phenolic compounds (Arencibia *et al.*, 2012; Jouili *et al.*, 2011).

Fungal elicitation clearly enhanced PhGs accumulation. Shoots elicited with the highest FE

dose accumulated 5-fold verbascoside than non-elicited shoots. PhGs enhancement agrees with that observed in cell suspension cultures of *Cistanche deserticola* elicited with *F. solani* oligosaccharides (Lu and Mei, 2003), chitosan (Cheng et al., 2006) or a yeast-derived elicitor (Cheng et al., 2005). PhGs role in plants has been little explored since researches on their pharmacological activities have received more attention. Verbascoside influences fungi morphology and decreases sporulation of *Fusarium culmorum*, *Bipolaris sorokiniana* and *Botrytis cinerea* (Nikonorova et al., 2009). Also inhibit growth of *Penicillium digitatum* (Oyourou et al., 2013). The effects of fungal elicitation on PhGs biosynthesis have not been tested in previous studies. Verbascoside and isoverbascoside were significantly increased after fungal elicitation, suggesting antioxidant protective roles of these compounds against the oxidative stress caused by *F. oxysporum*. This possibility is under further investigation.

## Conclusions

Fungal elicitation with WCOs from *F. oxysporum* increases PhGs accumulation in *C. tenuiflora* via the stimulation of POD and PAL and, without reduction in biomass.

## Acknowledgements

B.A. Cardenas-Sandoval is indebted to Consejo Nacional de Ciencia y Tecnología (CONACYT-MEXICO) and PIFI-IPN for the fellowship awarded as graduate student. Financial support from Secretaría de Investigación y Posgrado del IPN (grant 20140231) and by Consejo Nacional de Ciencia y Tecnología-México (grant 220007).

## Notation

CWOs DW	cell wall oligosaccharides dry weight
FE	fungal elicitation
GAE	gallic acid equivalents
PhGs	phenylethanoid glycosides
PAL	phenylalanine ammonia-lyase
POD	guaiacol peroxidase
ROS	reactive oxygen species
TPC	total phenolic compounds

## References

- Arencibia, A. D., Bernal, A., Zayas, C., Carmona, E., Cordero, C., González, G., García, R. and Santana, I. (2012). Hydrogen peroxide induced phenylpropanoids pathway eliciting a defensive response in plants micropropagated in Temporary Immersion Bioreactors (TIBs). *Plant Science* 195, 71-79.
- Boari, A. and Vurro, M. (2004). Evaluation of *Fusarium* spp. and other fungi as biological control agents of broomrape (*Orobancha ramosa*). *Biological Control* 30, 212-219.
- Bradford, M.M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72, 248-254.
- Cardenas-Sandoval, B.A., López-Laredo, A.R., Martínez-Bonfil, B.P., Bermúdez-Torres, K. and Trejo-Tapia, G. (2012). Advances in the phytochemistry of *Cuphea aequipetala*, *C. aequipetala* var. *hispida* and *C. lanceolata*: extraction and quantification of phenolic compounds and antioxidant activity. *Revista Mexicana de Ingeniería Química* 11, 401-413.
- Cheng, X., Wei, T., Guo, B., Ni, W. and Liu, C.-Z. (2005). *Cistanche deserticola* cell suspension cultures: Phenylethanoid glycosides biosynthesis and antioxidant activity. *Process Biochemistry* 40, 3119-3124.
- Cheng, X., Zhou, H.-Y., Cui, X., Ni, W. and Liu, C.-Z. (2006). Improvement of phenylethanoid glycosides biosynthesis in *Cistanche deserticola* cell suspension cultures by chitosan elicitor. *Journal of Biotechnology* 121, 253-260.
- De Wit, J.G.M.P., Hofman, J.E., and Aarts J.M.M.J.G. (1984). Origen of specific elicitors of chlorosis and necrosis occurring in intracellular fluids of compatible interactions of *Cladosporium fulvum* (syn. *Fulvia fulva*) and tomato. *Physiological Plant Pathology* 24, 17-23.
- Dubois, M., Gilles, K.A. Hamilton, J.K. Rebers, P.A. and Smith, F. 1956. Colorimetric method for determination of sugars and related substances. *Analytical Chemistry* 28, 350- 356.



- Dutsadee, C. and Nunta, C. (2008). Induction of peroxidase, scopoletin, phenolic compounds and resistance in *Hevea brasiliensis* by elicitor and a novel protein elicitor purified from *Phytophthora palmivora*. *Physiological and Molecular Plant Pathology* 72, 179-187.
- El Modafar, C., Tantaoui, A. and El Boustani, E.-S. (2001). Differential induction of phenylalanine ammonia-lyase activity in date palm roots in response to inoculation with *Fusarium oxysporum* f. sp. *albedinis* and to elicitation with fungal wall elicitor. *Journal of Plant Physiology* 158, 715-722.
- Funari, C. S., Gullo, F.P., Napolitano, A., Carneiro, R.L., Mendes-Giannini, M.J.S., Fusco-Almeida, A. M., Piacente, S., Pizzi, C. and Silva, D. H. S. (2012). Chemical and antifungal investigations of six *Lippia* species (Verbenaceae) from Brazil. *Food Chemistry* 135, 2086-2094.
- Gamborg, O. L., Miller, R. A. and Ojima, K. (1968). Nutrient requirements of suspension cultures of soybean root cells. *Experimental Cell Research* 50, 150-158.
- Gómez, Y. A. (2011). Identificación estructural de compuestos mayoritarios en plantas silvestres de *Castilleja tenuiflora* y su acumulación en cultivos *in vitro* de raíces. Tesis de Doctorado en Ciencias en Desarrollo de Productos Bióticos, Instituto Politécnico Nacional, Yautepec, Morelos, México.
- Gómez-Aguirre, Y. A., Zamilpa, A., González, M. and Trejo-Tapia, G. (2012). Adventitious root cultures of *Castilleja tenuiflora* Benth. as a source of phenylethanoid glycosides. *Industrial Crops and Products* 36, 188-195.
- Gyurkovska, V., Alipieva, K., Maciuk, A., Dimitrova, P., Ivanovska, N., Haas, C., Bley, T. and Georgiev, M. (2011). Anti-inflammatory activity of Devil's claw *in vitro* systems and their active constituents. *Food Chemistry* 125, 171-178.
- Jouili, H., Bouazizi, H. and El Ferjani, E. (2011). Plant peroxidases: biomarkers of metallic stress. *Acta Physiologiae Plantarum* 33, 2075-2082.
- Kollárová, K., Lišková, D. and Lux, A. (2007). Influence of galactoglucomannan oligosaccharides on root culture of *Karwinskia humboldtiana*. *Plant Cell, Tissue and Organ Culture* 91, 9-19.
- Kollárová, K., Richterová, D., Slováková, L., Henselová, M., Capek, P. and Lisková, D. (2009). Impact of galactoglucomannan oligosaccharides on elongation growth in intact mung bean plants. *Plant Science* 177, 324-330.
- Lichtenthaler, H. K. (1987). Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. In: *Methods in Enzymology*, vol. 148 (R.D. Lester Packer ed.), Pp. 350-382 Academic Press.
- López-Laredo, A., Gómez-Aguirre, Y., Medina-Pérez, V., Salcedo-Morales, G., Sepúlveda-Jiménez, G. and Trejo-Tapia, G. (2012). Variation in antioxidant properties and phenolics concentration in different organs of wild growing and greenhouse cultivated *Castilleja tenuiflora* Benth. *Acta Physiologiae Plantarum* 34, 2435-2442.
- Lu, C.-t. and Mei, X.-g. (2003). Improvement of phenylethanoid glycosides production by a fungal elicitor in cell suspension culture of *Cistanche deserticola*. *Biotechnology Letters* 25, 1437-1439.
- Martínez-Bonfil, B., Salcedo-Morales, G., López-Laredo, A., Ventura-Zapata, E., Evangelista-Lozano, S. and Trejo-Tapia, G. (2011). Shoot regeneration and determination of iridoid levels in the medicinal plant *Castilleja tenuiflora* Benth. *Plant Cell, Tissue and Organ Culture* 107, 195-203.
- Medina, V. (2011). Efecto de la reducción de nitrógeno en el metabolismo de compuestos fenólicos en brotes de *Castilleja tenuiflora* Benth. cultivados en biorreactor de inmersión temporal. Tesis de Maestría en Ciencias en Desarrollo de Productos Bióticos, Instituto Politécnico Nacional, Yautepec, Morelos, México.
- Moreno-Escobar, J. A., Bazaldúa, S., Villarreal, M. L., Bonilla-Barbosa, J. R., Mendoza, S. and Rodríguez-López, V. (2011). Cytotoxic and antioxidant activities of selected Lamiales species from Mexico. *Pharmaceutical Biology* 49, 1243-1248.

- Morkunas, I. and Gmerek, J. (2007). The possible involvement of peroxidase in defense of yellow lupine embryo axes against *Fusarium oxysporum*. *Journal of Plant Physiology* 164, 185-194.
- Ndambi, B., Cadisch, G., Elzein, A. and Heller, A. (2011). Colonization and control of *Striga hermonthica* by *Fusarium oxysporum* f. sp. strigae, a mycoherbicide component: An anatomical study. *Biological Control* 58, 149-159.
- Nikonorova, A. K., Egorov, C. A., Galkina, T. G., Grishin, E. V. and Babakov, A. V. (2009). Antifungal activity of phenolic glycoside verbascoside from *Plantago major* seeds. *Mikologiya i Fitopatologiya* 43, 52-57.
- Nita-Lazar, M., Heyraud, A., Gey, C., Braccini, I. and Lienart, Y. (2004). Novel oligosaccharides isolated from *Fusarium oxysporum* L. rapidly induce PAL activity in *Rubus* cells. *Acta Biochimica Polonica* 51, 625-634.
- Orozco-Cárdenas, M. L., Narváez-Vásquez, J. and Ryan, C. A. (2001). Hydrogen peroxide acts as a second messenger for the induction of defense genes in tomato plants in response to wounding, systemin, and methyl jasmonate. *The Plant Cell* 13, 179-191.
- Ortega García, J. G. (2010). Diagnóstico de hongos fitopatógenos de jitomate y efecto de *Trichoderma asperellum* Tc74 sobre *Fusarium* spp. Tesis de maestría en Ciencias en Manejo Agroecológico de Plagas y Enfermedades, Instituto Politécnico Nacional, Yautepec, Morelos, México.
- Oyourou, J. N., Combrinck, S., Regnier, T. and Marston, A. (2013). Purification, stability and antifungal activity of verbascoside from *Lippia javanica* and *Lantana camara* leaf extracts. *Industrial Crops and Products* 43, 820-826.
- Pettit, G. R., Numata, A., Takemura, T., Ode, R. H., Narula, A. S., Schmidt, J. M., Cragg, G. M. and Pase, C. P. (1990). Antineoplastic agents, 107. Isolation of acteoside and isoacteoside from *Castilleja linariaefolia*. *Journal of Natural Products* 53, 456-458.
- Sánchez, P. M., Villarreal, M. L., Herrera-Ruiz, M., Zamilpa, A., Jiménez-Ferrer, E. and Trejo-Tapia, G. (2013). *In vivo* anti-inflammatory and anti-ulcerogenic activities of extracts from wild growing and *in vitro* plants of *Castilleja tenuiflora* Benth. (Orobanchaceae). *Journal of Ethnopharmacology* 150, 1032-1037.
- Sánchez-Rangel, J.C., Benavides, J. and Jacobo-Velázquez, D.A. (2014). Abiotic stress based bioprocesses for the production of high value antioxidant phenolic compound in plants: an overview. *Revista Mexicana de Ingeniería Química* 13, 49-61.
- Singh, N., Shukla, N., Singh, P., Sharma, R., Rajendran, S. M., Maurya, R. and Palit, G. (2010). Verbascoside isolated from *Tectona grandis* mediates gastric protection in rats via inhibiting proton pump activity. *Fitoterapia* 81, 755-761.
- Stasolla, C. and Yeung, E. C. (2007). Cellular ascorbic acid regulates the activity of major peroxidases in the apical poles of germinating white spruce (*Picea glauca*) somatic embryos. *Plant Physiology and Biochemistry* 45, 188-198.
- Trejo-Tapia, G., Rosas-Romero, G., López-Laredo, A. R., Bermúdez-Torres, K. and Zamilpa, A. (2012). *In vitro* organ cultures of the cancer herb *Castilleja tenuiflora* Benth. as potential sources of iridoids and antioxidant compounds. In: *Biotechnological Production of Plant Secondary Metabolites*, (I. Orhan ed.), Pp. 87-106 Bentham Science Publishers.
- Yan, Q., Shi, M., Ng, J. and Wu, J. Y. (2006). Elicitor-induced rosmarinic acid accumulation and secondary metabolism enzyme activities in *Salvia miltiorrhiza* hairy roots. *Plant Science* 170, 853-858.
- Yuan, Y.-J., Li, C., Hu, Z.-D., Wu, J.-C. and Zeng, A.-P. (2002). Fungal elicitor-induced cell apoptosis in suspension cultures of *Taxus chinensis* var. mairei for taxol production. *Process Biochemistry* 38, 193-198.
- Zhao, J., Davis, L. C. and Verpoorte, R. (2005). Elicitor signal transduction leading to production of plant secondary metabolites. *Biotechnology Advances* 23, 283-333.