



Pesquisa Brasileira em Odontopediatria e
Clínica Integrada

ISSN: 1519-0501

apesb@terra.com.br

Universidade Federal da Paraíba
Brasil

ORDOBAZARI, Morteza; DAVOUDIAN, Masoude; ORDOBAZARI, Atousa
Comparison between Manual and Software Surgical Planning Predictions in Orthognathic Surgery
Pesquisa Brasileira em Odontopediatria e Clínica Integrada, vol. 10, núm. 2, mayo-agosto, 2010, pp.
291-295

Universidade Federal da Paraíba
Paraíba, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=63716962023>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

Comparison between Manual and Software Surgical Planning Predictions in Orthognathic Surgery

Comparação Entre os Prognósticos do Planejamento Cirúrgico Manual ou por Software em Cirurgia Ortognática

Morteza ORDOBAZARI¹, Masoude DAVOUDIAN², Atousa ORDOBAZARI³

¹Professor of Department of Orthodontics, School of Dentistry, Shaheed Beheshti University, Tehran, Iran.

²Assistant Professor, Department of Orthodontics, School of Dentistry, Kerman University, Kerman, Iran.

³Physician, California Pacific Medical Center (APRL), San Francisco, California, USA.

RESUMO

Objetivo: Avaliar comparativamente o prognóstico do planejamento cirúrgico por software com o método manual.

Método: Radiografias laterais cefalométricas e modelos de estudo pré e pós-cirurgia de vinte pacientes Classe III (12 mulheres e 8 homens, média de idade: $22,5 \pm 2,3$) que tinham se submetidos à cirurgia no Departamento de Ortodontia. Para o planejamento manual, com combinação de oclusogramas pré e pós-cirurgia com radiografias cefalométricas laterais de cada paciente, foram feitas análises tridimensionais. Prognósticos da movimentação esquelética foram elaborados. Em seguida, mensurou-se tridimensionalmente a movimentação esquelética. Quinze variáveis angulares e lineares foram avaliadas e os resultados analisados estatisticamente. Para cada paciente, o prognóstico manual e por software antes e após a cirurgia foram comparados entre si. O teste t pareado foi usado para verificar possíveis diferenças ($p < 0,05$).

Resultados: Não foram observadas diferenças entre os métodos.

Conclusão: O Software Dolphin Imaging (versão 10) tem boa acurácia para a predição do resultado pré e pós-cirúrgico de cirurgias ortognáticas.

ABSTRACT

Objective: To evaluate the proximity of computer imaging software surgical planning prediction methods, with standard manual methods.

Method: Lateral cephalometry radiographs and study cast records of pre and post surgery of twenty longface CI III patients (12 Females and 8 Males, mean age: 22.5 ± 2.3), that had been undergone surgery in Department of Orthodontics. For manual prediction, with combination of pre and post surgical occlusograms with lateral cephalometric radiograph tracings of each samples, three dimensional analysis was done. Predictions of skeletal movement with tacing overlay method, has prepared. Finally, measured the rate of skeletal movements at in three dimensional of spaces. Fifteen angular and linear variables were measured, and the results, analyzed using the statistical software statistics. For each patient, the manual and computer software prediction before and after surgery were compared with each other. The paired t-test was used to evaluate possible differences between manual and computer software prediction ($p < 0.05$).

Results: The difference of the variables of the software prediction in the pre and postsurgical stages in comparison with standard manual prediction method was insignificant.

Conclusion: Dolphin Imaging software (version 10.0) has a good accuracy for prediction of pre and postsurgical outcome of orthognathic samples.

DESCRIPTORES

Procedimentos cirúrgicos ortognáticos; Validação de programas de computador; Má oclusão de Angle classe III.

KEYWORDS

Orthognathic surgical procedures; Software validation; Malocclusion, Angle class III.

INTRODUCTION

The need for orthognathic surgery to correction of skeletal deformities is increasing, because recently, more adult people are going to do orthodontic treatments. Prediction of the outcome of orthognathic surgery is very important in treatment of these deformities, that usually has been performed with manual tracings¹.

Recently technologic advancement induce using computer, in planning and prediction of the outcome of orthogenetic surgery treatments¹⁻³. Several investigations has been performed to compare the accuracy of manual and software-aided predictions⁴⁻⁶.

The aim of this study was determining the proximity between predictions performed with Dolphin imaging software (version 10.0) and the standard manual method pre and postsurgery^{7,8}.

MATERIALS AND METHODS

Lateral cephalometry radiographs and study cast records of pre and post surgery of twenty longface CI III samples, that had been undergone orthognathic surgery were prepared. Casts of each patient were mounted in centric occlusion (Figure 1).



Figure 1. Mounted upper and lower casts in maximum intercuspation.

With use of Xerox copy, an image of the upper and lower occlusal aspect view of dental arches (occlusogram) was prepared (Figures 2 and 3). Combination of pre and post surgical occlusograms with lateral cephalometry radiographs tracing, three dimensional analysis was doing (Figure 4).




Figure 2. Occlusogram of the upper arch.



Figure 3. Occlusogram of the lower arch.





Figure 4. Combination of pre and post surgical occlusograms with lateral cephalometry radiograph tracing.


Postsurgical outcome was predicted manually, shown in the Figures (5 to 10) as following: Presurgical manual prediction (Figure 5); Incisor at rest line,(2 mm below the upper lip) (Figure 6); Autorotation of the mandible around the condyle (2 mm upper to rest line) (Figure 7); Reposition of the maxilla on mandible until A

point superimposes on N.Perp line (Figure 8); Reposition of the mandible to achieving maximum intercuspation

(Figure 9) and Postsurgical manual prediction (Figure 10).



Figures 5, 6 and 7. Presurgical manual prediction (Fig. 5), Incisor at rest line (2 mm below the upper lip) (Fig. 6) and Rotating the mandible around the condyle, 2 mm upper to rest line (Fig. 7).



Figures 8, 9 and 10. Reposition of maxilla on mandible until A superimposes on N.Perp line (Fig. 8), Reposition of mandible to achieving maximum intercuspation (Fig. 9) and Postsurgical manual prediction (Fig. 10).

Entering of the measurements from manual prediction tracing, in the treatment panel, was used

to predict postsurgical outcome with Dolphin imaging software, shown in the Figures 11 to15 as following:




Figure 11. Digitized cephalometry in Dolphin imaging software




Figure 12. Entering of the manual data to the treatment panel for prediction of presurgical outcome




Figure 13. Presurgical software prediction.




Figure 14. Entering of the manual data to the treatment panel for prediction of postsurgical outcome.




Figure 15. Postsurgical software prediction.

Based on literature review, the manual method was accepted as golden standard. For every patient, in the pre and postsurgical stages, the correlation between manual and software aided predictions were determined, respectively. For these comparisons, the paired t-test was computed ($p < 0.05$).

RESULTS

The below measurements were calculated for the real outcomes and predictions before and after the surgery: Angular measurements: U1 to SN, L1 to MEGO, L1 to A.pog; Linear measurements: U1/A.perp, L1/A.Pog., U1/PP, U6/PP, L1/MeGo, L6/MeGo.

Comparison between manual and software prediction of presurgical outcome: All variables were not statistically significant.

Comparison between manual and software prediction of postsurgical outcome: Except SNA, U1/A. Perp, other variables were not statistically significant.

DISCUSSION

While it is not confirmed through the repetitive examination of outcome measurements showing at least comparable or preferably improved accuracy when compared with that of the traditional manual technique, routine utilization of software aided cephalometric predictions cannot be justified. The data for prediction of pre and postsurgical stages of the manual and software-aided prediction were not significantly different, respectively.

Previous investigation on Quick ceph software, offered a reasonably accurate method of prediction⁹. The results of this study showed that Quick Ceph tended to overestimate horizontal landmark position and underestimate vertical changes. However, the mean differences and standard deviations were less than values reported for manually prepared prediction profiles.

Some researchers in their studies^{1,10} found similarity between manual and software method for the prediction of postsurgical outcome. Computer programs cannot consistently predict the skeletal changes occurring after orthognathic surgery but their results may be considered inside a clinically acceptable range¹¹. They stated last-minute changes by the surgeons as the meaningful cause of the errors in prediction. Previous study indicated that Version 8.0 of Dolphin Imaging Software needs to re-assessed for software errors that may result in clinically significant miscalculations, and to facilitate compensation of radiographic magnification when using linear measurements⁷. This finding was because of using surgeon manual data for movement of the jaws with the software during investigation.

In addition to difficulty in locating A point, according to others authors¹² positioning the apex of upper incisors in the cephalometric view is not so easy, therefore there

was significant difference for the SNA and U1/A.perp measurements in postsurgical comparisons.

The most important limitation in this study was probable imprecision in calculating the movements occurred in manual method based on different positions of surgical reference lines. The accuracy of the expert selecting the best profile would be another limiting factor in this investigation.

Recent advancement of computer technology in different clinical fields of medicine has made increasingly problematic to assess the efficacy of the technology before it becomes available in the market and used by numerous clinicians. Although a multitude of cephalometric analysis software programs have been commercially available for several years, there are few studies evaluating the reliability and the accuracy of measures by the programs. There are also few studies comparing differences across different versions of digital tracing softwares. It can be more easier for the practitioner to select technology that is reliable and accurate by referring to associated investigations.

This study provides a basis to assess other software programs. Further studies in this area will, no doubt, improve our ability to select the best program by more knowledge about the advantages and disadvantages of other softwares. But, for now, the orthodontic clinicians can use Dolphin imaging software (version 10) instead of traditional manual methods for prediction of orthognathic treatment of longface class III patients with clinically acceptable accuracy.

CONCLUSION

Dolphin Imaging software (version 10.0) has a good accuracy for prediction of presurgical and postsurgical outcome of longface CI III orthognathic patients.

REFERENCES

1. Gossett CB, Preston CB, Dunford R, Lampasso J. Prediction accuracy of computer-assisted surgical visual treatment objectives as compared with conventional visual treatment objectives. *J Oral maxillofac Surg* 2005; 63(5):609-17.
2. Lu CH, Ko EW, Huang CS. The accuracy of video imaging prediction in soft tissue outcome after bimaxillary orthognathic surgery. *J Oral Maxillofac Surg* 2003; 61(3):333-42.
3. Loh S, Yow M. Computer prediction of hard tissue profiles in orthognathic surgery. *Int J Adult Orthodon Orthognath Surg* 2002; 17(4):342-7.
4. Mankad B, Cisneros GJ, Freeman K, Eisig SB. Prediction accuracy of soft tissue profile in orthognathic surgery. *Int J Adult Orthodon Orthognath Surg* 1999; 14(1):19-26.

5. Sinclair PM, Kilpelainen P, Phillips C, White RP Jr, Rogers L, Sarver DM. The accuracy of video imaging in orthognathic surgery. *Am J Orthod Dentofacial Orthop* 1995; 107(2):177-85.
6. Smith JD, Thomas PM, Proffit WR. A comparison of current prediction imaging programs. *Am J Orthod Dentofacial Orthop* 2004; 125(5):527-36.
7. Power G, Breckon J, Sherriff M, McDonald F. Dolphin imaging software: An analysis of the accuracy of cephalometric digitization and orthognathic prediction. *Int J Oral Maxillofac Surg* 2005; 34(6):619-26.
8. Chunmaneechote P, Friede H. Mandibular setback osteotomy: facial soft tissue behavior and possibility to improve the accuracy of the soft tissue profile prediction with the use of a computerized cephalometric program: Quick Ceph Image Pro: v. 2.5. *Clin Orthod Res* 1999; 2(2):85-98.
9. Hing NR. The accuracy of computer generated prediction tracings. *Int J Oral Maxillofac Surg* 1989; 18(3):148-51.
10. Eckhardt CE, Cunningham SJ. How predictable is orthognathic surgery? *Eur J Orthod* 2006; 26(3):303-9.
11. Kaipatur N, Al-Thomali Y, Flores-Mir C. Accuracy of computer programs in predicting orthognathic surgery hard tissue response. *J Oral Maxillofac Surg* 2009; 67(8):1628-39.
12. Loh S, Heng JK, Ward-Booth P, Winchester L, McDonald F. A radiographic analysis of computer prediction in conjunction with orthognathic surgery. *Int J Oral Maxillofac Surg* 2001; 30(4):259-63.

Recebido/Received: 24/01/10
Revisado/Reviewed: 10/03/10
Aprovado/Approved: 15/04/10

Correspondência:

Professor Morteza Ordobazari
No. 4, 33 street, Alvand Avenue, Arjantin Square
Tehran Iran PC 1516666115
Telephone : +98 21 8877-2146
Fax : +98 21 8877-1909
E-mail: m_ordobazari@yahoo.com