



Vojnotehnicki glasnik/Military Technical
Courier

ISSN: 0042-8469

vojnotehnicki.glasnik@mod.gov.rs

University of Defence
Serbia

Abbas, Mujahid; Ali, Basit; Nazir, Talat; M. Dedovi, Nebojša; Bin-Mohsin, Bandar; N.
Radenovi, Stojan

Solutions and Ulam-Hyers stability of differential inclusions involving Suzuki type
multivalued mappings on b-metric spaces

Vojnotehnicki glasnik/Military Technical Courier, vol. 68, núm. 3, julio-septiembre, 2020,
pp. 438-487
University of Defence

Available in: <https://www.redalyc.org/articulo.oa?id=661770398003>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



SOLUTIONS AND ULAM-HYERS STABILITY OF DIFFERENTIAL INCLUSIONS INVOLVING SUZUKI TYPE MULTIVALUED MAPPINGS ON b -METRIC SPACES

Mujahid Abbas^a, Basit Ali^b, Talat Nazir^c, Nebojša M. Dedović^d, Bandar Bin-Mohsin^e, Stojan N. Radenović^f

^a Government College University, Department of Mathematics, Lahore, Islamic Republic of Pakistan;
University of Pretoria, Department of Mathematics and Applied Mathematics, Pretoria, Republic of South Africa,
e-mail: abbas.mujahid@gmail.com,
ORCID ID: <https://orcid.org/0000-0001-5528-1207>

^b University of Management and Technology, Department of Mathematics, Lahore, Islamic Republic of Pakistan,
e-mail: basit.aa@gmail.com,
ORCID ID: <https://orcid.org/0000-0003-4111-5974>

^c COMSATS University Islamabad, Department of Mathematics, Abbottabad Campus, Islamic Republic of Pakistan;
University of South Africa, Department of Mathematical Science, Science Campus, Johannesburg, Republic of South Africa,
e-mail: talat@ciit.net.pk,
ORCID ID: <https://orcid.org/0000-0001-6516-3212>

^d University of Novi Sad, Faculty of Agriculture, Department of Agricultural Engineering, Novi Sad, Republic of Serbia,
e-mail: nebojsa.dedovic@polj.uns.ac.rs, **corresponding author**,
ORCID ID: <https://orcid.org/0000-0002-4628-1405>

^e King Saud University, College of Science, Department of Mathematics, Riyadh, Kingdom of Saudi Arabia,
e-mail: balmohsen@ksu.edu.sa,
ORCID ID: <https://orcid.org/0000-0002-2160-4159>

^f University of Belgrade, Faculty of Mechanical Engineering, Belgrade, Republic of Serbia,
e-mail: radens@beotel.rs,
ORCID ID: <https://orcid.org/0000-0001-8254-6688>

DOI: 10.5937/vojtehg68-26718; <https://doi.org/10.5937/vojtehg68-26718>

FIELD: Mathematics

ARTICLE TYPE: Original Scientific Paper

ACKNOWLEDGMENT: The work of the fourth author Nebojša M. Dedović is supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia, project TR-37017.

Abstract:

Introduction/purpose: This paper presents coincidence and common fixed points of Suzuki type $(\alpha_* - \psi)$ – multivalued operators on b-metric spaces.

Methods: The limit shadowing property was discussed as well as the well-posedness and the Ulam-Hyers stability of the solution for the fixed point problem of such operators.

Results: The upper bound of the Hausdorff distance between the fixed point sets is obtained. Some examples are presented to support the obtained results.

Conclusion: The application of the obtained results establishes the existence of differential inclusion.

Keywords: b-metric space, multi-valued mapping, fixed point problems, Ulam-Hyers stability, initial value problem.

Introduction and preliminaries

Euclidean distance is an important measure of "nearness" between two real or complex numbers. Fréchet (1905) introduced the concept of a metric to obtain the distance between two arbitrary objects. Since then, this notion has been generalized further in one to many directions, see (An et al, 2015a), among which one of the most important generalizations is the concept of a b-metric initiated by (Czerwik, 1993). For more details of b-metric spaces see (Aleksić et al, 2018), (Hussain et al, 2012), (Kirk & Shahzad, 2014) and the references therein.

Definition 1.1 Let X be a nonempty set. A mapping $d: X \times X \rightarrow [0, +\infty)$ is said to be a b-metric on X if there exists some real constant $b \geq 1$ such that for any $x, y, z \in X$, the following condition holds:

- a₁):** $d(x, y) = 0$ if and only if $x = y$;
- a₂):** $d(x, y) = d(y, x)$;
- a₃):** $d(x, y) \leq bd(x, z) + bd(z, y)$

The pair (X, d) is termed a b-metric space with b-metric constant b . Every metric is b-metric for $b = 1$ but the converse does not hold in general (Ćirić et al, 2012), (Czerwik, 1993), (Singh & Prasad, 2008).

In the sequel, the letters, \mathbb{R}^+ , \mathbb{R} , \mathbb{N} and \mathbb{Z}^+ will denote the set of all nonnegative real numbers, the set of all real numbers, the set of all natural numbers and the set of all nonnegative integer numbers, respectively.

Let (X, d) be a b-metric space and $P(X)$ a collection of all subsets of X . Denote $Cl(X)$, $CB(X)$, and $K(X)$ by the collection of closed, closed and bounded and compact subsets of X , respectively.

Let $U, V \in P(X)$. The gap functional D , the excess generalized function ρ , the Pompeiu-Hausdorff generalized functional H , and the functional δ induced by a b-metric d on X are defined as:

$$\begin{aligned} (1) \quad D(U, V) &= \begin{cases} \inf_{u \in U, v \in V} d(u, v), & \text{if } U \neq V \neq \emptyset \neq U, \\ 0, & \text{if } U = V = \emptyset, \\ \infty, & \text{otherwise.} \end{cases} \\ (2) \quad \rho(U, V) &= \begin{cases} \sup_{u \in U} D(u, V), & \text{if } U \neq V \neq \emptyset \neq U, \\ 0, & \text{if } U = \emptyset, \\ \infty, & \text{if } V = \emptyset, U \neq \emptyset. \end{cases} \\ (3) \quad H(U, V) &= \begin{cases} \max\{\rho(U, V), \rho(V, U)\}, & \text{if } U \neq V \neq \emptyset \neq U, \\ 0, & \text{if } U = V = \emptyset, \\ \infty, & \text{otherwise.} \end{cases} \\ \delta(U, V) &= \begin{cases} \sup_{u \in U, v \in V} d(u, v), & \text{if } U \neq V \neq \emptyset \neq U, \\ 0, & \text{if } U = V = \emptyset, \\ \infty, & \text{otherwise.} \end{cases} \end{aligned}$$

An et al (2015b) studied the topological properties of b-metric spaces and stated that a b-metric is not necessarily continuous in each variable. If a b-metric is continuous in one variable, then it is continuous in other variable. A ball $B(u_0, \epsilon) = \{v \in X : d(u_0, v) < \epsilon\}$ in a b-metric space (X, d) is not necessarily an open set. A ball is an open set if d is continuous in one variable.

Let (X, d) be a b-metric space. We call (f, T) a hybrid pair of mappings if $f: X \rightarrow X$ and $T: X \rightarrow CB(X)$.

A mapping f is called a contraction if there is some real constant $r \in [0, 1)$ such that for any $u, v \in X$, we have $d(fu, fv) \leq rd(u, v)$.

A point u in X is a fixed point of f , if $u = fu$, a fixed point of T , if $u \in Tu$, a coincidence point of (f, T) if $fu \in T$, and a common fixed point of (f, T) if $u = fu \in Tu$. Denote $F(f), F(T)$ by the fixed points of f and T , respectively, and $C(f, T)$ and $F(f, T)$ by coincidence and common fixed point of (f, T) , respectively.

Definition 1.2, compare with (Abbas et al, 2012). A pair (f, T) is w -compatible if $f(Tu) \subseteq T(fu)$ for all $u \in C(f, T)$. The mapping f is T -weakly commuting at some point $u \in X$ if $f^2(u) \in T(fu)$.

Using an axiom of choice, Haghi et al (2011) proved the following lemma.

Lemma 1.3 (Haghi et al, 2011) Let $f: X \rightarrow X$ be a self-mapping of a nonempty set X , then there exists a subset $E \subseteq X$ such that $f(E) = f(X)$ and f is one-to-one on E .

Lemma 1.4, compare (Rus et al, 2003). Let (X, d) be a b -metric space, $U, V \in P(X)$. If there exists a $\lambda > 0$ such that for each $u \in U$, there exists a $v \in V$ such that $d(u, v) \leq \lambda$, and for each $v \in V$, there exists a $u \in U$ such that $d(u, v) \leq \lambda$, then $H(U, V) \leq \lambda$.

We need following lemmas given in (Czerwik, 1993), (Singh & Prasad, 2008).

Lemma 1.5 Let (X, d) be a b -metric space, $u, v \in X$, $\{u_n\}$ a sequence in X and $U, V \in CB(X)$. The following statements hold:

b₁:- $(CB(X), H)$ is a b -metric space and $(CB(X), H)$ is complete whenever (X, d) is complete;

b₂:- $D(u, V) \leq H(U, V)$ for all $u \in U$;

b₃:- $D(u, U) \leq bd(u, v) + bD(v, U)$;

b₄:- for $h > 1$ and $u \in U$, there is a $v \in V$ such that $d(u, v) \leq hH(U, V)$;

b₅:- for every $h > 0$ and $u \in U$, there is a $v \in V$ such that $d(u, v) \leq H(U, V) + h$;

b₆:- $D(u, U) = 0$ if and only if $u \in \overline{U} = U$;

b₇:- $d(u_0, u_n) \leq bd(u_0, u_1) + \dots + b^{n-1}d(u_{n-2}, u_{n-1}) + b^{n-1}d(u_{n-1}, u_n)$;

b₈:- $\{u_n\}$ is a Cauchy sequence if and only if for $\epsilon > 0$, there exists $n(\epsilon) \in \mathbb{N}$ such that for each $n, m \geq n(\epsilon)$ we have $d(u_n, u_m) < \epsilon$;

b₉:- $\{u_n\}$ is a convergent sequence if and only if there exists $u \in X$ such that for all $\epsilon > 0$ there exists $n(\epsilon) \in \mathbb{N}$ such that for all $n \geq n(\epsilon)$, $d(u_n, u) < \epsilon$.

A sequence $\{u_n\}$ is convergent to $u \in X$ if and only if $\lim_{n \rightarrow +\infty} d(u_n, u) = 0$.

A subset $Y \subset X$ is closed if and only if for each sequence $\{u_n\}$ in Y that converges to an element u , $u \in Y$. A subset $Y \subset X$ is bounded if $\text{diam}(Y)$ is finite, where $\text{diam}(Y) = \sup\{d(u, v) : u, v \in Y\}$. A b-metric space (X, d) is said to be complete if every Cauchy sequence in X is convergent in X .

Following are some known classes of mappings given in (Berinde, 1993), (Berinde, 1996), (Berinde, 1997), (Bota et al, 2015), (Rus, 2001).

Let $\psi, \varphi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$, then

$$\mathbf{c}_1\text{-} \Psi_1 = \left\{ \psi: \psi \text{ is increasing, } \lim_{n \rightarrow +\infty} \psi^n(t) = 0, \text{ for any } t \geq 0 \right\}.$$

The elements in this class are called comparison functions. If $\psi \in \Psi_1$, then the n^{th} iterate of ψ is a comparison function, ψ is continuous at $t = 0$, and $\psi(t) < t$, for any $t > 0$.

$$\mathbf{c}_2\text{-} \Psi_2 = \{ \psi: \sum_{n=1}^{\infty} \psi^n(t) < +\infty \text{ for all } t > 0 \text{ and } \psi \text{ is nondecreasing} \}.$$

$$\mathbf{c}_3\text{-} \Psi_3 = \left\{ \begin{array}{l} \psi: \psi \text{ is increasing, there exists a } n_0 \in \mathbb{N}, a \in (0, 1), \\ \text{a sequence } u_n \geq 0 \text{ such that } \sum_{n=1}^{\infty} u_n < +\infty \text{ and} \\ \psi^{n+1}(t) \leq a\psi^n(t) + u_n \text{ for all } n \geq n_0, t \geq 0 \end{array} \right\}$$

is called the class of (c) –comparison functions.

$$\mathbf{c}_4\text{-} \Psi_4 = \left\{ \begin{array}{l} \psi: \psi \text{ is increasing, there exists a } n_0 \in \mathbb{N}, a \in (0, 1), \\ b \geq 1, \text{ a sequence } u_n \geq 0 \text{ such that } \sum_{n=1}^{\infty} u_n < +\infty \text{ and} \\ b^{n+1}\psi^{n+1}(t) \leq ab^n\psi^n(t) + u_n \text{ for all } n \geq n_0, t \geq 0 \end{array} \right\}$$

is known as the class of (b) –comparison functions.

Note that $\Psi_2 \subseteq \Psi_1$. If $b = 1$, then $\Psi_3 = \Psi_4$.

Lemma 1.6 (Berinde, 1993) *If $\psi \in \Psi_4$ with $b \geq 1$, then the series $\sum_{n=0}^{\infty} b^n \psi^n(t)$ converges for all $t \in \mathbb{R}^+$, and $r_b(t) = \sum_{n=0}^{\infty} b^n \psi^n(t)$ is increasing and continuous at $t = 0$.*

In the light of the above lemma, $\Psi_4 \subseteq \Psi_1$.

Lemma 1.7 (Păcurar, 2010) *If $\psi \in \Psi_4$ with $b \geq 1$, and $\{a_n\} \subseteq \mathbb{R}^+$ is such that $\lim_{n \rightarrow +\infty} a_n = 0$, then*

$$\lim_{k \rightarrow +\infty} \sum_{n=0}^{\infty} b^{k-n} \psi^{k-n}(a_n) = 0.$$

Example 1.8 Let $\psi(t) = qt$, where $q \in [0, 1)$ and $t \in \mathbb{R}^+$. Consider $\sum_{k=0}^{\infty} u_k(t)$, where $u_k(t) = b^k \psi^k(t)$ and $b \geq 1$. If $t = 0$, then $\sum_{k=0}^{\infty} u_k(t)$

converges trivially. If $t > 0$, then by the generalized ratio test (Berinde, 1993), $\sum_{k=0}^{\infty} u_k(t)$ is convergent for any $t > 0$. Hence for some $n_0 \in \mathbb{N}$, $b^{n+1}\psi^{n+1}(t) \leq ab^n\psi^n(t) + u_n$ for all $n \geq n_0$ and $t \geq 0$. Consequently, we have $\psi \in \Psi_4$.

The Banach contraction principle (BCP) (Banach, 1922) states that a contraction mapping on a complete metric space has a unique fixed point.

Let $\alpha: X \times X \rightarrow \mathbb{R}^+$. A mapping $f: X \rightarrow X$ is called an α -admissible if for all $u, v \in X$, $\alpha(u, v) \geq 1$ implies that $\alpha(fu, fv) \geq 1$. Samet et al (2012), Theorem 1, obtained the following generalization of BCP.

Theorem 1.9 *Let (X, d) be a complete metric space and $f: X \rightarrow X$ an α -admissible mapping. Suppose that there is an element u_0 in X with $\alpha(u_0, fu_0) \geq 1$. If for any $u, v \in X$, there exists $\psi \in \Psi_2$ such that $\alpha(u, v)d(fu, fv) \leq \psi(d(u, v))$, then $F(f)$ is nonempty provided that f is continuous.*

Suzuki (2008) provided an interesting generalization of BCP that characterizes metric completeness.

For some other important generalizations of BCP, see (Berinde, 1993), (Berinde, 1996), (Berinde, 1997), (Bhaskar & Lakshmikantham, 2006), (Nieto & Rodríguez-López, 2005), (Nieto & Rodríguez-López, 2007), (Ran & Reurings, 2004) and references therein. A number of fixed point theorems have been obtained in b -metric spaces (Aleksić et al, 2019a), (Aleksić et al, 2019b), (Ali & Abbas, 2017), (An et al, 2015a), (An et al, 2015a), (Ćirić et al, 2012), (Chifu & Petruşel, 2014), (Czerwik, 1993), (Karapinar et al, 2020), (Latif, 2015), (Mitrović, 2019), (Păcurar, 2010).

The development of the metric fixed point theory of multivalued mappings was initiated by (Nadler, 1969). He introduced the concept of set-valued contraction mappings and extended the Banach contraction principle to set-valued mappings by using the Hausdorff metric as follows.

Theorem 1.10 *Let (X, d) be a complete metric space. If a multivalued mapping $T: X \rightarrow CB(X)$ satisfies $H(Tu, Tv) \leq rd(u, v)$ for all $u, v \in X$ and for some $r \in [0, 1)$, then $F(T)$ is nonempty.*

The fixed point theory of multivalued mappings provides a useful machinery to analyze the problems of pure, applied and computational

mathematics which can be reformulated in the form of an inclusion for an appropriate multivalued mapping.

For more results in this direction, we refer to (Abbas et al, 2012), (Abbas et al, 2013), (Asl et al, 2012), (Rus et al, 2003), (Mitrović et al, 2020).

Khojasteh et al (2014) proved a new type of the fixed point theorem for multivalued mappings in metric spaces as follows.

Theorem 1.11 *Let (X, d) be a complete metric space. If a multivalued mapping $T: X \rightarrow CB(X)$ satisfies*

$$H(Tu, Tv) \leq \left(\frac{D(u, Tv) + D(v, Tu)}{1 + \delta(u, Tu) + \delta(v, Tv)} \right) d(u, v) \quad (1.1)$$

for all $u, v \in X$, then $F(T)$ is nonempty.

Recently, Rhoades (2015) improved the result of Khojasteh for two multivalued mappings as follows.

Theorem 1.12 (Rhoades, 2015) *Let (X, d) be a complete metric space. If multivalued mappings $S, T: X \rightarrow CB(X)$ satisfy $H(Su, Tv) \leq n_{S,T}(u, v)m_{S,T}(u, v)$ for all $u, v \in X$, then $F(T) \cap F(S)$ is nonempty, where*

$$\begin{aligned} n_{S,T}(u, v) &= \left(\frac{\max\{d(u, v), D(u, Su) + D(v, Tv), D(u, Tv) + D(v, Su)\}}{1 + \delta(u, Su) + \delta(v, Tv)} \right) \\ m_{S,T}(u, v) &= \max \left\{ d(u, v), D(u, Su), D(v, Tv), \frac{D(u, Tv) + D(v, Su)}{2} \right\}. \end{aligned} \quad (1.2)$$

If $S = T$ in above theorem then we get the following result.

Theorem 1.13, (Rhoades, 2015). *Let (X, d) be a complete metric space. If a multivalued mapping $T: X \rightarrow CB(X)$ satisfies for all $u, v \in X$, $H(Tu, Tv) \leq n_{T,T}(u, v)m_{T,T}(u, v)$, then $F(T)$ is nonempty.*

Let $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $U, V \in P(X)$. Define $\alpha_*(U, V) = \inf_{u \in U, v \in V} \alpha(u, v)$.

A multivalued mapping $T: X \rightarrow Cl(X)$ is called α_* – admissible mapping if for any $u, v \in X$, $\alpha(u, v) \geq 1$ implies that $\alpha_*(Tu, Tv) \geq 1$. The concepts of α_* – admissible mapping coincides with α – admissible mapping in case of a single valued mapping.

Asl et al (2012), Theorem 1, defined $(\alpha_* - \psi)$ – contractive multifunctions and proved the following result.

Theorem 1.14 Let (X, d) be a complete metric space and $T: X \rightarrow Cl(X)$ an α_* – admissible mapping that satisfies $\alpha_*(Tu, Tv)H(Tu, Tv) \leq \psi(d(u, v))$ for all $u, v \in X$ and $\psi \in \Psi_2$. Moreover, if there exists a $u_0 \in X$ and $u_1 \in Tu_0$ such that $\alpha(u_0, u_1) \geq 1$, then $F(T)$ is nonempty provided that if $\{u_n\}$ is a sequence in X such that $\alpha(u_n, u_{n+1}) \geq 1$ for all n and $\lim_{n \rightarrow +\infty} u_n = u$, then $\alpha(u_n, u) \geq 1$ for all n .

In what follows, we assume that a b-metric d is continuous in one variable.

Definition 1.15, compare with (Rus et al, 2003). Let (X, d) be a b-metric space. A mapping $T: X \rightarrow Cl(X)$ is called a multivalued weakly Picard operator (MWP operator), if for all $u \in X$ and for some $v \in Tu$, there exists a sequence $\{u_n\}$ satisfying (a₁) $u_0 = u, u_1 = v$, (a₂) $u_{n+1} \in Tu_n$ for all $n \geq 0$, (a₃) $\{u_n\}$ converges to some $z \in F(T)$. The sequence $\{u_n\}$ satisfying (a₁) and (a₂) is called a sequence of successive approximations (ssa) of T starting from (u, v) . If T is a single-valued mapping, then we call it a Picard operator if it satisfies (a₁) to (a₃).

Recently Bota et al (2015) proved the fixed point theorem for $(\alpha_* - \psi)$ – contractive multivalued mappings as follows.

Theorem 1.16 Let (X, d) be a complete b-metric space, $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $T: X \rightarrow Cl(X)$ an α_* – admissible multivalued operator that satisfies

$$\alpha_*(Tu, Tv)H(Tu, Tv) \leq \psi(d(u, v))$$

for all $u, v \in X$ and $\psi \in \Psi_4$. Assume that there exists a $u_0 \in X$ and $u_1 \in Tu_0$ such that $\alpha(u_0, u_1) \geq 1$. Then T is a MWP operator provided that if there is a sequence $\{u_n\}$ in X such that $u_n \rightarrow u$, then $\alpha(u_n, u) \geq 1$ for all $n \in \mathbb{N}_1$.

A mapping $T: X \rightarrow Cl(X)$ is called $(\alpha_* - f)$ –admissible mapping if $u, v \in X, \alpha(fu, fv) \geq 1$ implies that $\alpha_*(Tu, Tv) \geq 1$.

Let (X, d) be a b-metric space, $g: X \rightarrow X$ and $T: X \rightarrow Cl(X)$. Set

$$N_T(u, v) = \frac{\max\{d(u, v), D(u, Tu) + D(v, Tv), D(u, Tv) + D(v, Tu)\}}{b(\delta(u, Tu) + \delta(v, Tv) + 1)},$$

$$M_T(u, v) = \max\left\{d(u, v), D(u, Tu), D(v, Tv), \frac{D(u, Tv) + D(v, Tu)}{2b}\right\},$$

and

$$N_{g,T}(u, v) = \frac{\max\{d(gu, gv), D(gu, Tu) + D(gv, Tv), D(gu, Tv) + D(gv, Tu)\}}{b(\delta(gu, Tu) + \delta(gv, Tv) + 1)}$$

$$M_{g,T}(u, v) = \max\left\{d(gu, gv), D(gu, Tu), D(gv, Tv), \frac{D(gu, Tv) + D(gv, Tu)}{2b}\right\}.$$

We now give the following definitions.

Definition 1.17 Let (X, d) be a b -metric space. A mapping $T: X \rightarrow Cl(X)$ is called Suzuki type $(\alpha_* - \psi)$ – multivalued operator if there exists a $\psi \in \Psi_4$ such that

$$\frac{1}{2}D(u, Tu) \leq bd(u, v) \quad (1.3)$$

implies

$$\alpha_*(Tu, Tv)H(Tu, Tv) \leq \max\{1, N_T(u, v)\}\psi(M_T(u, v)) \quad (1.4)$$

for all $u, v \in X$.

If in the above definition we replace a mapping T by a single valued mapping $f: X \rightarrow X$, then we call it a Suzuki type $(\alpha_* - \psi)$ – operator.

Definition 1.18 Let (X, d) be a b -metric space. A hybrid pair (g, T) is called a Suzuki type $(\alpha_* - \psi)$ – hybrid pair of operators if there exists a $\psi \in \Psi_4$ such that

$$\frac{1}{2}D(gu, Tu) \leq bd(gu, gv) \quad (1.5)$$

implies

$$\alpha_*(Tu, Tv)H(Tu, Tv) \leq \max\{1, N_{g,T}(u, v)\}\psi(M_{g,T}(u, v)) \quad (1.6)$$

for all $u, v \in X$.

In case T is replaced by a single valued mapping $f: X \rightarrow X$, we call it a Suzuki type $(\alpha_* - \psi)$ – pair of operators.

Fixed point of Suzuki type $(\alpha_* - \psi)$ – multivalued operators on b -metric spaces

In this section, we prove that Suzuki type $(\alpha_* - \psi)$ – multivalued operators are MWP operators.

Theorem 2.1 *Let (X, d) be a complete b -metric space, $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $T: X \rightarrow Cl(X)$ a Suzuki type $(\alpha_* - \psi)$ – multivalued operator. Further, assume that T is α_* – admissible mapping and there exists $x_0 \in X$ and $x_1 \in Tx_0$ such that $\alpha(x_0, x_1) \geq 1$. If for any sequence $\{x_n\}$ converging to x in X , we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$, then*

d₁-: *T is an MWP operator.*

d₂-: *If there is some $u \in F(T)$ such that $u \neq z$ and $\alpha(z, u) \geq 1$, then $d(z, u) > \frac{1}{2}$ provided that $1 \leq N_T(x, y)$ for all $x, y \in X$.*

Proof. (d₁) By the given assumption, there exists $x_0 \in X$ and $x_1 \in Tx_0$ such that $\alpha(x_0, x_1) \geq 1$. If $x_0 = x_1$, then $x_0 \in Tx_0$. Define a sequence $\{x_n\}$ in X by $x_n = x_1 = x_0$ for all $n \in \mathbb{Z}^+$. Thus $x_n \in Tx_n$ for all $n \geq 0$ and $\{x_n\}$ converges to $x = x_0 \in F(T)$ and hence T is an MWP operator. Let $x_0 \neq x_1$. Since T is α_* – admissible mapping, $\alpha_*(x_0, x_1) \geq 1$ implies that $\alpha_*(Tx_0, Tx_1) \geq 1$. As T is a Suzuki type $(\alpha_* - \psi)$ – multivalued operator and

$$\frac{1}{2}D(x_0, Tx_0) \leq d(x_0, x_1) \leq bd(x_0, x_1), \quad (2.1)$$

so from (1.4), we obtain

$$\begin{aligned} &< D(x_1, Tx_1) \leq H(Tx_0, Tx_1) < \alpha_*(Tx_0, Tx_1)H(Tx_0, Tx_1) \\ &\leq \max\{1, N_T(x_0, x_1)\}\psi(M_T(x_0, x_1)) \\ &= \max\left\{1, \left(\frac{\max\{d(x_0, x_1), D(x_0, Tx_0) + D(x_1, Tx_1), D(x_0, Tx_1) + D(x_1, Tx_0)\}}{b(1 + \delta(x_0, Tx_0) + \delta(x_1, Tx_1))}\right)\right\} \\ &\quad \psi\left(\max\left\{d(x_0, x_1), D(x_0, Tx_0), D(x_1, Tx_1), \frac{D(x_0, Tx_1) + D(x_1, Tx_0)}{2b}\right\}\right) \end{aligned}$$

$$\begin{aligned}
 &\leq \max \left\{ 1, \left(\frac{\max\{d(x_0, x_1), d(x_0, x_1) + D(x_1, Tx_1), bd(x_0, x_1) + bD(x_1, Tx_1)\}}{b(1 + d(x_0, x_1) + D(x_1, Tx_1))} \right) \right\} \\
 &\leq \psi \left(\max \left\{ d(x_0, x_1), D(x_1, Tx_1), \frac{d(x_0, x_1) + D(x_1, Tx_1)}{2} \right\} \right) \\
 &\leq \max \left\{ 1, \left(\frac{d(x_0, x_1) + D(x_1, Tx_1)}{1 + d(x_0, x_1) + D(x_1, Tx_1)} \right) \right\} \\
 &\leq \psi \left(\max \left\{ d(x_0, x_1), D(x_1, Tx_1), \frac{d(x_0, x_1) + D(x_1, Tx_1)}{2} \right\} \right) \\
 &\leq \psi(\max\{d(x_0, x_1), D(x_1, Tx_1)\}).
 \end{aligned}$$

That is

$$0 < D(x_1, Tx_1) \leq \psi(\max\{d(x_0, x_1), D(x_1, Tx_1)\}). \quad (2.2)$$

If $\max\{d(x_0, x_1), D(x_1, Tx_1)\} = D(x_1, Tx_1)$, then (2.2) implies that

$$0 < D(x_1, Tx_1) \leq \psi(D(x_1, Tx_1)). \quad (2.3)$$

As $D(x_1, Tx_1) > 0$ and $\psi \in \Psi_4$, (2.3) give

$$0 < D(x_1, Tx_1) \leq \psi(D(x_1, Tx_1)) < D(x_1, Tx_1),$$

a contradiction. Hence $\max\{d(x_0, x_1), D(x_1, Tx_1)\} = d(x_0, x_1)$. From (2.3), it follows that

$$0 < D(x_1, Tx_1) \leq \psi(d(x_0, x_1)). \quad (2.4)$$

Let $q > 1$. We may choose $x_2 \in Tx_1$ such that

$$0 < D(x_1, Tx_1) \leq d(x_1, x_2) < qD(x_1, Tx_1) \leq q\psi(d(x_0, x_1)).$$

That is

$$0 < d(x_1, x_2) < qD(x_1, Tx_1) \leq q\psi(d(x_0, x_1)). \quad (2.5)$$

Since $\alpha(x_1, x_2) \geq \alpha(Tx_0, Tx_1) \geq 1$, we get $\alpha(Tx_1, Tx_2) \geq 1$. Set $c_0 = d(x_0, x_1) > 0$, then from (2.5) we get $x_1 \neq x_2$ and $d(x_1, x_2) < q\psi(c_0)$. As $\psi \in \Psi_4$, so

$$\psi(d(x_1, x_2)) < \psi(q\psi(c_0)). \quad (2.6)$$

If $q_1 = \frac{\psi(q\psi(c_0))}{\psi(d(x_1, x_2))}$, then by (2.6) $q_1 > 1$. Now, if $x_2 \in Tx_2$ then the proof is finished. Let $x_2 \notin Tx_2$. Note that

$$\frac{1}{2}D(x_1, Tx_1) \leq d(x_1, x_2) \leq bd(x_1, x_2).$$

By (1.4)

$$\begin{aligned} &< D(x_2, Tx_2) \leq H(Tx_1, Tx_2) < \alpha_*(Tx_1, Tx_2)H(Tx_1, Tx_2) \\ &\leq \max\{1, N_T(x_1, x_2)\}\psi(M_T(x_1, x_2)) \\ &\quad \max\left\{1, \left(\frac{\max\{d(x_1, x_2), D(x_1, Tx_1) + D(x_2, Tx_2), D(x_1, Tx_2) + D(x_2, Tx_1)\}}{b(1 + \delta(x_1, Tx_1) + \delta(x_2, Tx_2))}\right)\right\} \\ &= \psi\left(\max\left\{d(x_1, x_2), D(x_1, Tx_1), D(x_2, Tx_2), \frac{D(x_1, Tx_2) + D(x_2, Tx_1)}{2b}\right\}\right) \\ &\leq \max\left\{1, \left(\frac{\max\{d(x_1, x_2), d(x_1, x_2) + D(x_2, Tx_2), b(d(x_1, x_2) + D(x_2, Tx_2))\}}{b(1 + d(x_1, x_2) + D(x_2, Tx_2))}\right)\right\} \\ &\leq \psi\left(\max\left\{d(x_1, x_2), d(x_1, x_2), D(x_2, Tx_2), \frac{d(x_1, x_2) + D(x_2, Tx_2)}{2}\right\}\right) \\ &\leq \max\left\{1, \left(\frac{d(x_1, x_2) + D(x_2, Tx_2)}{1 + d(x_1, x_2) + D(x_2, Tx_2)}\right)\right\} \\ &\leq \psi\left(\max\left\{d(x_1, x_2), d(x_1, x_2), D(x_2, Tx_2), \frac{d(x_1, x_2) + D(x_2, Tx_2)}{2}\right\}\right) \\ &\leq \psi(\max\{d(x_1, x_2), D(x_2, Tx_2)\}). \end{aligned}$$

That is

$$0 < D(x_2, Tx_2) \leq \psi(\max\{d(x_1, x_2), D(x_2, Tx_2)\}). \quad (2.7)$$

If $\max\{d(x_1, x_2), D(x_2, Tx_2)\} = D(x_2, Tx_2)$, then

$$0 < D(x_2, Tx_2) \leq \psi(D(x_2, Tx_2)). \quad (2.8)$$

Now $D(x_2, Tx_2) > 0$, $\psi \in \Psi_4$ and (2.8) give

$$0 < D(x_2, Tx_2) \leq \psi(D(x_2, Tx_2)) < D(x_2, Tx_2),$$

a contradiction. Hence

$$0 < D(x_2, Tx_2) \leq \psi(d(x_1, x_2)). \quad (2.9)$$

We may choose $x_3 \in Tx_2$ such that

$$0 < D(x_2, Tx_2) \leq d(x_2, x_3) < q_1 D(x_2, Tx_2) \leq q_1 \psi(d(x_1, x_2)) = \psi(q\psi(c_0)).$$

That is

$$0 < d(x_2, x_3) < q_1 D(x_2, Tx_2) \leq q_1 \psi(d(x_1, x_2)) = \psi(q\psi(c_0)). \quad (2.10)$$

As $\alpha(x_2, x_3) \geq \alpha(Tx_1, Tx_2) \geq 1$, so $\alpha(Tx_2, Tx_3) \geq 1$. From (2.10), we get $x_2 \neq x_3$ and

$$\psi(d(x_2, x_3)) < \psi^2(q\psi(c_0)). \quad (2.11)$$

Set $q_2 = \frac{\psi^2(q\psi(c_0))}{\psi(d(x_2, x_3))} > 1$. If $x_3 \in Tx_3$ then we are done. Suppose that $x_3 \notin Tx_3$. Similarly, we obtain $x_4 \in Tx_3$ such that

$$0 < d(x_3, x_4) < q_1 D(x_3, Tx_3) \leq q_2 \psi(d(x_2, x_3)) = \psi^2(q\psi(c_0)). \quad (2.12)$$

Continuing this way, we can obtain a sequence $\{x_n\}$ in X such that $x_{n+1} \in Tx_n, x_{n+1} \neq x_n, \alpha(x_{n+1}, x_{n+2}) \geq 1$, and it satisfies:

$$0 < D(x_{n+1}, Tx_{n+1}) \leq \psi(d(x_n, x_{n+1})) \quad (2.13)$$

and

$$0 < d(x_{n+1}, x_{n+2}) < \psi^n(q\psi(c_0)) \quad (2.14)$$

for all $n \in \mathbb{Z}^+$. From (2.14), for $n, m \in \mathbb{N}$ with $m > n$, we have

$$\begin{aligned} d(x_n, x_m) &\leq bd(x_n, x_{n+1}) + b^2 d(x_{n+1}, x_{n+2}) + \dots \\ &\quad + b^{m-n} d(x_{m-2}, x_{m-1}) + b^{m-n} d(x_{m-1}, x_m) \\ &\leq b\psi^{n-1}(q\psi(c_0)) + b^2 \psi^n(q\psi(c_0)) + \dots \\ &\quad + b^{m-n} \psi^{m-3}(q\psi(c_0)) + b^{m-n} \psi^{m-2}(q\psi(c_0)) \\ &= \frac{1}{b^{n-2}} (b^{n-1} \psi^{n-1}(q\psi(c_0)) + b^n \psi^n(q\psi(c_0)) + \dots \\ &\quad + b^{m-2} \psi^{m-2}(q\psi(c_0))) \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{b^{n-2}} \sum_{i=n-1}^{m-2} b^i \psi^i(q\psi(c_0)) \\
 &= \frac{1}{b^{n-2}} \left(\sum_{i=0}^{m-2} b^i \psi^i(q\psi(c_0)) - \sum_{i=0}^{n-2} b^i \psi^i(q\psi(c_0)) \right).
 \end{aligned}$$

That is

$$d(x_n, x_m) \leq \frac{1}{b^{n-2}} \left(\sum_{i=0}^{m-2} b^i \psi^i(q\psi(c_0)) - \sum_{i=0}^{n-2} b^i \psi^i(q\psi(c_0)) \right). \quad (2.15)$$

Set $S_n = \sum_{i=0}^n b^i \psi^i(q\psi(c_0))$. Then from (2.15) we obtain that

$$d(x_n, x_m) \leq \frac{1}{b^{n-2}} (S_{m-2} - S_{n-2}). \quad (2.16)$$

By Lemma 1.6, $\sum_{i=0}^{\infty} b^i \psi^i(t)$ converges for any $t \geq 0$. Hence $\lim_{n \rightarrow +\infty} S_{n-2} = S$ for some $S \in \mathbb{R}^+$. If $b = 1$, then from (2.16) we get $\lim_{n \rightarrow +\infty} d(x_n, x_m) \leq \lim_{n \rightarrow +\infty} S_{m-1} - \lim_{n \rightarrow +\infty} S_{n-1} = 0$. If $b > 1$, then from (2.16) we have

$$\lim_{n \rightarrow +\infty} d(x_n, x_m) \leq \lim_{n \rightarrow +\infty} \frac{1}{b^{n-1}} (S_{m-1} - S_{n-1}) \leq \lim_{n \rightarrow +\infty} \frac{S_{m-1}}{b^{n-1}} = 0$$

for all $m, n \in \mathbb{N}$. Hence $\{x_n\}$ is a Cauchy sequence in X . There exists $z \in X$ such that

$$\lim_{n \rightarrow +\infty} d(x_n, z) = 0. \quad (2.17)$$

Now we show that $z \in F(T)$. If $D(z, Tz) > 0$, then we claim that one of the following two inequalities

$$\frac{1}{2} D(x_n, Tx_n) \leq bd(x_n, z) \quad (2.18)$$

$$\frac{1}{2}D(x_{n+1}, Tx_{n+1}) \leq bd(x_{n+1}, z) \quad (2.19)$$

holds for all $n \in \mathbb{Z}^+$. Assume on the contrary that there exists an $n_0 \in \mathbb{Z}^+$ such that

$$\frac{1}{2}D(x_{n_0}, Tx_{n_0}) > bd(x_{n_0}, z) \quad (2.20)$$

and

$$\frac{1}{2}D(x_{n_0+1}, Tx_{n_0+1}) > bd(x_{n_0+1}, z). \quad (2.21)$$

Now from (2.13), (2.20) and (2.21), we have

$$\begin{aligned} d(x_{n_0}, x_{n_0+1}) &\leq bd(x_{n_0}, z) + bd(z, x_{n_0+1}) \\ &< \frac{1}{2}D(x_{n_0}, Tx_{n_0}) + \frac{1}{2}D(x_{n_0+1}, Tx_{n_0+1}) \\ &\leq \frac{1}{2}d(x_{n_0}, x_{n_0+1}) + \frac{1}{2}\psi(d(x_{n_0}, x_{n_0+1})) \\ &< \frac{1}{2}d(x_{n_0}, x_{n_0+1}) + \frac{1}{2}d(x_{n_0}, x_{n_0+1}) \\ &= d(x_{n_0}, x_{n_0+1}) \end{aligned}$$

a contradiction. Hence either (2.18) or (2.19) holds for an infinite subset N_1 of \mathbb{Z}^+ . By the given assumption, it follows that $\alpha(x_n, z) \geq 1$. As T is α_* -admissible, $\alpha(Tx_n, Tz) \geq 1$. Now if (2.18) holds for all $n \in \mathbb{Z}^+$, then from (1.4) we get

$$\begin{aligned} D(x_{n+1}, Tz) &\leq H(Tx_n, Tz) < \alpha_*(Tx_n, Tz)H(Tx_n, Tz) \\ &\leq \max\{1, N_T(x_n, z)\}\psi(M_T(x_n, z)) \\ &= \max\left\{1, \left(\frac{\max\{d(x_n, z), D(x_n, Tx_n) + D(z, Tz), D(x_n, Tz) + D(z, Tx_n)\}}{b(1 + \delta(x_n, Tx_n) + \delta(z, Tz))}\right)\right\} \\ &= \psi\left(\max\left\{d(x_n, z), D(x_n, Tx_n), D(z, Tz), \frac{D(x_n, Tz) + D(z, Tx_n)}{2b}\right\}\right) \\ &\leq \max\left\{1, \left(\frac{\max\{d(x_n, z), d(x_n, x_{n+1}) + D(z, Tz), D(x_n, Tz) + d(z, x_{n+1})\}}{b(1 + d(x_n, x_{n+1}) + D(z, Tz))}\right)\right\} \\ &\leq \psi\left(\max\left\{d(x_n, z), d(x_n, x_{n+1}), D(z, Tz), \frac{D(x_n, Tz) + d(z, x_{n+1})}{2b}\right\}\right). \end{aligned}$$

On taking limit as $n \rightarrow +\infty$, we have

$$\begin{aligned} & \lim_{n \rightarrow +\infty} D(x_{n+1}, Tz) \\ & \leq \lim_{n \rightarrow +\infty} \max \left\{ 1, \left(\frac{\max\{d(x_n, z), d(x_n, x_{n+1}) + D(z, Tz), D(x_n, Tz) + d(z, x_{n+1})\}}{b(1 + d(x_n, x_{n+1}) + D(z, Tz))} \right) \right\} \\ & \leq \lim_{n \rightarrow +\infty} \psi \left(\max \left\{ d(x_n, z), d(x_n, x_{n+1}), D(z, Tz), \frac{D(x_n, Tz) + d(z, x_{n+1})}{2b} \right\} \right). \end{aligned}$$

As ψ is continuous at 0, $\psi \in \Psi_4$ and $D(z, Tz) > 0$, we have

$$D(z, Tz) \leq \max \left\{ 1, \frac{D(z, Tz)}{b(1 + D(z, Tz))} \right\} \psi(D(z, Tz)) = \psi(D(z, Tz)) < D(z, Tz).$$

a contradiction. Consequently, $z \in Tz$. Similarly, we obtain $z \in Tz$ when (2.19) holds for an infinite subset N_1 of \mathbb{Z}^+ .

To prove part (d₂), let $u \in F(T)$ such that $u \neq z$ and $\alpha(z, u) \geq 1$. Since T is α_* -admissible, $\alpha_*(Tz, Tu) \geq 1$. Now $\frac{1}{2}D(z, Tz) = 0 \leq d(z, u)$ implies that

$$\begin{aligned} d(z, u) & \leq bD(z, Tz) + bD(Tz, u) \leq bH(Tz, Tu) \\ & \leq b\alpha_*(Tz, Tu)H(H(Tz, Tu)) \\ & \leq b \max \left\{ 1, \left(\frac{\max\{d(z, u), D(z, Tz) + D(u, Tu), D(z, Tu) + D(u, Tz)\}}{b(1 + \delta(z, Tz) + \delta(u, Tu))} \right) \right\} \\ & \leq \psi \left(\max \left\{ d(z, u), D(z, Tz), D(u, Tu), \frac{D(z, Tu) + D(u, Tz)}{2b} \right\} \right) \\ & \leq b \max \left\{ 1, \left(\frac{\max\{d(z, u), d(z, z) + d(u, u), d(z, u) + d(u, z)\}}{b(1 + d(z, z) + d(u, u))} \right) \right\} \\ & \leq \psi \left(\max \left\{ d(z, u), d(z, z), d(u, u), \frac{d(z, u) + d(u, z)}{2b} \right\} \right) \\ & \leq b \max \left\{ 1, \left(\frac{2d(z, u)}{b} \right) \right\} \psi(d(z, u)). \end{aligned}$$

Now, $\max\{1, N_T(x, y)\} = N_T(x, y)$ gives

$$d(z, u) \leq 2d(z, u)\psi(d(z, u)) < 2d^2(z, u)$$

and hence $d(z, u) > \frac{1}{2}$.

Corollary 2.2 Let (X, d) be a complete b -metric space, $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $T: X \rightarrow Cl(X)$ an α_* – admissible mapping such that

$$\frac{1}{2}D(x, Tx) \leq bd(x, y)$$

implies that

$$\alpha_*(Tx, Ty)H(Tx, Ty) \leq \max\{1, N_T(x, y)\}\psi(d(x, y))$$

for all $x, y \in X, \psi \in \Psi_4$. Further, assume that there exists $x_0 \in X$ and $x_1 \in Tx_0$ such that $\alpha(x_0, x_1) \geq 1$. If for any sequence $\{x_n\}$ converging to x in X , we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$ then

e₁-: T is an MWP operator

e₂-: If there is some $u \in F(T)$ such that $u \neq z$ and $\alpha(z, u) \geq 1$, then $d(z, u) > \frac{1}{2}$ provided that

$$\max\{1, N_T(x, y)\} = N_T(x, y) \text{ for all } x, y \in X.$$

Corollary 2.3 Let (X, d) be a complete b -metric space, $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $T: X \rightarrow Cl(X)$ an α_* – admissible mapping such that

$$\frac{1}{2}D(x, Tx) \leq bd(x, y)$$

implies that

$$\alpha_*(Tx, Ty)H(Tx, Ty) \leq \max\left\{1, \frac{d(x, y)}{b(1 + \delta(x, Tx) + \delta(y, Ty))}\right\}\psi(d(x, y))$$

for all $x, y \in X, \psi \in \Psi_4$. Further, assume that there exists $x_0 \in X$ and $x_1 \in Tx_0$ such that $\alpha(x_0, x_1) \geq 1$. If for any sequence $\{x_n\}$ converging to x in X , we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$ then

e₃-: T is an MWP operator.

e₄-: If there is some $u \in F(T)$ such that $u \neq z$ and $\alpha(z, u) \geq 1$, then $d(z, u) > 1$ provided that

$$\max \left\{ 1, \frac{d(x, y)}{b(1 + \delta(x, Tx) + \delta(y, Ty))} \right\} = \frac{d(x, y)}{b(1 + \delta(x, Tx) + \delta(y, Ty))}$$

for all $x, y \in X$.

Proof. Follows from Corollary 2.3.

Corollary 2.4 Let (X, d) be a complete b-metric space, $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $T: X \rightarrow Cl(X)$ an α_* – admissible mapping such that

$$\frac{1}{2} D(x, Tx) \leq bd(x, y)$$

implies that

$$\alpha_*(Tx, Ty)H(Tx, Ty) \leq N_T(x, y)\psi(M_T(x, y))$$

for all $x, y \in X, \psi \in \Psi_4$. Further, assume that there exists $x_0 \in X$ and $x_1 \in Tx_0$ such that $\alpha(x_0, x_1) \geq 1$. If for any sequence $\{x_n\}$ converging to x in X , we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$ then

e₅-: T is an MWP operator.

e₆-: If there is some $u \in F(T)$ such that $u \neq z$ and $\alpha(z, u) \geq 1$, then $d(z, u) > \frac{1}{2}$.

Proof. Take $\max\{1, N_T(x, y)\} = N_T(x, y)$ in Theorem 2.1.

Corollary 2.5 Let (X, d) be a complete b-metric space, $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $T: X \rightarrow Cl(X)$ an α_* –admissible mapping such that

$$\begin{aligned} \frac{1}{2} D(x, Tx) \leq bd(x, y) \text{ implies that } \alpha_*(Tx, Ty)H(Tx, Ty) \\ \leq N_T(x, y)\psi(d(x, y)) \end{aligned} \quad (2.22)$$

for all $x, y \in X, \psi \in \Psi_4$. Further, assume that there exists $x_0 \in X$ and $x_1 \in Tx_0$ such that $\alpha(x_0, x_1) \geq 1$. If for any sequence $\{x_n\}$ converging to x in X , we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$ then

e₇-: T is an MWP operator.

e₈-: If there is some $u \in F(T)$ such that $u \neq z$ and $\alpha(z, u) \geq 1$, then $d(z, u) > \frac{1}{2}$.

Proof. Take $M_T(x, y) = d(x, y)$ in Corollary 2.4.

The following Corollary is a Suzuki type generalization of (Asl et al, 2012), Theorem 2.1 (Bota et al, 2015), Theorem 1 (Mohammadi, 2013), Theorem 3.1 (Samet et al, 2012), Theorem 2.2 and references therein in the context of b-metric spaces.

Corollary 2.6 Let (X, d) be a complete b-metric space, $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $T: X \rightarrow Cl(X)$ an α_* – admissible mapping such that

$$\frac{1}{2}D(x, Tx) \leq bd(x, y) \text{ implies that} \quad (2.23)$$

$$\alpha_*(Tx, Ty)H(Tx, Ty) \leq \psi(M_T(x, y))$$

for all $x, y \in X, \psi \in \Psi_4$. Further, assume that there exists $x_0 \in X$ and $x_1 \in Tx_0$ such that $\alpha(x_0, x_1) \geq 1$. Then T is an MWP operator provided that for any sequence $\{x_n\}$ converging to x in X , we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$.

Proof. Take $\max\{1, N_T(x, y)\} = 1$ in Theorem 2.1.

Corollary 2.7 Let (X, d) be a complete b-metric space, $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $T: X \rightarrow Cl(X)$ an α_* – admissible mapping such that

$$\alpha_*(Tx, Ty)H(Tx, Ty) \leq \psi(M_T(x, y)) \quad (2.24)$$

for all $x, y \in X, \psi \in \Psi_4$. Further, assume that there exists $x_0 \in X$ and $x_1 \in Tx_0$ such that $\alpha(x_0, x_1) \geq 1$. Then T is an MWP operator provided that for any sequence $\{x_n\}$ converging to x in X , we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$.

Corollary 2.8 Let (X, d) be a complete b-metric space, $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $T: X \rightarrow Cl(X)$ an α_* – admissible mapping such that

$$\frac{1}{2}D(x, Tx) \leq bd(x, y)$$

implies that

$$\alpha_*(Tx, Ty)H(Tx, Ty) \leq \psi(d(x, y)) \quad (2.25)$$

for all $x, y \in X, \psi \in \Psi_4$. Further, assume that there exists $x_0 \in X$ and $x_1 \in Tx_0$ such that $\alpha(x_0, x_1) \geq 1$. Then T is an MWP operator provided that for any sequence $\{x_n\}$ converging to x in X , we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$.

Proof. Take $M_T(x, y) = d(x, y)$ in Corollary 2.6.

Now we state Theorem 2.1 in the context of single valued mapping $f: X \rightarrow X$, where X is a b -metric space. The existence of a fixed point follows immediately from Theorem 2.1. To prove the uniqueness of the fixed point, we need the condition H, given as follows:

(H): for all $x, y \in X$, there exists a $z \in X$ such that $\alpha(x, z) \geq 1$ and $\alpha(y, z) \geq 1$.

Corollary 2.9 Let (X, d) be a complete b -metric space and $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $f: X \rightarrow X$ an α – admissible mapping such that

$$\frac{1}{2}d(x, fx) \leq bd(x, y)$$

implies that

$$\alpha(fx, fy)d(fx, fy) \leq \max\{1, N_f(x, y)\}\psi(M_f(x, y)) \quad (2.26)$$

for all $x, y \in X, \psi \in \Psi_4$. Further, assume that there exists $x_0 \in X$ and $x_1 = fx_0$ such that $\alpha(x_0, x_1) \geq 1$ and for any sequence $\{x_n\}$ converging to x in X , we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$. If the condition (H) is satisfied, then f is a Picard operator and for an arbitrary $z \in X$, the sequence $\{f^n z\}$ converges to some $w \in F(f)$ and

e₉-: $F(f) = \{w\}$ if $\max\{1, N_f(x, y)\} = 1$,

e₁₀-: $d(u, w) > \frac{b}{2}$ for any $u \in F(f)$ such that $u \neq w$ provided that $\max\{1, N_f(x, y)\} = N_f(x, y)$.

Proof. Theorem 2.1, f is a Picard operator and $F(f)$ is nonempty. Let $u, v \in F(f)$ such that $u \neq v$. By the condition (H), there exists a $z \in X$ such that $\alpha(u, z) \geq 1$ and $\alpha(v, z) \geq 1$. Note that $\{f^n z\}$ is a Picard sequence which converges to some $w \in F(f)$. As f is an α – admissible mapping, so for all $n \geq 1$ we have $\alpha(u, f^n z) \geq 1$ and $\alpha(v, f^n z) \geq 1$. Since

$$\frac{1}{2}d(u, fu) = 0 \leq bd(u, f^{n-1}z),$$

by (2.26) we have

$$\begin{aligned} d(u, f^n z) &= d(fu, f^n z) \\ &\leq \alpha(fu, f f^{n-1} z) d(fu, f f^{n-1} z) \\ &\leq \max\{1, N_f(u, f^{n-1} z)\} \psi(M_f(u, f f^{n-1} z)) \\ &\leq \max\left\{1, \left(\frac{\max\{d(u, f^{n-1} z), d(u, fu) + d(f^{n-1} z, f f^{n-1} z), d(u, f f^{n-1} z) + d(f^{n-1} z, fu)\}}{b(1 + d(u, fu) + d(f^{n-1} z, f f^{n-1} z))}\right)\right\} \\ &\quad \psi\left(\max\left\{d(u, f^{n-1} z), d(u, fu), d(f^{n-1} z, f f^{n-1} z), \frac{d(u, f f^{n-1} z) + d(f^{n-1} z, fu)}{2b}\right\}\right) \\ &\leq \max\left\{1, \left(\frac{\max\{d(u, f^{n-1} z), d(u, u) + d(f^{n-1} z, f^n z), d(u, f^n z) + d(f^{n-1} z, u)\}}{b(1 + d(u, u) + d(f^{n-1} z, f^n z))}\right)\right\} \\ &\quad \psi\left(\max\left\{d(u, f^{n-1} z), d(u, u), d(f^{n-1} z, f^n z), \frac{d(u, f^n z) + d(f^{n-1} z, u)}{2b}\right\}\right). \end{aligned}$$

On taking limit as $n \rightarrow +\infty$, we obtain that

$$\begin{aligned} d(u, w) &\leq \max\left\{1, \left(\frac{\max\{d(u, w), d(u, u) + d(w, w), d(u, w) + d(w, u)\}}{b(1 + d(u, u) + d(w, w))}\right)\right\} \\ &\quad \psi\left(\max\left\{d(u, w), d(u, u), d(w, w), \frac{d(u, w) + d(w, u)}{2b}\right\}\right) \\ &= \max\left\{1, \left(\frac{2d(u, w)}{b}\right)\right\} \psi(d(u, w)). \end{aligned}$$

That is

$$d(u, w) \leq \max\left\{1, \left(\frac{2d(u, w)}{b}\right)\right\} \psi(d(u, w)). \quad (2.27)$$

Now, If $\max\{1, N_f(x, y)\} = 1$, and $w \neq u$, then we have

$$d(u, w) \leq \psi(d(u, w)) < d(u, w). \quad (2.28)$$

Also, if $\max\{1, N_f(x, y)\} = 1$, and $v \neq w$, then we have

$$d(v, w) \leq \psi(d(v, w)). \quad (2.29)$$

A contradiction in both cases. Thus $w = u = v$, and hence $F(f)$ is singleton. If $\max\{1, N_f(x, y)\} = N_f(x, y)$ and $u \neq w$ then from (2.27) we get

$$d(u, w) \leq \frac{2d(u, w)}{b} \psi(d(u, w)) < \frac{2}{b} d^2(u, w)$$

and $d(u, w) > \frac{b}{2}$.

Corollary 2.10 Let (X, d) be a complete b-metric space and $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $f: X \rightarrow X$ an α – admissible mapping such that

$$\frac{1}{2}d(x, fx) \leq bd(x, y) \text{ implies that } \alpha(fx, fy)d(fx, fy) \leq \psi(M_f(x, y))$$

for all $x, y \in X, \psi \in \Psi_4$. Further, assume that there exists $x_0 \in X$ and $x_1 = fx_0$ such that $\alpha(x_0, x_1) \geq 1$ and for any sequence $\{x_n\}$ converging to x in X , we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$. If the condition (H) is satisfied, then f is a Picard operator and for an arbitrary $z \in X$, the sequence $\{f^n z\}$ converges to $w \in F(f)$ and $F(f) = \{w\}$.

Corollary 2.11 Let (X, d) be a complete b-metric space and $\alpha: X \times X \rightarrow \mathbb{R}^+$. Let $f: X \rightarrow X$ be an α – admissible mapping that satisfies

$$\frac{1}{2}d(x, fx) \leq bd(x, y)$$

implies that

$$\alpha(fx, fy)d(fx, fy) \leq \max\{1, N_f(x, y)\}\psi(d(x, y))$$

for all $x, y \in X, \psi \in \Psi_4$. Moreover, suppose that there exists $x_0 \in X$ and $x_1 = fx_0$ such that $\alpha(x_0, x_1) \geq 1$ and if there is a sequence $\{x_n\}$ in X such that $x_n \rightarrow x$, then $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$. Further, assume that there exists $x_0 \in X$ and $x_1 = fx_0$ such that $\alpha(x_0, x_1) \geq 1$ and for any sequence $\{x_n\}$ converging to x in X , we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$. If the condition (H) is satisfied, then f is a Picard operator and for an arbitrary $z \in X$, the sequence $\{f^n z\}$ converges to $w \in F(f)$ and $F(f) = \{w\}$. Also, $d(u, w) > \frac{b}{2}$ for any $u \in F(f)$ such that $u \neq w$ provided that $\max\{1, N_f(x, y)\} = N_f(x, y)$.

Example 2.12 Let $X = \{x_1, x_2, x_3, x_4, x_5\}$ and $d: X \times X \rightarrow \mathbb{R}^+$ be defined as

$$\begin{aligned} d(x_2, x_5) &= d(x_3, x_4) = d(x_3, x_5) = d(x_2, x_4) = 6, \\ d(x_2, x_3) &= 9, d(x_1, x_4) = d(x_1, x_5) = 10, \\ d(x_1, x_2) &= d(x_1, x_3) = 4, d(x_4, x_5) = 1, \\ d(x, x) &= 0 \text{ and } d(x, y) = d(y, x) \text{ for all } x, y \in X. \end{aligned}$$

As $9 = d(x_2, x_3) \not\leq d(x_2, x_1) + d(x_1, x_3) = 8$, so d is not a metric on X . Indeed, (X, d) is a b-metric space with $b = \frac{9}{8} > 1$. Consider a mapping $T: X \rightarrow Cl(X)$ defined by $Tx_1 = Tx_2 = Tx_3 = \{x_1\}$, $Tx_4 = \{x_2\}$ and $Tx_5 = \{x_3\}$. If we take $\psi(t) = \frac{9}{10}t$ for $t \in \mathbb{R}^+$, then $\psi \in \Psi_4$ (see 1.8). If mapping $\alpha: X \times X \rightarrow \mathbb{R}^+$ is defined as $\alpha(x_i, x_j) = 1$ for all $i, j \in \{1, 2, 3, 4, 5\}$, then T is an α_* -admissible mapping. For $x, y \in \{x_1, x_2, x_3\}$, we have $H(Tx, Ty) = 0 \leq \max\{1, N_T(x, y)\}\psi(M_T(x, y))$. For (x, y) when $x \in \{x_1, x_2, x_3\}$ and $y \in \{x_4, x_5\}$, we obtain that

$$\begin{aligned} \alpha(Tx_1, Tx_4)H(Tx_1, Tx_4) &= d(x_1, x_2) = 4 \leq 9 = \psi(d(x_1, x_4)) \\ &\leq \max\{1, N_T(x_1, x_4)\}\psi(M_T(x_1, x_4)), \\ \alpha(Tx_2, Tx_4)H(Tx_2, Tx_4) &= d(x_1, x_2) = 4 \leq \frac{54}{10} = \psi(d(x_2, x_4)) \\ &\leq \max\{1, N_T(x_2, x_4)\}\psi(M_T(x_2, x_4)), \\ \alpha(Tx_3, Tx_4)H(Tx_3, Tx_4) &= d(x_1, x_2) = 4 \leq \frac{54}{10} = \psi(d(x_3, x_4)) \\ &\leq \max\{1, N_T(x_3, x_4)\}\psi(M_T(x_3, x_4)), \\ \alpha(Tx_1, Tx_5)H(Tx_1, Tx_5) &= d(x_1, x_3) = 4 \leq 9 = \psi(d(x_1, x_5)) \\ &\leq \max\{1, N_T(x_1, x_5)\}\psi(M_T(x_1, x_5)), \\ \alpha(Tx_2, Tx_5)H(Tx_2, Tx_5) &= d(x_1, x_3) = 4 \leq \frac{54}{10} = \psi(d(x_2, x_5)) \\ &\leq \max\{1, N_T(x_2, x_5)\}\psi(M_T(x_2, x_5)), \\ \alpha(Tx_3, Tx_5)H(Tx_3, Tx_5) &= d(x_1, x_3) = 4 \leq \frac{54}{10} = \psi(d(x_3, x_5)) \\ &\leq \max\{1, N_T(x_3, x_5)\}\psi(M_T(x_3, x_5)). \end{aligned}$$

Note that

$$\frac{1}{2}D(x_4, Tx_4) = \frac{1}{2}d(x_4, x_2) = 3 > \frac{9}{8} = bd(x_4, x_5),$$

and

$$\frac{1}{2}D(x_5, Tx_5) = \frac{1}{2}d(x_5, x_3) = 3 > \frac{5}{4} = bd(x_5, x_4).$$

Hence

$$\frac{1}{2}(D(x, Tx) \leq bd(x, y)) \leq 0$$

implies

$$\alpha(Tx, Ty)H(Tx, Ty) \leq \max\{1, N_T(x, y)\}\psi(M_T(x, y))$$

holds for all $x, y \in X$. Thus all the conditions of Theorem 2.1 are satisfied. On the other hand, if we take $x = x_4, y = x_5$, then $\alpha(Tx_4, Tx_5)H(Tx_4, Tx_5) = d(x_2, x_3) = 9 > \psi(d(x_4, x_5)) = \psi(1) = \frac{9}{10}$ and $\alpha(Tx_4, Tx_5)H(Tx_4, Tx_5) \not\leq \psi(d(x_4, x_5))$. Consequently, Theorem 1.16 in (Bota et al, 2015) does not hold in this case.

The following example illustrates an assumption $\max\{1, N_T(x, y)\} > 1$.

Example 2.13 Let $X = \{x_1, x_2, x_3\}$ and $d: X \times X \rightarrow \mathbb{R}^+$ be defined as

$$\begin{aligned} d(x_1, x_2) &= 4, d(x_1, x_3) = 1, d(x_2, x_3) = 2, \\ d(x, x) &= 0 \text{ and } d(x, y) = d(y, x) \text{ for all } x, y \in X. \end{aligned}$$

As $4 = d(x_1, x_2) \not\leq d(x_1, x_3) + d(x_3, x_2) = 3$, so d is not a metric on X . Indeed, (X, d) is a b -metric space with $b = \frac{4}{3} > 1$. Consider a mapping $T: X \rightarrow Cl(X)$ defined by

$$Tx = \begin{cases} \{x_2\} & \text{if } x = x_1, \\ \{x_1\} & \text{if } x = x_2, \\ \{x_2\} & \text{if } x = x_3. \end{cases}$$

If we take $\psi(t) = \frac{8}{9}t$ for $t \in \mathbb{R}^+$, then $\psi \in \Psi_4$ (see Example 1.8). If $\alpha: X \times X \rightarrow \mathbb{R}^+$ is defined as $\alpha(x_i, x_j) = 1$ for all $i, j \in \{1, 2, 3\}$, then T is α_* -admissible. Note that

$$\begin{aligned} N_T(x_1, x_2) &= \frac{3\max\{d(x_1, x_2), D(x_1, Tx_1) + D(x_2, Tx_2), D(x_1, Tx_2) + D(x_2, Tx_1)\}}{4(1 + \delta(x_1, Tx_2) + \delta(x_2, Tx_1))} \\ &= \frac{3\max\{d(x_1, x_2), d(x_1, x_2) + d(x_2, x_1), d(x_1, x_1) + d(x_2, x_2)\}}{4(1 + d(x_1, x_1) + d(x_2, x_2))} \\ &= \frac{3\max\{4, 8, 0\}}{4(1 + 0)} = 6 > 1. \end{aligned}$$

Hence $\max\{1, N_T(x, y)\} = 6 > 1$. Note that

$$\begin{aligned} N_T(x_1, x_3) &= \frac{3\max\{d(x_1, x_3), D(x_1, Tx_1) + D(x_3, Tx_3), D(x_1, Tx_3) + D(x_3, Tx_1)\}}{4(1 + \delta(x_1, Tx_3) + \delta(x_3, Tx_1))} \\ &= \frac{3\max\{d(x_1, x_3), d(x_1, x_2) + d(x_3, x_3), d(x_1, x_3) + d(x_3, x_2)\}}{4(1 + d(x_1, x_3) + d(x_3, x_2))} \\ &= \frac{3\max\{1, 4, 3\}}{4(1 + 1 + 2)} = \frac{3}{4}, \end{aligned}$$

and

$$\begin{aligned} N_T(x_2, x_3) &= \frac{3\max\{d(x_2, x_3), D(x_2, Tx_2) + D(x_3, Tx_3), D(x_2, Tx_3) + D(x_3, Tx_2)\}}{4(1 + \delta(x_2, Tx_3) + \delta(x_3, Tx_2))} \\ &= \frac{3\max\{d(x_2, x_3), d(x_2, x_1) + d(x_3, x_3), d(x_2, x_3) + d(x_3, x_1)\}}{4(1 + d(x_2, x_3) + d(x_3, x_1))} \\ &= \frac{3\max\{2, 4, 3\}}{4(1 + 2 + 1)} = \frac{3}{4}. \end{aligned}$$

Also,

$$\begin{aligned} \alpha(Tx_1, Tx_2)H(Tx_1, Tx_2) &= d(x_2, x_1) = 4 < \frac{64}{3} = N_T(x_1, x_2)\psi(d(x_1, x_2)) \\ &\leq \max\{1, N_T(x_1, x_2)\psi(M_T(x_1, x_2))\}, \\ \alpha(Tx_1, Tx_3)H(Tx_1, Tx_3) &= d(x_2, x_3) = 2 < \frac{8}{3} = N_T(x_1, x_3)\psi(d(x_1, Tx_1)) \\ &\leq \max\{1, N_T(x_1, x_3)\psi(M_T(x_1, x_3))\}, \\ \alpha(Tx_2, Tx_3)H(Tx_2, Tx_3) &= d(x_1, x_3) = 1 < \frac{4}{3} = N_T(x_2, x_3)\psi(d(x_2, x_3)) \\ &\leq \max\{1, N_T(x_2, x_3)\psi(M_T(x_2, x_3))\}. \end{aligned}$$

Thus all the conditions of Theorem 2.1 are satisfied. On the other hand, if we take $x = x_1, y = x_2$, then $\alpha(Tx_1, Tx_2)H(Tx_1, Tx_2) = d(x_2, x_3) = 4 > \psi(d(x_1, x_2)) = \psi(4) = \frac{32}{9}$. Hence $\alpha(Tx_1, Tx_2)H(Tx_1, Tx_2) \not\leq \psi(d(x_1, x_2))$. Consequently Theorem 1.16 in (Bota et al, 2015) is not applicable in this case which is a generalization of Theorems 1.14 and 1.9.

For $b = 1$, Theorem 2.1 reduces to the following important Corollary.

Corollary 2.14 Let (X, d) be a complete metric space, $\alpha: X \times X \rightarrow \mathbb{R}^+$ and $T: X \rightarrow Cl(X)$ satisfies the following implication

$$\frac{1}{2}D(x, Tx) \leq d(x, y) \text{ implies } \max\{1, n_{T,T}(x, y)\}\psi(m_{T,T}(x, y))$$

for all $x, y \in X$ and $\psi \in \Psi_4$ where $n_{T,T}(x, y)$ and $m_{T,T}(x, y)$ are the same as given in (1.2), Further, assume that there exists $x_0 \in X$ and $x_1 \in Tx_0$ such that $\alpha(x_0, x_1) \geq 1$ and for any sequence $\{x_n\}$ converging to x in X , we have $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$. Then

e₁₁:- T is an MWP operator.

e₁₂:- If there is some $u \in F(T)$ such that $u \neq z$ and $\alpha(z, u) \geq 1$, then $d(z, u) > \frac{1}{2}$ provided that

$$\max\{1, n_{T,T}(x, y)\} = n_{T,T}(x, y).$$

Next we present an example which shows that Corollary 2.14 is a potential generalization of Theorems 1.14, 1.9, 1.11, 1.13.

Example 2.15 Let $X = \{x_1, x_2, x_3, x_4, x_5\}$ and $d: X \times X \rightarrow \mathbb{R}^+$ be defined by

$$\begin{aligned} d(x_2, x_5) &= d(x_3, x_4) = d(x_3, x_5) = d(x_2, x_4) = 5, \\ d(x_1, x_4) &= d(x_1, x_5) = 9, d(x_1, x_2) = d(x_1, x_3) = 4, \\ d(x_4, x_5) &= 2, d(x_2, x_3) = 8, \\ d(x, x) &= 0 \text{ and } d(x, y) = d(y, x) \text{ for all } x, y \in X. \end{aligned}$$

Note that d is a metric on X . Consider a mapping $T: X \rightarrow Cl(X)$ defined by $Tx_1 = Tx_2 = Tx_3 = \{x_1\}$, $Tx_4 = \{x_2\}$ and $Tx_5 = \{x_3\}$. If we take $\psi(t) = \frac{8}{9}t$ for $t \in \mathbb{R}^+$, then $\psi \in \Psi_i$ for each $i = 1, 2, 3, 4$ (see Example 1.8). If $\alpha: X \times X \rightarrow \mathbb{R}^+$ is defined as $\alpha(x_i, x_j) = 1$ for all $i, j \in \{1, 2, 3, 4, 5\}$, then T is α_* -admissible mapping. For $x, y \in \{x_1, x_2, x_3\}$, we have $H(Tx, Ty) = 0 \leq \max\{1, N_T(x, y)\}\psi(M_T(x, y))$. For (x, y) , when $x \in \{x_1, x_2, x_3\}$ and $y \in \{x_4, x_5\}$, we obtain that

$$\begin{aligned} \alpha(Tx_1, Tx_4)H(\alpha(Tx_1, Tx_4)) &= d(x_1, x_2) = 4 \leq 8 = \psi(d(x_1, x_4)) \\ &\leq \max\{1, n_{T,T}(x_1, x_4)\}\psi(m_{T,T}(x_1, x_4)), \\ \alpha(Tx_2, Tx_4)H(Tx_2, Tx_4) &= d(x_1, x_2) = 4 \leq \frac{40}{9} = \psi(d(x_2, x_4)) \\ &\leq \max\{1, n_{T,T}(x_2, x_4)\}\psi(m_{T,T}(x_2, x_4)), \\ \alpha(Tx_3, Tx_4)H(Tx_3, Tx_4) &= d(x_1, x_2) = 4 \leq \frac{40}{9} = \psi(d(x_3, x_4)) \end{aligned}$$

$$\begin{aligned}
 &\leq \max\{1, n_{T,T}(x_3, x_4)\psi(m_{T,T}(x_3, x_4)), \\
 \alpha(Tx_1, Tx_5)H(Tx_1, Tx_5) &= d(x_1, x_3) = 4 \leq 8 = \psi(d(x_1, x_5)) \\
 &\leq \max\{1, n_{T,T}(x_1, x_5)\psi(m_{T,T}(x_1, x_5)), \\
 \alpha(Tx_2, Tx_5)H(Tx_2, Tx_5) &= d(x_1, x_3) = 4 \leq \frac{40}{9} = \psi(d(x_2, x_5)) \\
 &\leq \max\{1, n_{T,T}(x_2, x_5)\psi(m_{T,T}(x_2, x_5)), \\
 \alpha(Tx_3, Tx_5)H(Tx_3, Tx_5) &= d(x_1, x_3) = 4 \leq \frac{40}{9} = \psi(d(x_3, x_5)) \\
 &\leq \max\{1, n_{T,T}(x_3, x_5)\psi(m_{T,T}(x_3, x_5)).
 \end{aligned}$$

Note that $\frac{1}{2}D(x_4, Tx_4) = \frac{1}{2}d(x_4, x_2) = \frac{5}{2} > 1 = d(x_4, x_5)$, and $\frac{1}{2}D(x_5, Tx_5) = \frac{1}{2}d(x_5, x_3) = \frac{5}{2} > 1 = d(x_5, x_4)$.

Hence

$$\frac{1}{2}(D(x, Tx) \leq d(x, y))$$

implies

$$\alpha(Tx, Ty)H(Tx, Ty) \leq \max\{1, N_T(x, y)\psi(M_{TT}(x, y))$$

holds for all $x, y \in X$. Thus all the conditions of Corollary 2.14 are satisfied. On the other hand, if we take $x = x_4, y = x_5$, then

$$\alpha(Tx_4, Tx_5)H(Tx_4, Tx_5) = d(x_2, x_3) = 8 > \psi(d(x_4, x_5)) = \psi(1) = \frac{8}{9}.$$

Hence,

$$\alpha(Tx_4, Tx_5)H(Tx_4, Tx_5) = 8 \not\leq \frac{8}{9} = \psi(d(x_4, x_5)).$$

Consequently, Theorem 1.14 is not applicable in this case. Note that Theorem 1.14 is a generalization of Theorem 1.9. Now

$$\begin{aligned} n_{T,T}(x_4, x_5) &= \frac{\max\{d(x_4, x_5), D(x_4, Tx_4) + D(x_5, Tx_5), D(x_4, Tx_5) + D(x_5, Tx_4)\}}{1 + \delta(x_4, Tx_4) + \delta(x_5, Tx_5)} \\ &= \frac{\max\{d(x_4, x_5), d(x_4, x_2) + d(x_5, x_3), d(x_4, x_3) + d(x_5, x_2)\}}{1 + d(x_4, x_2) + d(x_5, x_3)} \\ &= \frac{\max\{1, 10, 10\}}{11} = \frac{10}{11} \neq 1 \end{aligned}$$

$$\begin{aligned} m_{T,T}(x_4, x_5) &= \max\left\{d(x_4, x_5), D(x_4, Tx_4), D(x_5, Tx_5), \frac{D(x_4, Tx_5) + D(x_5, Tx_4)}{2}\right\} \\ &= \max\{1, 5, 5, 5\} = 5 \end{aligned}$$

implies that $\alpha(Tx_4, Tx_5)H(Tx_4, Tx_5) = 8 \not\geq \frac{50}{11} = n_{T,T}(x_4, x_5)m_{T,T}(x_4, x_5)$. Hence, Theorem 1.13 which is a generalization of Theorem 1.11 does not hold in this case.

Coincidence and common fixed point results in b-metric spaces

As an application of Theorem 2.1, we obtain the existence of coincidence and common fixed point of Suzuki type $(\alpha_* - \psi)$ – hybrid pair of operators in b-metric spaces.

Theorem 3.1 Let (X, d) be a b-metric space and (g, T) a Suzuki type $(\alpha_* - \psi)$ – hybrid pair of operators such that T an $(\alpha_* - g)$ –admissible mapping. Suppose that there exists $x_0 \in X$ and $gx_1 \in Tx_0$ such that $\alpha(gx_0, gx_1) \geq 1$ and for any sequence $\{x_n\}$ in X with $gx_n \rightarrow gx$, we have $\alpha(gx_n, gx) \geq 1$ for all $n \in \mathbb{Z}^+$. Then there exists x in X such that $gx \in Tx$ provided that $T(X) \subseteq g(X)$ and $g(X)$ is complete. Moreover, if there is some $gy \in Ty$ such that $gx \neq gy$ and $\alpha(gx, gy) \geq 1$ then $d(gx, gy) \geq \frac{1}{2}$. Further, $F(g, T)$ is nonempty if any of the following conditions hold:

f₁-: The hybrid pair (g, T) is w – compatible, $\lim_{n \rightarrow +\infty} g^n(x) = w$ for some $w \in X$ and $x \in C(g, T)$ and g is continuous at w .

f₂-: The mapping g is T – weakly commuting at some $x \in C(g, T)$ and $g^2x = gx$.

f₃-: The mapping g is continuous at at some $x \in C(g, T)$ and $\lim_{n \rightarrow +\infty} g^n(w) = x$ for some $w \in X$.

Proof. Lemma 1.3, there is a set $E \subseteq X$ such that $g: E \rightarrow X$ is one-to-one and $g(E) = g(X)$. Define a mapping $T: g(E) \rightarrow CB(X)$ by $Tgx = Tx$ for all $g(x) \in g(E)$. The mapping T is well defined because g is one-to-one. Since (g, T) is a Suzuki type $(\alpha_* - \psi)$ – hybrid pair of operators, therefore

$$\begin{cases} \frac{1}{2}D(gx, Tx) \leq bd(gx, gy) \\ \text{implies} \\ \alpha_*(Tx, Ty)H(Tx, Ty) \leq \max\{1, N_{g,T}(x, y)\}\psi(M_{g,T}(x, y)) \\ = \max\left\{1, \frac{\max\{d(gx, gy), D(gx, Tx) + D(gy, Ty), D(gx, Ty) + D(gy, Tx)\}}{b(1 + \delta(gx, Tx) + \delta(gy, Ty))}\right\} \\ \psi\left(\max\left\{d(gx, gy), D(gx, Tx), D(gy, Ty), \frac{D(gx, Ty) + D(gy, Tx)}{2b}\right\}\right) \end{cases}$$

for all $x, y \in X$ for some $\psi \in \Psi_4$ and $\varphi \in \Phi$. Thus

$$\begin{cases} \frac{1}{2}D(gx, Tgx) \leq bd(gx, gy) \\ \text{implies} \\ \alpha_*(Tgx, Tgy)H(Tgx, Tgy) \leq \max\{1, N_T(x, y)\}\psi(M_T(x, y)) \\ = \max\left\{1, \frac{\max\{d(gx, gy), D(gx, Tgx) + D(gy, Tgy), D(gx, Tgy) + D(gy, Tgx)\}}{b(1 + \delta(gx, Tgx) + \delta(gy, Tgy))}\right\} \\ \psi\left(\max\left\{d(gx, gy), D(gx, Tgx), D(gy, Tgy), \frac{D(gx, Tgy) + D(gy, Tgx)}{2b}\right\}\right) \end{cases}$$

for all $gx, gy \in g(E)$. Since $g(E) = g(X)$ is complete. By the given assumption, there exists $x_0 \in X$ and $gx_1 \in Tx_0$ such that $\alpha(gx_0, gx_1) \geq 1$. As T is $(\alpha_* - g)$ – admissible, we have $\alpha(Tx, Ty) \geq 1$ which implies that $\alpha(Tgx, Tgy) \geq 1$. Thus T is α_* – admissible. Hence T satisfies all the conditions of Theorem 2.1. Consequently, T is an MWP operator on $g(E)$, and we obtain a point $u \in g(E)$ such that $u \in Tu$. Since $u \in g(E)$, there is a point x in X such that $gx = u$. This implies that $gx \in Tgx = Tx$. By Theorem 2.1 if there is some $w \in Tw$ such that $u \neq w$ and $\alpha(u, w) \geq 1$, then we have $d(u, w) \geq \frac{1}{2}$ if $\max\{1, N_{g,T}(x, y)\} = N_{g,T}(x, y)$. For $w \in Tw$ there is a y in X such that $gy = w$ and $gy \in Tgy =$

Ty . Consequently, $d(gx, gy) \geq \frac{1}{2}$. Now we prove that $F(g, T) \neq \emptyset$. First consider the case when (C_1) holds. Since the pair (g, T) is w – compatible and $\lim_{n \rightarrow +\infty} g^n(x) = u$ for some $u \in X$, the continuity of g at u implies that $gu = u$ and $\lim_{n \rightarrow +\infty} g^n(x) = gu$. Now $\alpha(g^n(x), gu) \geq 1$ and $(\alpha_* - g)$ – admissibility of T imply that $\alpha(Tg^{n-1}(x), Tu) \geq 1$. By w – compatibility of the pair (g, T) , we have $g^n(x) \in T(g^{n-1}(x))$, that is $g^n(x) \in C(g, T)$ for all $n \in \mathbb{N}$. Note that

$$\frac{1}{2}D(g^n(x), T(g^{n-1}(x))) \leq d(g^n(x), g^n(x)) = 0 \leq bd(gg^{n-1}(x), gu).$$

Since (g, T) is a generalized Suzuki type $(\alpha_* - \psi)$ – hybrid pair of operators, therefore

$$\begin{aligned} & D(g^n x, Tu) \\ & \leq H(Tg^{n-1}x, Tu) \leq \alpha(Tg^{n-1}(x), Tu)H(Tg^{n-1}x, Tu) \\ & \leq \max \left\{ 1, \frac{\max\{d(g^n x, gu), D(g^n x, Tg^{n-1}x) + D(gy, Tu), D(g^n x, Tu) + D(gu, Tg^{n-1}x)\}}{b(1 + \delta(g^n x, Tg^{n-1}x) + \delta(gu, Tu))} \right\} \\ & \leq \psi \max \left\{ d(g^n x, gu), D(g^n x, Tg^{n-1}x), D(gu, Tu), \frac{D(g^n x, Tu) + D(gu, Tg^{n-1}x)}{2b} \right\} \\ & \leq \max \left\{ 1, \frac{\max\{d(g^n x, gu), d(g^n x, g^n x) + D(gu, Tu), D(g^n x, Tu) + d(gu, g^n x)\}}{b(1 + d(g^n x, g^n x) + D(gu, Tu))} \right\} \\ & \leq \psi \max \left\{ d(g^n x, gu), d(g^n x, g^n x), D(gu, Tu), \frac{D(g^n x, Tu) + d(gu, g^n x)}{2b} \right\}. \end{aligned}$$

On taking limit as $n \rightarrow +\infty$, we obtain that

$$D(gu, Tu) \leq \max \left\{ 1, \frac{D(gu, Tu)}{b(1 + D(gu, Tu))} \right\} \psi(D(gu, Tu)) = \psi(D(gu, Tu)).$$

If $D(gu, Tu) > 0$, then we have $D(gu, Tu) < D(gu, Tu)$, a contradiction. Consequently, $u = gu \in Tu$. That is $F(g, T) \neq \emptyset$. Now let (C_2) hold, then $g^2x = gx$ for some $x \in C(g, T)$. By T – weakly commuting of g , we have $gx = g^2x \in Tgx$. Hence $gx \in F(g, T)$. In case (C_3) holds, $\lim_{n \rightarrow +\infty} g^n(u) = x$ for some $u \in X$ and $x \in C(g, T)$. By continuity of g , we obtain that $x = gx \in Tx$.

Hence $F(g, T) \neq \emptyset$.

Corollary 3.2 Let (X, d) be a b-metric space and (g, T) a hybrid pair such that T is an $(\alpha_* - g)$ - admissible. If there exists a $\psi \in \Psi_4$ such that

$$\frac{1}{2}D(gx, Tx) \leq bd(gx, gy)$$

implies that

$$\alpha_*(Tx, Ty)H(Tx, Ty) \leq \max\{1, N_{g,T}(x, y)\}\psi(d(x, y))$$

for all $x, y \in X$. Suppose that there exists $x_0 \in X$ and $gx_1 \in Tx_0$ such that $\alpha(gx_0, gx_1) \geq 1$ and for any sequence $\{x_n\}$ in X such that $gx_n \rightarrow gx$, we have $\alpha(gx_n, gx) \geq 1$ for all $n \in \mathbb{Z}^+$. Then there exists x in X such that $gx \in Tx$ provided that $T(X) \subseteq g(X)$ and $g(X)$ is complete. Moreover, if there is some $gy \in Ty$ such that $gx \neq gy$ and $\alpha(gx, gy) \geq 1$, then $d(gx, gy) \geq \frac{1}{2}$. Further, $F(g, T)$ is nonempty if the conditions (j₁)-(j₃) Theorem 3.1 hold.

Corollary 3.3 Let (X, d) be a b-metric space and (g, T) a hybrid pair such that T is an $(\alpha_* - g)$ - admissible. If there exists a $\psi \in \Psi_4$ such that

$$\frac{1}{2}D(gx, Tx) \leq bd(gx, gy)$$

implies

$$\alpha_*(Tx, Ty)H(Tx, Ty) \leq \psi(d(x, y))$$

for all $x, y \in X$. Suppose that there exists $x_0 \in X$ and $gx_1 \in Tx_0$ such that $\alpha(gx_0, gx_1) \geq 1$ and for any sequence $\{x_n\}$ in X such that $gx_n \rightarrow gx$, we have $\alpha(gx_n, gx) \geq 1$ for all $n \in \mathbb{Z}^+$. Then there exists x in X such that $gx \in Tx$ provided that $T(X) \subseteq g(X)$ and $g(X)$ is complete. Moreover, if there is some $gy \in Ty$ such that $gx \neq gy$ and $\alpha(gx, gy) \geq 1$, then $d(gx, gy) \geq \frac{1}{2}$. Further, $F(g, T)$ is nonempty if the conditions (j₁)-(j₃) in Theorem 3.1 hold.

Data dependence of fixed point sets and Ulam-Hyers stability results

Consider the following class of functions

$$\Theta = \{\xi: \mathbb{R}^+ \rightarrow \mathbb{R}^+ \text{ such that } \xi \text{ is increasing and continuous at } 0\}.$$

Let (X, d) be a b-metric space and $T: X \rightarrow P(X)$. The fixed point problem of T is to find an $x \in X$ such that

$$x \in Tx. \tag{4.1}$$

Inequality (4.1) is also known as fixed point inclusion. The fixed point inclusion is said to be generalized Ulam-Hyers stable if there exists a function $\xi \in \Theta$ such that for each $\varepsilon > 0$ and for each solution u_* of the inequality

$$D(u, Tu) \leq \varepsilon \quad (4.2)$$

there exists a solution z_* of the fixed point problem (4.1) such that $d(u_*, z_*) \leq \xi(\varepsilon)$.

Further, if there exists a $c > 0$ such that $\xi(t) = ct$ for each $t \in \mathbb{R}^+$, then the fixed point problem (4.1) is said to be Ulam-Hyers stable. Let $F(T)$ and U be the sets of solutions of (4.1) and (4.2), respectively. For more on Ulam-Hyers stability of fixed point problems, we refer the interested reader to (Ulam, 1964), (Lazar, 2012), (Petru et al, 2011), (Rus, 2009), (Hyers, 1941). Let (X, d) be a b -metric space and $T: X \rightarrow Cl(X)$ be a multivalued mapping then $E(T) = \{x \in X: \{x\} = Tx\}$.

Define a multivalued operator $T^\infty: G(T) \rightarrow P(F(T))$ by

$$T^\infty(x, y) = \{z \in F(T): \text{there is an ssa of } T \text{ at } (x, y) \text{ converging to } z\}$$

where $G(T) = \{(x, y): x \in X, y \in Tx\}$ is a graph of T .

A selection of $T: X \rightarrow P(X)$ is a single valued mapping $t: X \rightarrow X$ such that $tx \in Tx$ for all $x \in X$.

Definition 4.1 (Rus et al, 2003). Let (X, d) be a metric space and $c > 0$. An MWP operator $T: X \rightarrow P(X)$ is called ac – multivalued weakly Picard (briefly c – MWP) operator if there exists a selection t^∞ of T^∞ such that $d(x, t^\infty(x, y)) \leq cd(x, y)$ for all $(x, y) \in G(T)$.

One of the main results concerning c – MWP operators is the following:

Theorem 4.2 (Rus, 2001). Let (X, d) be a metric space and $T_1, T_2: X \rightarrow P(X)$. If T_i is a c_i – MWP operator for each $i \in \{1, 2\}$ and there exists $\lambda > 0$ such that $H(T_1x, T_2x) \leq \lambda$, for all $x \in X$. Then $H(F(T_1), F(T_2)) \leq \lambda \max\{c_1, c_2\}$.

Now we prove the following result.

Theorem 4.3 Let (X, d) be a complete b -metric space and $\alpha: X \times X \rightarrow \mathbb{R}^+$. Suppose that

g₁:- for each $i \in \{1, 2\}$, $T_i: X \rightarrow Cl(X)$ are multivalued operators such that

$$\frac{1}{2}D(x, T_i x) \leq bd(x, y)$$

implies that

$$\alpha_*(T_i x, T_i y)H(T_i x, T_i y) \leq \max\{1, N_{T_i}(x, y)\}\psi_i(d(x, y)) \quad (4.3)$$

for all $x, y \in X, \psi_i \in \Psi_4$.

g₂:- for each $i \in \{1, 2\}$, T_i is α_* – admissible mapping,

g₃:- there exists $x_0 \in X$ and $x_1 \in T_i x_0$ such that $\alpha(x_0, x_1) \geq 1$ for each $i \in \{1, 2\}$,

g₄:- if there is a sequence $\{x_n\}$ in X such that $x_n \rightarrow x$, then $\alpha(x_n, x) \geq 1$ for all $n \in \mathbb{Z}^+$,

g₅:- there exists $\lambda > 0$ such that $H(T_1 x, T_2 x) \leq \lambda$, for all $x \in X$.

Then $Fix(T_i) \in Cl(X), i \in \{1, 2\}$ and each T_i is an MWP operator such that

$$H(Fix(T_1), Fix(T_2)) \leq b\max\{\lambda_1, \lambda_2\}$$

where $\lambda_i = \sum_{k=0}^{\infty} b^k \psi_i^k(\lambda)$ for each $i \in \{1, 2\}$.

Proof. From Theorem 2.1, it follows that $Fix(T_i) \neq \emptyset$ for each $i \in \{1, 2\}$. Let $\{x_n\}$ be a sequence in $Fix(T_1)$ such that $x_n \rightarrow z$ as $n \rightarrow +\infty$. This implies that $\alpha(x_n, z) \geq 1$. Since T_1 is α_* – admissible mapping, $\alpha(T_1 x_n, T_1 z) \geq 1$. As

$$\frac{1}{2}D(x_n, T_1 x_n) = 0 \leq bd(z, x_n),$$

so we get

$$\begin{aligned}
 & D(z, T_1 z) \\
 & \leq bd(z, x_n) + bD(x_n, T_1 z) \\
 & \leq bd(z, x_n) + bH(T_1 z, T_1 x_n) \leq bd(z, x_n) + b\alpha(T_1 x_n, T_1 z)H(T_1 z, T_1 x_n) \\
 & \leq bd(z, x_n) + \max\{1, N_{T_1}(x, y)\}\psi_1(d(x, y)) \\
 & \leq \left(b \max \left\{ 1, \frac{\max\{d(z, x_n), D(z, T_1 z) + D(x_n, T_1 x_n)\}}{b(1 + \delta(z, T_1 z) + \delta(x_n, T_1 x_n))} \right\} \right) \psi_1(d(z, x_n)) \\
 & \quad + bd(z, x_n) \\
 & \leq \left(b \max \left\{ 1, \frac{\max\{d(z, x_n), D(z, T_1 z) + d(x_n, x_n)\}}{b(1 + D(z, T_1 z) + d(x_n, x_n))} \right\} \right) \psi_1(d(z, x_n)) \\
 & \quad + bd(z, x_n)
 \end{aligned}$$

On taking limit as $n \rightarrow +\infty$, we obtain that $D(z, T_1 z) \leq 0$, that is, $z \in T_1 z$ and hence $F(T_1)$ is closed.

Similarly, $F(T_2)$ is a closed subset of X .

From Corollary 2.2, we conclude that T_i for each $i \in \{1, 2\}$ is an MWP operator.

By a similar process as followed in Theorem 2.1 starting from $x_1 \in F(T_1)$ and $x_2 \in T_2 x_1$, we obtain a sequence $\{x_n\}$ such that $x_{n+1} \in T_2 x_n$ for all $n \geq 1$, $x_{n+1} \neq x_n$, $\alpha(x_{n+1}, x_{n+2}) \geq 1$, $0 < D(x_{n+1}, T x_{n+1}) \leq \psi_2(d(x_n, x_{n+1}))$ and

$$0 < d(x_{n+1}, x_{n+2}) < \psi_2^n(q\psi(c_0)) \quad (4.4)$$

for all $n \geq 1$, where $c_0 = d(x_1, x_2)$.

Following the arguments similar to those in the proof of Theorem 2.1, we conclude that $\{x_n\}$ is a Cauchy sequence and there is an element u in X such that $x_n \rightarrow u$ as $n \rightarrow +\infty$ and $u \in T_2 u$.

Note that

$$\begin{aligned}
 & d(x_n, x_{n+p}) \\
 & \leq bd(x_n, x_{n+1}) + b^2d(x_{n+1}, x_{n+2}) + \dots + b^{p-1}d(x_{n+p-2}, x_{n+p-1}) \\
 & \quad + b^pd(x_{n+p-1}, x_{n+p}) \\
 & \leq b\psi_2^{n-1}(q\psi_2(c_0)) + b^2\psi_2^n(q\psi_2(c_0)) + \dots + b^{p-1}\psi_2^{n+p-3}(q\psi_2(c_0)) \\
 & \quad + b^p\psi_2^{n+p-2}(q\psi_2(c_0)) \\
 & \leq \frac{1}{b^{n-2}}(b^{n-1}\psi_2^{n-1}(q\psi_2(c_0)) + b^n\psi_2^n(q\psi_2(c_0)) + \dots + b^{n+p-2}\psi_2^{n+p-2}(q\psi_2(c_0))) \\
 & = \frac{1}{b^{n-2}} \sum_{k=n-1}^{n+p-2} b^k\psi_2^k(q\psi_2(c_0)) \leq \frac{1}{b^{n-2}} \sum_{k=n-1}^{n+p-2} b^k\psi_2^k(q\psi_2(\lambda)) \\
 & = \frac{1}{b^{n-2}} \left(\sum_{k=0}^{n+p-2} b^k\psi_2^k(q\psi_2(\lambda)) - \sum_{k=0}^{n-1} b^k\psi_2^k(q\psi_2(\lambda)) + b^{n-1}\psi_2^{n-1}(q\psi_2(\lambda)) \right).
 \end{aligned}$$

That is,

$$\begin{aligned}
 d(x_n, x_{n+p}) & \leq \frac{1}{b^{n-2}} \left(\sum_{k=0}^{n+p-2} b^k\psi_2^k(q\psi_2(\lambda)) - \sum_{k=0}^{n-1} b^k\psi_2^k(q\psi_2(\lambda)) \right. \\
 & \quad \left. + b^{n-1}\psi_2^{n-1}(q\psi_2(\lambda)) \right).
 \end{aligned} \tag{4.5}$$

On taking limit as $p \rightarrow +\infty$, we get

$$\begin{aligned}
 d(x_n, u) & \leq \frac{1}{b^{n-2}} \left(\sum_{k=0}^{\infty} b^k\psi_2^k(q\psi_2(\lambda)) - \sum_{k=0}^{n-1} b^k\psi_2^k(q\psi_2(\lambda)) \right. \\
 & \quad \left. + b^{n-1}\psi_2^{n-1}(q\psi_2(\lambda)) \right).
 \end{aligned} \tag{4.6}$$

By Lemma 1.6, $\sum_{k=0}^{\infty} b^k\psi_2^k(t)$ converges for any $t \geq 0$, there exists $\lambda_2 \geq 0$ such that $\sum_{k=0}^{\infty} b^k\psi_2^k(\lambda) = \lambda_2$ and hence

$$d(x_n, u) \leq \frac{1}{b^{n-2}} \left(\lambda_2 - \sum_{k=0}^{n-1} b^k \psi_2^k(q\psi(\lambda)) + b^{n-1} \psi_2^{n-1}(q\psi_2(\lambda)) \right). \quad (4.7)$$

For $n = 1$, we get $d(x_1, u) \leq b\lambda_2$. Thus for $x_0 \in F(T_1)$, there exists $u \in F(T_2)$ such that $d(x_0, u) \leq b\lambda_2$. Similarly for each $z_0 \in F(T_2)$, we get $v \in F(T_1)$ and $\lambda_1 \geq 0$ such that $d(z_0, v) \leq b\lambda_1$. It follows from Lemma 1.4 that

$$H(F(T_1), F(T_2)) \leq b \max\{\lambda_1, \lambda_2\}.$$

Now we discuss the Ulam-Hyers stability results.

Theorem 4.4 *Let (X, d) be a b -metric space and $T: X \rightarrow Cl(X)$. Assume that all the hypotheses of Corollary 2.3 hold. Then we have*

h₁-: *The fixed point inclusion (4.1) is ζ_i^{-1} – generalized Ulam-Hyers stable for $i = 1, 2$, provided that for each $x \in F(T)$ there exists $z \in U$ such that $\alpha(x, z) \geq 1$, where $\zeta_1, \zeta_2: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ defined by $\zeta_1(t) = t - b^2\psi(t)$, $\zeta_2(t) = t - b\psi(t)$ are strictly increasing, onto and continuous at $t = 0$.*

h₂-: *If $E(T) \neq \emptyset$, then the fixed point inclusion (4.1) is ζ_i^{-1} – generalized Ulam-Hyers stable for $i = 3, 4$, provided that for $x \in E(T)$ there exists $z \in U$ such that $\alpha(x, z) \geq 1$, $\zeta_3, \zeta_4: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ defined by $\zeta_3(t) = t - b\psi(t)$, $\zeta_4(t) = t - t\psi(t)$ are strictly increasing, onto and continuous at $t = 0$.*

h₃-: *(Estimate between the fixed point sets of two multivalued mappings) If $S: X \rightarrow Cl(X)$ is such that for $x \in F(S)$ there exists $z \in F(T)$ with $\alpha(x, z) \geq 1$ and for $x \in F(T)$ there exists $z \in F(S)$ with $\alpha(x, z) \geq 1$, and $H(S(x), T(x)) \leq \lambda$ for all $x \in X$, then $H(F(S), F(T)) \leq \max_{i=1,2} \zeta_i^{-1}(b^2\lambda)$ where ζ_i is same as in (h₁) for each $i = 1, 2$.*

h₄-: *(Estimate between the fixed point sets of two multivalued mappings) If $S: X \rightarrow Cl(X)$ is such that for $x \in F(S)$, there exists $z \in E(T)$ with $\alpha(x, z) \geq 1$ and for $x \in E(T)$ there exists $z \in F(S)$ with $\alpha(x, z) \geq 1$, and $H(S(x), T(x)) \leq \lambda$ for all $x \in X$, then $H(F(S), F(T)) \leq \max_{i=3,4} \zeta_i^{-1}(b\lambda)$ where ζ_i is same as in (h₂) for each $i = 3, 4$.*

h₅-: *(Well-posedness of the fixed point problem with respect to b -metric d) If for any sequence $\{x_n\}$ in X , there exists a unique point $x^* \in E(T)$ such that $\alpha(x_n, x^*) \geq 1$ and $\lim_{n \rightarrow +\infty} D(x_n, Tx_n) = 0$, then $\lim_{n \rightarrow +\infty} d(x_n, x^*) = 0$.*

h₆-: (Well-posedness of the fixed point problem with respect to b -metric H) If for any sequence $\{x_n\}$ in X , there exists a unique point $x^* \in E(T)$ such that $\alpha(x_n, x^*) \geq 1$ and $\lim_{n \rightarrow +\infty} H(\{x_n\}, Tx_n) = 0$, then $\lim_{n \rightarrow +\infty} d(x_n, x^*) = 0$.

h₇-: (Limit shadowing property of the multivalued operators) If for any sequence $\{x_n\}$ in X , there exists a unique point $x^* \in E(T)$ with $\alpha(x_n, x^*) \geq 1$ and $\lim_{n \rightarrow +\infty} D(x_n, Tx_n) = 0$, then there exists a sequence of successive approximations $\{y_n\}$ such that $\lim_{n \rightarrow +\infty} d(x_n, y_n) = 0$.

Proof. (h₁) From Corollary 2.3, T is an MWP operator and hence $F(T)$ is nonempty. If $x^* \in F(T)$, then by given condition there exists a $y^* \in U$ such that $\alpha(x^*, y^*) \geq 1$. The α_* -admissibility of T gives that $\alpha(Tx^*, Ty^*) \geq 1$. Since $y^* \in U$, for any given $\varepsilon > 0$, we have $D(y^*, Ty^*) \leq \varepsilon$. Note that

$$\frac{1}{2}D(x^*, Tx^*) = 0 \leq bd(x^*, y^*).$$

Then

$$\begin{aligned} d(x^*, y^*) &\leq bD(x^*, Tx^*) + bD(Tx^*, y^*) = bD(Tx^*, y^*) \\ &\leq b^2(H(Tx^*, Ty^*) + D(Ty^*, y^*)) \\ &\leq b^2(\alpha(Tx^*, Ty^*)H(Tx^*, Ty^*) + \varepsilon) \\ (u-3) \quad &\leq b^2 \left(\max \left\{ 1, \frac{d(x^*, y^*)}{b(1 + \delta(x^*, Tx^*) + \delta(y^*, Ty^*))} \right\} \psi(d(x^*, y^*)) + \varepsilon \right) \\ &\leq b^2 \left(\max \left\{ 1, \frac{d(x^*, y^*)}{b(1 + d(x^*, x^*) + D(y^*, Ty^*))} \right\} \psi(d(x^*, y^*)) + \varepsilon \right) \\ &\leq b^2 \left(\max \left\{ 1, \frac{d(x^*, y^*)}{b} \right\} \psi(d(x^*, y^*)) + \varepsilon \right). \end{aligned}$$

If $\max \left\{ 1, \frac{d(x^*, y^*)}{b} \right\} = 1$, then we have $d(x^*, y^*) \leq b^2(\psi(d(x^*, y^*)) + \varepsilon)$.

If $\zeta_1(d(x^*, y^*)) = d(x^*, y^*) - b^2\psi(d(x^*, y^*))$, then from the above inequality we get $\zeta_1(d(x^*, y^*)) \leq b^2\varepsilon$ and hence $d(x^*, y^*) \leq \zeta_1^{-1}(b^2\varepsilon)$. Consequently, the fixed point inclusion (4.1) is ξ -generalized Ulam-Hyers stable, where $\xi = \zeta_1^{-1}$.

If $\max \left\{ 1, \frac{d(x^*, y^*)}{b} \right\} = \frac{d(x^*, y^*)}{b}$, then $d(x^*, y^*) \geq b$. From (4.1) we obtain

that

$$\begin{aligned} d(x^*, y^*) &\leq b^2 \left(\frac{d(x^*, y^*)}{b} \psi(d(x^*, y^*)) + \varepsilon \right) \\ &\leq bd(x^*, y^*) \psi(d(x^*, y^*)) + b^2 \varepsilon \\ &\leq bd(x^*, y^*) \psi(d(x^*, y^*)) + b^2 \varepsilon. \end{aligned}$$

Now if $\zeta_2(d(x^*, y^*)) = d(x^*, y^*) - bd(x^*, y^*) \psi(d(x^*, y^*))$, then from the above inequality we get $\zeta_2(d(x^*, y^*)) \leq b^2 \varepsilon$ and hence $d(x^*, y^*) \leq \zeta_2^{-1}(b^2 \varepsilon)$. Consequently, the fixed point inclusion (4.1) is ξ – generalized Ulam-Hyers stable, where $\xi = \zeta_2^{-1}$.

(h₂) Let $E(T) \neq \emptyset$, and $x^* \in E(T)$ then

$$d(x^*, y^*) = D(Tx^*, y^*) \leq b(H(Tx^*, Ty^*) + D(Ty^*, y^*)).$$

Following the arguments similar to those in the proof of (h₁), the result follows.

(h₃) Let $x^* \in F(S)$, then there exists a $y^* \in F(T)$ such that $\alpha(x^*, y^*) \geq$

1. By α_* – admissibility of T we get $\alpha(Tx^*, Ty^*) \geq 1$. Note that

$$\frac{1}{2}D(y^*, Ty^*) = 0 \leq bd(x^*, y^*).$$

Then by the given assumption on T , we obtain that

$$\begin{aligned} d(x^*, y^*) &\leq bD(x^*, Sx^*) + bD(Sx^*, y^*) \\ &= bD(Sx^*, y^*) \leq b^2(H(Sx^*, Tx^*) + D(Tx^*, y^*)) \\ &\leq b^2(H(Sx^*, Tx^*) + H(Tx^*, Ty^*)) \leq b^2(\lambda + H(Tx^*, Ty^*)) \\ &\leq b^2(\lambda + \alpha(Tx^*, Ty^*)H(Tx^*, Ty^*)) \\ &\leq b^2 \left(\lambda + \max \left\{ 1, \frac{d(x^*, y^*)}{b(1+\delta(x^*, Tx^*)+\delta(y^*, Ty^*))} \right\} \psi(d(x^*, y^*)) \right) \\ &\leq b^2 \left(\lambda + \max \left\{ 1, \frac{d(x^*, y^*)}{b(1+d(x^*, x^*)+D(y^*, Ty^*))} \right\} \psi(d(x^*, y^*)) \right) \\ &\leq b^2 \left(\lambda + \max \left\{ 1, \frac{d(x^*, y^*)}{b(1+d(x^*, x^*)+D(y^*, Ty^*))} \right\} \psi(d(x^*, y^*)) \right) \\ &\leq b^2(\lambda + \psi(d(x^*, y^*))). \end{aligned}$$

If $\max \left\{ 1, \frac{d(x^*, y^*)}{b} \right\} = 1$, and

$$\zeta_1(d(x^*, y^*)) = d(x^*, y^*) - b^2(\psi(d(x^*, y^*))),$$

then from the above inequality we get $\zeta_1(d(x^*, y^*)) \leq b^2 \lambda$. Consequently, for every $x^* \in F(S)$, there exists a $y^* \in F(T)$ such that $d(x^*, y^*) \leq \zeta_1^{-1}(b^2 \lambda)$. Similarly, it can be proved that for every $y^* \in F(T)$, there exists

a $x^* \in F(T)$ such that $d(x^*, y^*) \leq \zeta_1^{-1}(b^2\lambda)$. Hence by Lemma 1.4, we obtain that

$$H(F(S), F(T)) \leq \zeta_1^{-1}(b^2\lambda).$$

If $\max\left\{1, \frac{d(x^*, y^*)}{b}\right\} = \frac{d(x^*, y^*)}{b}$, then for $\zeta_1(d(x^*, y^*)) = d(x^*, y^*) - bd(x^*, y^*)\psi(d(x^*, y^*))$ we get

$$H(F(S), F(T)) \leq \zeta_2^{-1}(b^2\lambda).$$

Consequently,

$$H(F(S), F(T)) \leq \max_{i=1,2} \zeta_i^{-1}(b^2\lambda).$$

(h₄) This can be proved on the similar lines as in (h₃) using the definition of $E(T)$.

(h₅) If $\{x_n\}$ is a sequence in X , there exists a unique $x^* \in E(T)$ such that $\alpha(x_n, x^*) \geq 1$ and $\lim_{n \rightarrow +\infty} D(x_n, Tx_n) = 0$. Then there exists $u_n \in Tx_n$ such that $\lim_{n \rightarrow +\infty} D(x_n, Tx_n) = \lim_{n \rightarrow +\infty} d(x_n, u_n) = 0$. Since T is α_* -admissible, $\alpha(Tx_n, Tx^*) \geq 1$. As $\frac{1}{2}D(x^*, Tx^*) = 0 \leq bd(x_n, x^*)$, by given assumption we have

$$\begin{aligned} d(x_n, x^*) &\leq b(D(x_n, Tx_n) + D(Tx_n, x^*)) \\ &\leq b(D(x_n, Tx_n) + H(Tx_n, Tx^*)) \\ &\leq b(D(x_n, Tx_n) + \alpha(Tx_n, Tx^*)H(Tx_n, Tx^*)) \\ &\leq b(D(x_n, Tx_n) \\ &\quad + \max\left\{1, \frac{d(x_n, x^*)}{b(1+D(x_n, Tx_n)+D(x^*, Tx^*))}\right\} \psi(d(x_n, x^*))) \\ &\leq b(D(x_n, Tx_n) \\ &\quad + \max\left\{1, \frac{d(x_n, x^*)}{b(1+D(x_n, Tx_n))}\right\} \psi(d(x_n, x^*))) \\ &\leq b\left(D(x_n, Tx_n) + \max\left\{1, \frac{d(x_n, x^*)}{b}\right\} \psi(d(x_n, x^*))\right). \end{aligned}$$

If $\max\left\{1, \frac{d(x_n, x^*)}{b}\right\} = 1$, then we have

$$d(x_n, x^*) - b\psi(d(x_n, x^*)) \leq bD(x_n, Tx_n).$$

That is, $\zeta_3(d(x_n, x^*)) \leq bD(x_n, Tx_n)$. Similarly, if $\max\left\{1, \frac{d(x_n, x^*)}{b}\right\} = \frac{d(x_n, x^*)}{b}$, we get $\zeta_4(d(x_n, x^*)) \leq bD(x_n, Tx_n)$. This implies for each $i \in \{3, 4\}$ we get $d(x_n, x^*) \leq \zeta_i^{-1}(bD(x_n, Tx_n))$. On taking limit as $n \rightarrow +\infty$ and using the continuity of ζ_i at 0, for each $i \in \{3, 4\}$ we get the desired result.

(h₆) follows from (h₅) as $D(x_n, Tx_n) \leq H(\{x_n\}, Tx_n)$.

(h₇) From (h₅) it is clear that $\lim_{n \rightarrow +\infty} d(x_n, x^*) = 0$. Since $x^* \in E(T)$, so there exists a sequence of successive approximations defined by $y_n = x^*$ for all n such that $\lim_{n \rightarrow +\infty} d(x_n, y_n) = \lim_{n \rightarrow +\infty} d(x_n, x^*) = 0$.

Existence and stability of solutions of differential inclusions

Let $Cl_c(\mathbb{R})$ be collection of nonempty closed and convex subsets of \mathbb{R} and $F: \mathbb{R} \rightarrow Cl_c(\mathbb{R})$ a lower semicontinuous multivalued mapping. Consider the initial value problem

$$\begin{aligned} x'(t) &\in F(x(t)), \text{ for } t \in J, \\ x(t) &= x_0 \text{ for } t = a_1, \\ x &\in C(J), \end{aligned} \quad (4.8)$$

where $J = [a_1, a_2]$ and $C(J)$ is a Banach space of absolutely continuous real valued functions defined on J . Since \mathbb{R} with usual metric is paracompact, F is a lower semicontinuous multivalued mapping with $F(u)$ closed and convex for each $u \in \mathbb{R}$, by Michael's Theorem (Michael, 1956), there exists a continuous function $f: \mathbb{R} \rightarrow \mathbb{R}$ such that $f(u) \in F(u)$ for all $u \in \mathbb{R}$.

Now consider the following initial value problem

$$\begin{aligned} x'(t) &= f(x(t)), \text{ for } t \in J, \\ x(t) &= x_0 \text{ for } t = a_1, \\ x &\in C(J). \end{aligned} \quad (4.9)$$

Note that the solution of problem (4.9) is a solution of problem (4.8). Integrating from a_1 to t , we obtain

$$\int_{a_1}^t x'(s) ds = \int_{a_1}^t f(x(s)) ds,$$

that is,

$$x(t) = x_0 + \int_{a_1}^t f(x(s)) ds, \text{ for } t \in J. \quad (4.10)$$

On the other hand, if (4.10) holds then (4.9) holds. Thus (4.9) and (4.10) are equivalent.

Suppose that $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfies the following hypotheses:

$\int_{a_1}^t f(x(s))ds = 0$, for $t \in J$ if and only if $x(t) = x_0(t)$ for all $t \in J$.

There exists a nonnegative real number L_f such that $L_f(a_2 - a_1) < \frac{1}{b^2}$, where b is b-metric constant and for all $u, v \in \mathbb{R}$, the relation $\|f(u) - f(v)\| \leq L_f \|u(t) - v(t)\|$ holds.

Define $T: X \rightarrow X$, where $X = C(J)$ by

$$T(x(t)) = x_0 + \int_{a_1}^t f(x(s))ds, \text{ for } t \in J. \quad (4.11)$$

Let $d: C(J) \times C(J) \rightarrow \mathbb{R}^+$ be defined as

$$d(x, y) := \max_{t \in J} \|x(t) - y(t)\|^2.$$

Then $(C(J), d)$ is complete b-metric space. Define $\alpha: C(J) \times C(J) \rightarrow \mathbb{R}^+$ by

$$\alpha(x, y) = \begin{cases} k & \text{for } x \neq x_0, y \neq y_0, \text{ where } k \geq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Let $\psi: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be defined as $\psi(t) = L_f^2(a_2 - a_1)t$. Clearly, $\psi \in \Psi_4$.

First we show that mapping T is α_* -admissible. As $\alpha(x, y) \geq 1$ implies that $\alpha(x, y) = k$. For $x \neq x_0$ and $y \neq x_0$, from (i₁-) we have $Tx \neq x_0$ and $Ty \neq x_0$ on J . It follows that $\alpha_*(Tx, Ty) = k \geq 1$ and hence T is α_* -admissible. Now, by (i₂-) for all $x, y \in X$,

$$\begin{aligned} d(Tx, Ty) &= \max_{t \in J} \left\| \int_{a_1}^t f(x(s))ds - \int_{a_1}^t f(y(s))ds \right\|^2 \\ &\leq \max_{t \in J} \int_{a_1}^t \|f(x(s)) - f(y(s))\|^2 ds \end{aligned}$$

$$\begin{aligned}
 &\leq \max_{t \in J} \int_{a_1}^t L_f^2 \|x(s) - y(s)\|^2 ds \\
 &\leq L_f^2 \max_{t \in J} \int_{a_1}^t \max_{s \in J} \|x(s) - y(s)\|^2 ds \\
 &= L_f^2 d(x, y) \max_{t \in J} \int_{a_1}^t ds = L_f^2 (a_2 - a_1) d(x, y) = \psi(d(x, y)).
 \end{aligned}$$

By Corollary 2.3, we obtain the solution of problem (4.9) which provides the solution of problem (4.8) as well. Define the mappings $\zeta_1, \zeta_2: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ by

$$\begin{aligned}
 \zeta_1(t) &= t - b^2 L_f^2 (a_2 - a_1) t = t - b^2 \psi(t), \\
 \zeta_2(t) &= t - b L_f^2 (a_2 - a_1) t^2 = t - b t \psi(t),
 \end{aligned}$$

where $\psi(t) = L_f^2 (a_2 - a_1) t$. Clearly the mapping ζ_1 is strictly increasing and onto. Consequently, all the axioms of Theorem 4.4 hold with mapping ζ_1 . Hence the fixed point inclusion (4.8) is ζ_1^{-1} -generalized Ulam-Hyers stable. Now $\frac{d}{dt} \zeta_2(t) > 0$ if $1 - 2b L_f^2 (a_2 - a_1) t > 0$. As $b L_f^2 (a_2 - a_1) < 1$, hence the fixed point inclusion (4.8) is ζ_2^{-1} -generalized Ulam-Hyers stable if $t < \frac{1}{2}$.

Let $F: J \times \mathbb{R} \times \mathbb{R} \rightarrow Cl_c(\mathbb{R})$ be a lower semicontinuous multivalued mapping. Consider the initial value problem

$$\begin{aligned}
 x'(t) &\in F(t, x(t), x(t-h)), \text{ for } t \in J, \\
 x(t) &= x_0 \text{ for } t \in [a_1 - h, a_1], \\
 x &\in C(J),
 \end{aligned} \tag{4.12}$$

where h is a positive real number. Then there exists a selection f such that $f(s, u, v) \in F(s, u, v)$ for all $u, v \in \mathbb{R}$ and $s \in [a_1 - h, a_1]$, see (Michael, 1956). Note that any solution of the problem

$$\begin{aligned}
 x'(t) &= f(t, x(t), x(t-h)), \text{ for } t \in J, \\
 x(t) &= x_0 \text{ for } t \in [a_1 - h, a_1], \\
 x &\in C(J)
 \end{aligned} \tag{4.13}$$

is a solution for problem (4.12). Further, (4.13) is equivalent to

$$\begin{aligned} x(t) &= x_0 + \int_{a_1}^t f(s, x(s), x(s-h))ds, \text{ for } t \in J, \\ x(t) &= x_0 \text{ for } t \in [a_1-h, a_1]. \end{aligned}$$

We suppose that f satisfies the following hypotheses:

$$\int_{a_1}^t f(s, x(s), x(s-h))ds = 0, \text{ for } t \in J \text{ if and only if } x = x_0 \text{ on } J.$$

$$\|f(t, u_1, v_1) - f(t, u_2, v_2)\| \leq L_f(\|u_1 - u_2\| + \|v_1 - v_2\|),$$

for all $u_1, u_2, v_1, v_2 \in \mathbb{R}$, where $L_f(a_2 - a_1) < \frac{1}{4b^2}$ and b is a metric constant.

Define the operator $T: Y \rightarrow Y$, where $Y = C[a_1 - h, a_2] \times \mathbb{R} \times \mathbb{R}$ by

$$T(x(t)) = \begin{cases} x_0 + \int_{a_1}^t f(s, x(s), x(s-h))ds, & \text{for } t \in J, \\ x(t) = x_0 & \text{for } t \in [a_1 - h, a_1]. \end{cases} \quad (4.14)$$

From the definition of α , the admissibility of T follows. Now by (i₄-) for all $x, y \in Y$, we have

$$\begin{aligned} d(Tx, Ty) &= \max_{t \in J} \left\| \int_{a_1}^t f(s, x(s), x(s-h))ds - \int_{a_1}^t f(s, x(s), y(s-h))ds \right\|^2 \\ &\leq \max_{t \in J} \int_{a_1}^t \|f(s, x(s), x(s-h)) - f(s, x(s), y(s-h))\|^2 ds \\ &\leq \max_{t \in J} \int_{a_1}^t L_f^2 \|x(s) - y(s)\|^2 ds \\ &= 4L_f^2 \max_{t \in J} \int_{a_1}^t \|x(s) - y(s)\|^2 ds \\ &\leq 4L_f^2(a_2 - a_1)\psi(d(x, y)), \end{aligned}$$

where $\psi(t) = 4L_f^2(a_2 - a_1)t$. By Corollary 2.3, we obtain the solution of problem (4.13) which is also being the selection is the solution for (4.12).

If $\zeta_1(t) = t - 4b^2L_f^2(a_2 - a_1)t$ and $\zeta_2(t) = t - 4bL_f^2(a_2 - a_1)t^2$, then the

fixed point inclusion (4.12) is ζ_1^{-1} – generalized Ulam-Hyers stable. Now $\frac{d}{dt}\zeta_2(t) > 0$ if $1 - 8bL_f^2(a_2 - a_1)t > 0$. As $4bL_f^2(a_2 - a_1) < 1$, hence the fixed point inclusion (4.12) is ζ_2^{-1} – generalized Ulam-Hyers stable if $t < \frac{1}{2}$.

References

- Abbas, M., Ali, B., & Vetro, C. 2013. A Suzuki type fixed point theorem for a generalized multivalued mapping on partial Hausdorff metric spaces. *Topology and its Applications*, 160(3), pp.553-563. Available at: <https://doi.org/10.1016/j.topol.2013.01.006>.
- Abbas, M., Ćirić, Lj., Damjanović, B., & Khan, M.A. 2012. Coupled coincidence and common fixed point theorems for hybrid pair of mappings on a test. *Fixed Point Theory and Applications*, art.number:4. Available at: <https://doi.org/10.1186/1687-1812-2012-4>.
- Aleksić, S., Došenović, T., Mitrović, D.Z., & Radenović, S. 2019a. Remarks on common fixed point results for generalized α_* – ψ - contraction multivalued mappings in b -metric spaces. *Advances in Fixed Point Theory*, 9(1), pp.1-16. Available at: <https://doi.org/10.28919/afpt/3731>.
- Aleksić, S., Huang, H, Mitrović, D.Z., & Radenović, S. 2018. Remarks on some fixed point results in b -metric spaces. *Journal of Fixed Point Theory and Applications*, 20(art.number:147). Available at: <https://doi.org/10.1007/s11784-018-0626-2>.
- Aleksić, S., Mitrović, Z.D., & Radenović, S. 2019b. Picard sequences in b -metric spaces. *Fixed Point Theory*, 21(1), pp.35-46. Available at: <https://doi.org/10.24193/fpt-ro.2020.1.03>.
- Ali, B., & Abbas, M. 2017. Existence Ulam–Hyers stability of fixed point problem of generalized Suzuki type (α_*, ψ_φ) -contractive multivalued operators. *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas (RACSAM)*, 111(4), pp.1129-1146. Available at: <https://doi.org/10.1007/s13398-016-0351-x>.
- An, T.V., Dung, N.V., Kadelburg, Z., & Radenović, S. 2015a. Various generalizations of metric spaces and fixed point theorems. *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas (RACSAM)*, 109, pp.175-198. Available at: <https://doi.org/10.1007/s13398-014-0173-7>.
- An, T.V., Tuyen, L.Q., & Dung, N.V. 2015b. Stone-type theorem on b -metric spaces and applications. *Topology and its Applications*, 185-186, pp.50-64. Available at: <https://doi.org/10.1016/j.topol.2015.02.005>.
- Asl, J.H., Rezapour, Sh., & Shahzad, N. 2012. On fixed points of $\alpha - \psi$ -contractive multifunctions. *Fixed Point Theory and Applications*, art.number:212. Available at: <https://doi.org/10.1186/1687-1812-2012-212>.



Banach, S. 1922. Sur les opérations dans les ensembles abstraits et leur applications aux équations intégrales. *Fundamenta Mathematicae*, 3, pp.133-181 (in French). Available at: <https://doi.org/10.4064/fm-3-1-133-181>.

Berinde, V. 1993. Generalized contractions in quasimetric spaces. In: *Seminar on Fixed Point Theory*, Babeş-Bolyai University, Cluj-Napoca, pp.3-9.

Berinde, V. 1996. Sequences of operators and fixed points in quasimetric spaces. *Studia Universitatis Babeş-Bolyai Mathematica*, 41(4), pp.23-27.

Berinde, V. 1997. *Contractiuni generalizate si aplicatii*. Baia Mare: Editura Club Press (in Romanian).

Bhaskar, T.G., & Lakshmikantham, V. 2006. Fixed point theorems in partially ordered metric spaces and applications. *Nonlinear Analysis: Theory, Methods & Applications*, 65(7), pp.1379-1393. Available at: <https://doi.org/10.1016/j.na.2005.10.017>.

Bota, M.F., Ilea, V., Karapinar, E., & Mleşniţ, O. 2015. On α_* - ψ -contractive multivalued operators in b -metric spaces and applications. *Applied Mathematics & Information Sciences*, 9(5), pp.2611-2620 [online]. Available at: <http://www.naturalspublishing.com/Article.Asp?ArtclD=9576> [Accessed: 21 May 2020].

Chifu, C., & Petruşel, G. 2014. Fixed points for multivalued contractions in b -metric spaces with applications to fractals. *Taiwanese Journal of Mathematics*, 18(5), pp.1365-1375. Available at: <https://doi.org/10.11650/tjm.18.2014.4137>.

Czerwik, S. 1993. Contraction mappings in b -metric spaces. *Acta Mathematica et Informatica Universitatis Ostraviensis*, 1(1), pp.5-11 [online]. Available at: <https://dml.cz/dmlcz/120469> [Accessed: 21 May 2020].

Ćirić, Lj., Abbas, M., Rajović, M., & Ali, B. 2012. Suzuki type fixed point theorems for generalized multi-valued mappings on a set endowed with two b -metric. *Applied Mathematics and Computation*, 219(4), pp.1712-1723. Available at: <https://doi.org/10.1016/j.amc.2012.08.011>.

Fréchet, M. 1905. La notion d'écart et le calcul fonctionnel. *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences*, 140(1), pp.772-774 [online]. Available at: <https://www.biodiversitylibrary.org/item/31541#page/778/mode/1up> (in French) [Accessed: 21 May 2020].

Haghi, R.H., Rezapour, Sh., & Shahzad, N. 2011. Some fixed point generalisations are not real generalizations. *Nonlinear Analysis: Theory, Methods & Applications*, 74(5), pp.1799-1803. Available at: <https://doi.org/10.1016/j.na.2010.10.052>.

Hussain, N., Ćorić, D., Kadelburg, Z., & Radenović, S. 2012. Suzuki-type fixed point results in metric type spaces. *Fixed Point Theory and Applications*, art.number:126. Available at: <https://doi.org/10.1186/1687-1812-2012-126>.

Hyers, D.H. 1941. On the stability of the linear functional equation. In: *Proceedings of the National Academy of Sciences of the USA*, 27(4), pp.222-224, April 15. Available at: <https://doi.org/10.1073/pnas.27.4.222>.

Karapinar, E., Mitrović, Z.D., Ozturk, A., & Radenović, S. 2020. On a theorem of Ćirić in b -metric spaces. *Rendiconti del Circolo Matematico di Palermo Series 2*. Available at: <https://doi.org/10.1007/s12215-020-00491-9>.

- Khojasteh, F., Abbas, M., & Costache, S. 2014. Two new types of fixed point theorems in complete metric spaces. *Variational Analysis, Optimization, and Fixed Point Theory*, art.ID325840. Available at: <https://doi.org/10.1155/2014/325840>.
- Kirk, W., & Shahzad, N. 2014. *Fixed Point Theory in Distance Spaces*. Cham, Switzerland: Springer International Publishing.
- Latif, A., Parvaneh, V., Salimi, P., & Al-Mazrooei, A.E. 2015. Various Suzuki type theorems in b -metric spaces. *Journal of Nonlinear Sciences and Applications (JNSA)*, 8(4), pp.363-377. Available at: <https://doi.org/10.22436/jnsa.008.04.09>.
- Lazar, V.L. 2012. Ulam-Hyers stability for partial differential inclusions. *Electronic Journal of Qualitative Theory of Differential Equations*, 21, pp.1-19. Available at: <https://doi.org/10.14232/ejqtde.2012.1.21>.
- Michael, E. 1956. Continuous selections. I. *Annals of Mathematics*, 63(2), pp.361-382. Available at: <https://doi.org/10.2307/1969615>.
- Mitrović, Z.D. 2019. A note on the results of Suzuki, Miculescu and Mihail. *Journal of Fixed Point Theory and Applications*, 21(art.number:24). Available at: <https://doi.org/10.1007/s11784-019-0663-5>.
- Mitrović, Z.D., Parvaneh, V., Mlaiki, N., Hussain, N., & Radenović, S. 2020. On some new generalizations of Nadler contraction in b -metric spaces. *Cogent Mathematics and Statistics*, 7(1), art.number:1760189. Available at: <https://doi.org/10.1080/25742558.2020.1760189>.
- Mohammadi, B., Dinu, S., & Rezapour, Sh. 2013. Fixed points of Suzuki type quasi-contractions. *U.P.B. Scientific Bulletin, Series A*, 75(3), pp.3-12 [online]. Available at: https://www.scientificbulletin.upb.ro/rev_docs_arhiva/full89_199620.pdf. [Accessed: 21 May 2020].
- Nadler, Jr. S.B. 1969. Multi-valued contraction mappings. *Pacific Journal of Mathematics*, 30(2), pp.475-488. Available at: <https://doi.org/10.2140/pjm.1969.30.475>.
- Nieto, J.J., & Rodríguez-López, R. 2005. Contractive mapping theorems in partially ordered sets and applications to ordinary differential equation. *Order*, 22, pp.223-239. Available at: <https://doi.org/10.1007/s11083-005-9018-5>.
- Nieto, J.J., & Rodríguez-López, R. 2007. Existence and uniqueness of fixed point in partially ordered sets and applications to ordinary differential equations. *Acta Mathematica Sinica, English Series*, 23, pp.2205-2212. Available at: <https://doi.org/10.1007/s10114-005-0769-0>.
- Păcurar, M. 2010. A fixed point result for φ -contractions on b -metric spaces without the boundedness assumption. *Fasciculi Mathematici*, 43, pp.127-137 [online]. Available at: [http://www.math.put.poznan.pl/artykuly/FM43\(2010\)-PacurarM.pdf](http://www.math.put.poznan.pl/artykuly/FM43(2010)-PacurarM.pdf) [Accessed: 21 May 2020].
- Petru, T.P., Petrusel, A., & Yao, J.-C. 2011. Ulam-Hyers stability for operatorial equations inclusions via nonself operators. *Taiwanese Journal of Mathematics*, 15(5), pp.2195-2212. Available at: <https://doi.org/10.11650/twjrm/1500406430>.



Ran, A.C.M., & Reurings, M.C.B. 2004. A fixed point theorem in partially ordered sets and some applications to matrix equations. *Proceedings of the American Mathematical Society*, 132(5), pp.1435-1443. Available at: <https://doi.org/10.1090/S0002-9939-03-07220-4>.

Rhoads, B.E. 2015. Two new fixed point theorem. *Gen. Math. Notes*, 27(2), pp.123-132 [online]. Available at: https://www.emis.de/journals/GMN/yahoo_site_admin/assets/docs/12_GMN-7082-V27N2.154194604.pdf[Accessed: 21 May 2020].

Rus, I. A. 2001. *Generalized contractions and applications*. Cluj: University Press.

Rus, I. A. 2009. Remarks on Ulam stability of the operatorial equations. *Fixed Point Theory*, 10(2), pp.305-320 [online]. Available at: [http://www.math.ubbcluj.ro/~nodeacj/vol__10\(2009\)_no_2.php](http://www.math.ubbcluj.ro/~nodeacj/vol__10(2009)_no_2.php)[Accessed: 21 May 2020].

Rus, I. A., Petruşel, A., & Sîntămărian, A. 2003. Data dependence of the fixed point set of some multivalued weakly Picard operators. *Nonlinear Analysis: Theory, Methods & Applications*, 52(8), pp.1947-1959. Available at: [https://doi.org/10.1016/S0362-546X\(02\)00288-2](https://doi.org/10.1016/S0362-546X(02)00288-2).

Samet, B., Vetro, C., & Vetro, P. 2012. Fixed point theorems for $\alpha - \psi$ -contractive type mappings. *Nonlinear Analysis: Theory, Methods & Applications*, 75(4), pp.2154-2165. Available at: <https://doi.org/10.1016/j.na.2011.10.014>.

Singh, S.L., & Prasad, B. 2008. Some coincidence theorems and stability of iterative procedures. *Computers & Mathematics with Applications*, 55(11), pp.2512-2520. Available at: <https://doi.org/10.1016/j.camwa.2007.10.026>.

Suzuki, T. 2008. A generalized Banach contraction principle that characterizes metric completeness. *Proceedings of the American Mathematical Society*, 136(5), pp.1861-1869. Available at: <https://doi.org/10.1090/S0002-9939-07-09055-7>.

Ulam, S.M. 1964. *Problems in Modern Mathematics*. New York, NY: John Wiley and Sons.

РЕШЕНИЯ И УСТОЙЧИВОСТЬ ДИФФЕРЕНЦИАЛЬНЫХ
ВКЛЮЧЕНИЙ ПО УЛАМУ-ХАЙЕРСУ, ВКЛЮЧАЯ
РАЗНОВИДНОСТИ МНОГОЗНАЧНЫХ ОТОБРАЖЕНИЙ ПО
СУДЗУКИ В b -МЕТРИЧЕСКИХ ПРОСТРАНСТВАХ

Муджахид Абас^а, Басит Али^б, Талат Назир^в, Небойша М. Дедович^г,
Бандар Бин-Мохсин^д, Стоян Н. Раденович^е

^а Правительственный колледж в Лахоре - Университет, кафедра математики, г. Лахор, Исламская Республика Пакистан;
Преторийский университет, кафедра математики и прикладной математики, г. Претория, Южно-Африканская Республика

^б Университет менеджмента и технологий, кафедра математики, г. Лахор, Исламская Республика Пакистан

^в Университет COMSATS в Исламабаде, кафедра математики, Кампус в г. Абботтабад, Исламская Республика Пакистан;
Университет Южной Африки, кафедра математических наук, Научный кампус, г. Йоханнесбург, Южно-Африканская Республика

^г Нови-Садский университет, Сельскохозяйственный факультет, Департамент сельскохозяйственного машиностроения, г. Нови-Сад, Республика Сербия, **корреспондент**

^д Университет им. Короля Сауда, Колледж физико-математических наук, кафедра математики, Эр-Рияд, Королевство Саудовская Аравия

^е Белградский университет, Машиностроительный факультет, г. Белград, Республика Сербия

РУБРИКА ГРНТИ: 27.00.00 МАТЕМАТИКА;

27.25.17 Метрическая теория функций,

27.39.27 Нелинейный функциональный анализ

ВИД СТАТЬИ: оригинальная научная статья

Резюме:

Введение/цель: В данной статье представлены совпадения и общие неподвижные точки многозначного отображения типа Судзуки в b -метрических пространствах.

Методы: Обсуждаются предельные свойства, корректность и устойчивость решений задач неподвижной точкой таких отображений по методу Улама-Хайерса.

Результаты: Получена верхняя граница расстояния Хаусдорфа между неподвижными точками множеств. В качестве доказательства полученных результатов, в статье приведено несколько примеров.

Выводы: Применение полученных результатов доказывает существование дифференциальных включений.

Ключевые слова: b -метрические пространства, многозначные отображения, неподвижная точка и задачи, Улам-Хайерс стабильность, начальная задача.

РЕШЕЊА И УЛАМ-ХИЕРОВА СТАБИЛНОСТ ДИФЕРЕНЦИЈАЛНИХ ИНКЛУЗИЈА, УКЉУЧУЈУЋИ СУЗУКИЈЕВЕ ВРСТЕ ВИШЕЗНАЧНОГ ПРЕСЛИКАВАЊА НА b -МЕТРИЧКИМ ПРОСТОРИМА

Муџахид Абас^а, Басит Али^б, Талат Назир^в, Небојша М. Дедовић^г,
Бандар Бин-Мохсин^д, Стојан Н. Раденовић^ђ

^а Владин колеџ – Универзитет, Одсек математике,
Лахоре, Исламска Република Пакистан;
Универзитет у Преторији, Одсек за математику и примењену
математику, Преторија, Јужноафричка Република

^б Универзитет за менаџмент и технологију, Одсек за математику,
Лахоре, Исламска Република Пакистан

^в Универзитет COMSATS у Исламабаду, Одсек за математику,
Кампус Аботабад, Исламска Република Пакистан;
Универзитет Јужне Африке, Одсек за математичке науке,
Научни кампус, Јоханесбург, Јужноафричка Република

^г Универзитет у Новом Саду, Пољопривредни факултет,
Департман за пољопривреду технику,
Нови Сад, Република Србија, **аутор за преписку**

^д Универзитет краља Сауда, Колеџ природно-математичких наука,
Одсек за математику, Ријад, Краљевина Саудијска Арабија,

^ђ Универзитет у Београду, Машински факултет,
Београд, Република Србија

ОБЛАСТ: математика

ВРСТА ЧЛАНКА: оригинални научни рад

Сажетак:

Увод/циљ: У раду су представљене коинцидентне и заједничке фиксне тачке Сузукијеве врсте вишезначног пресликавања на b -метричким просторима.

Метод: Анализирана су гранична својства, добра постављеност и Улам-Хиерова стабилност решења за фиксни проблем вишезначних пресликавања.

Резултати: Добијена је горња граница Хаусдорфовог растојања између фиксних тачака скупова. Наведени су примери који подржавају добијене резултате.

Закључак: Применом представљених резултата установљена је егзистенција диференцијалне инклузије.

Кључне речи: b -метрички простори, вишезначно пресликавање, фиксна тачка и проблеми, Улам-Хиерова стабилност, почетни проблем.

Paper received on / Дата получения работы / Датум пријема чланка: 25.05.2020.
Manuscript corrections submitted on / Дата получения исправленной версии работы /
Датум достављања исправки рукописа: 03.06.2020.
Paper accepted for publishing on / Дата окончательного согласования работы / Датум
коначног прихватања чланка за објављивање: 05.06.2020.

© 2020 The Authors. Published by Vojnotehnički glasnik / Military Technical Courier
(www.vtg.mod.gov.rs, втг.мо.упр.срб). This article is an open access article distributed under the
terms and conditions of the Creative Commons Attribution license
(<http://creativecommons.org/licenses/by/3.0/rs/>).

© 2020 Авторы. Опубликовано в «Военно-технический вестник / Vojnotehnički glasnik / Military
Technical Courier» (www.vtg.mod.gov.rs, втг.мо.упр.срб). Данная статья в открытом доступе и
распространяется в соответствии с лицензией «Creative Commons»
(<http://creativecommons.org/licenses/by/3.0/rs/>).

© 2020 Аутори. Објавио Војнотехнички гласник / Vojnotehnički glasnik / Military Technical Courier
(www.vtg.mod.gov.rs, втг.мо.упр.срб). Ово је чланак отвореног приступа и дистрибуира се у
складу са Creative Commons licencom (<http://creativecommons.org/licenses/by/3.0/rs/>).

