



Darwiniana

ISSN: 0011-6793

sdenham@darwin.edu.ar

Instituto de Botánica Darwinion

Argentina

Amarilla, Leonardo D.; Chiapella, Jorge O.; Nagahama, Nicolás; Anton, Ana M.
INCLUSION OF DASYOCHLOA IN THE AMPHITROPICAL GENUS MUNROA (POACEAE,
CHLORIDOIDEAE) BASED ON MORPHOLOGICAL EVIDENCE

Darwiniana, vol. 1, núm. 2, julio-diciembre, 2013, pp. 241-252

Instituto de Botánica Darwinion

Buenos Aires, Argentina

Available in: <http://www.redalyc.org/articulo.oa?id=66929156008>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



INCLUSION OF *DASYOCHLOA* IN THE AMPHITROPICAL GENUS *MUNROA* (POACEAE, CHLORIDOIDEAE) BASED ON MORPHOLOGICAL EVIDENCE

Leonardo D. Amarilla, Jorge O. Chiapella, Nicolás Nagahama & Ana M. Anton

Instituto Multidisciplinario de Biología Vegetal (CONICET), Universidad Nacional de Córdoba Vélez Sarsfield 1611 - X5016GCA, Casilla de Correo 495, 5000 Córdoba, Córdoba, Argentina; amarillaleonardo@conicet.gov.ar (author for correspondence).

Abstract. Amarilla, L. D.; J. O. Chiapella, N. Nagahama & A. M. Anton. 2013. Inclusion of *Dasyochloa* in the amphitropical genus *Munroa* (Poaceae, Chloridoideae) based on morphological evidence. *Darwiniana*, nueva serie 1(2): 241-252.

Munroa is a morphologically heterogeneous American genus of grasses, including five species with disjunct distribution in North and South America. *Dasyochloa* is a monotypic grass genus, common in western North America, with a controversial taxonomic position, because it has been alternatively included in *Erioneuron*, *Koeleria*, *Triodia*, and *Uralespis*. We evaluated the relationship of *Dasyochloa* with *Munroa* using a phylogenetic analysis of qualitative and quantitative morphological data. *Dasyochloa* was not supported as an independent genus from *Munroa*, as *Dasyochloa* and *Munroa* species formed a monophyletic group defined by seven synapomorphies. On this basis, we transfer *Dasyochloa* to *Munroa*, and propose the new combination *M. pulchella* together with the emended diagnosis of *Munroa*.

Keywords. America; amphitropical distribution; Chloridoideae; *Dasyochloa*; *Munroa*; phylogeny; taxonomy.

Resumen. Amarilla, L. D.; J. O. Chiapella, N. Nagahama & A. M. Anton. 2013. Inclusión de *Dasyochloa* en el género anfítropical *Munroa* (Poaceae, Chloridoideae) sobre la base de evidencia morfológica. *Darwiniana*, nueva serie 1(2): 241-252.

Munroa es un género Americano morfológicamente heterogéneo, con cinco especies y distribución disyunta en América del Norte y del Sur. *Dasyochloa* es un género monotípico, común en el oeste de Norteamérica, con una historia taxonómica compleja porque ha sido incluido alternativamente en *Erioneuron*, *Koeleria*, *Triodia* y *Uralespis*. En este trabajo evaluamos las relaciones de *Dasyochloa* y *Munroa* con base en un análisis filogenético con caracteres morfológicos cualitativos y cuantitativos. *Dasyochloa* no resultó independiente de *Munroa*, ya que ambos géneros formaron un grupo monofilético definido por siete sinapomorfías. Sobre esta base, transferimos *Dasyochloa* a *Munroa* y proponemos la nueva combinación *M. pulchella*, junto con una enmienda en la diagnosis de *Munroa*.

Palabras clave. América; Chloridoideae; *Dasyochloa*; distribución anfítropical; *Munroa*; filogenia; taxonomía.

INTRODUCTION

The monotypic genus *Dasyochloa* Willd. ex Rydb. was founded by Rydberg (1906) based on *Triodia pulchella* Kunth [= *D. pulchella* (Kunth) Willd. ex Rydb.]. This species occurs in North America, in arid and semiarid regions of the western USA and northern Mexico. The identity of

Dasyochloa as an independent genus has been questioned several times since its establishment, and the taxon has been placed under several genera, including *Triodia* R. Br. (Kunth, 1815), *Koeleria* Pers. (Sprengel, [1824] 1825), *Uralespis* Nutt. (Kunth, 1829) and *Erioneuron* Nash (Tateoka, 1961) (for a complete explanation of the complicated taxonomical history of *Dasyochloa*, see

Valdés-Reyna & Hatch, 1997). Sánchez (1983) and Valdés-Reyna & Hatch (1995), using anatomical data, considered *Dasyochloa* as an independent monotypic genus separated from *Erioneuron*. Valdés-Reyna & Hatch (1997), using morphological data and protein electrophoresis accepted a monotypic *Dasyochloa* with *D. pulchella* as its single species.

Munroa Torr. was described by Torrey (1857) based on *Crypsis squarrosa* Nutt. [= *M. squarrosa* (Nutt.) Torr.], and Philippi (1870) described the first South American taxon, *M. mendocina* Phil. Later, the addition of *M. decumbens* Phil., *M. andina* Phil. and *M. argentina* Griseb., raised the number of accepted taxa to five. Anton & Hunziker (1978) revised the genus and established three sections based on the articulation of the rachilla: sect. *Munroa* (including only *M. squarrosa*), sect. *Hemimunroa* (Parodi) Anton & Hunz. (including *M. andina* and *M. decumbens*), and sect. *Apelytron* (Parodi) Anton & Hunz. (including *M. argentina* and *M. mendocina*).

Munroa occurs in two disjunct regions: in western North America (from Alberta and Saskatchewan to Sonora), and western South America, including mountain areas ranging from southern Peru through Bolivia and Chile and north-western Argentina (Anton & Hunziker, 1978). *Munroa squarrosa* occurs from southern Canada to northern Mexico and in its southern range has a nearly complete overlapping distribution with *D. pulchella* (Valdés-Reyna, 2003a); the other taxa, *M. andina*, *M. decumbens*, *M. mendocina* and *M. argentina*, inhabit arid and semi-arid regions above 1,200 m a. s. l. in South America (Anton & Hunziker, 1978).

The first hypothesis of a close relationship between *Dasyochloa* and *Munroa* was made by Parodi (1934), who noted similarities in several characters: spikelets in a short raceme not exceeding the blades of the fascicle, short leaves sheathed, ligules reduced to a fringe of short hairs, subsessile spikelets included in distal leaf sheaths, caryopses laterally compressed, papillate stigmas, and the special branching type, which consists of stems with the first internode well developed in length, and the successive internodes shorter and of variable length (see also Anton & Hunziker, 1978; Hunziker & Anton, 1979). Sánchez (1983, 1984) also found similarities in the leaf anatomy of *Dasy-*

ochloa and *Munroa*, and a recent molecular phylogenetic study of the Chloridoideae (Peterson et al., 2010a) showed *D. pulchella* as a sister taxon of *Munroa* in a well-supported clade.

Considering the sister relationship of *Dasyochloa* and *Munroa* in previous molecular phylogenies, their morphological affinities, and the overlapping distribution of *M. squarrosa* and *D. pulchella*, we hypothesize that *Dasyochloa* and *Munroa* form a single taxon. To test this hypothesis, we performed a cladistic analysis (parsimony) using morphological characters (discrete and continuous) to evaluate the phylogenetic relationship of *Dasyochloa* and *Munroa*.

MATERIAL AND METHODS

Plant material

Collections were made between 2010–2012, covering western Argentina and Bolivia, northern Chile and north western Mexico. Voucher specimens were deposited at CORD and XAL (Thiers, 2013). Field-collected specimens were identified using the keys and descriptions published by Anton & Hunziker (1978) and Valdés-Reyna & Hatch (1997). Plants used in the analysis were in full bloom, and were selected from different localities to cover the geographic range (Fig. 1) and represent the morphological variability of each species. Additional herbarium specimens were analysed from BAA, BAB, CORD, LIL, LPB, XAL, and UTC (Appendix 1).

Phylogenetic analysis

The data matrix was generated from direct observations of herbarium vouchers and from the literature (Parodi, 1934; Cáceres, 1950; Anton & Hunziker, 1978; Sánchez, 1983, 1984; Valdés-Reyna & Hatch, 1997; Allred, 2003; Peterson, 2003; Valdés-Reyna, 2003a, b, c; Stephenson & Saarela, 2007; Nicora & Rúgolo de Agrasar, 2009) and includes 11 taxa: all species of *Munroa* and *Dasyochloa pulchella* as ingroup, and *Erioneuron pilosum* (Buckley) Nash, *Blepharidachne bigelovii* (S. Watson) Hack., *Muhlenbergia montana* (Nutt.) Hitchc., and *Aristida adscensionis* L. as outgroups. The tree was rooted in *Brachyelytrum erectum* (Schreb.) P. Beauv. The outgroups and the root




Fig. 1. Geographic localities of the collections of *Munroa* and *Dasyochloa* (= *M. pulchella*) species.

were selected according to the phylogeny of Peterson et al. (2010).

Phylogenetic analysis and character optimization were implemented on a data matrix of 63 vegetative and reproductive characters, 20 were quantitative and 43 were qualitative. The quantitative characters were expressed as intervals of minimum and maximum values, and qualitative characters were binary or multistate. Details of the character coding and the resulting data matrix are shown in Appendix 2 and Appendix 3.

Phylogenetic analysis under maximum parsimony criteria was carried out using TNT v. 1.1 (Goloboff et al., 2008), and characters were treated as non-additive and equal-weighted. This analysis was performed with 1,000 random addition sequences (RAS) and posterior exchange branches with tree bisection-reconnection (TBR) algorithm, saving 10 trees per replicate. Support and stability was estimated by Jackknife method (Farris et al., 1996), resampling 1,000 times and expressing support values as absolute frequencies (Jk) and GC values (present group/contradicted group) (Goloboff et al., 2003) following the methodology employed by De Gennaro & Scatagliini (2012). Synapomorphies were mapped only in *Dasyochloa* and *Munroa* clade. Retention index (RI) and consistency index (CI) were calculated according to Farris (1989) and Kluge & Farris (1969) respectively.

The electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Code of Nomenclature for algae, fungi, and plants, and hence the new names contained in the electronic version are effectively published under the Code from the electronic edition alone. The online version of this work is archived and available from Instituto de Botánica Darwinion and the digital repositories cited in <http://www.ojs.darwin.edu.ar/index.php/darwiniana/about/editorialPolicies#custom-0>.

RESULTS

Phylogenetic analysis generated one most parsimonious tree (72 steps, CI = 0.6; RI = 0.7). *Dasyochloa pulchella* and *Munroa* formed a monophyletic and supported clade (Jk = 92, GC = 90) and *D. pulchella* is sister to *Munroa* (Fig. 2). *Munroa*

species formed a clade (Jk = 88, GC = 86) which is divided into two clades and one subclade: *M. andina*-*M. decumbens* clade (Jk = 75, GC = 67), *M. squarrosa*-*M. mendocina*-*M. argentina* clade (Jk = 65, GC = 53), and *M. mendocina*-*M. argentina* subclade (Jk < 50, GC = 23).

Seven synapomorphies were detected for the *Dasyochloa*-*Munroa* clade and two synapomorphies for the *Munroa* clade (Fig. 2). Synapomorphies of the *Dasyochloa*-*Munroa* clade are: fertile branch length (4–15 cm), stolon length (3–9 cm), special branching type (present), raceme concealed in the sheath (present), bicellular microhairs with distal cell hemispherical (present), stigmas papillose (present), and caryopsis dorsiventrally compressed (present). Synapomorphies of the *Munroa* clade are: ligule length (0.4–2 mm) and strand of hairs on the lemma of terminal spikelet (present).

DISCUSSION

Our results confirm the phylogenetic affinity between *Dasyochloa* and *Munroa*, with a higher support and number of synapomorphies for the *Dasyochloa*-*Munroa* clade than those for the *Munroa* subclade. The synapomorphies of the *Dasyochloa*-*Munroa* clade: branching type, raceme concealed in the sheath, stigmas papillose and caryopses dorsiventrally compressed, had been already recognized by Parodi (1934) as diagnostic characters defining *Munroa*. The close relationship between *Dasyochloa* and *Munroa* is also supported by previous results obtained by Columbus et al. (2007), where *M. squarrosa* and *D. pulchella* formed a clade, and by Peterson et al. (2010: 590, Fig. 3; 2010b: 1544, Fig. 3) where *D. pulchella* was sister to *Munroa* (*M. argentina* and *M. andina*). Based on previous and present evidence, we synonymized *Dasyochloa* to *Munroa*.

TAXONOMIC TREATMENT

Munroa Torr., Pacif. Railr. Rep. 4 (5, no. 4) [Whipple]: 158. 1856 [Sep 1857], nom. et orth. cons., **emend.** Amarilla. TYPE SPECIES: *Munroa squarrosa* (Nutt.) Torr. (= *Crypsis squarrosa* Nutt.).




Fig. 2. Single most parsimonious tree based on cladistic analysis of 63 quantitative and qualitative morphological characters. Numbers below the branches indicate Jk and GC values respectively. Synapomorphies of *Dasyochloa*-*Munroa* clade and *Munroa* clade are shown on the left (character number, name, and state).

Dasyochloa Willd. ex Rydb., Agric. Exp. Sta. Agric. Coll. Colorado Bull. 100: 18, 37. 1906, **syn. nov.** TYPE SPECIES: *Dasyochloa pulchella* (Kunth) Willd. ex Rydb. (= *Triodia pulchella* Kunth).

Annual or perennial, gynomonoeious or monoeious; culms tufted, branched, often decumbent and stoloniferous, rooting at the lower nodes, forming short spur shoots at the end of long internodes. Basal sheaths shorter than the internodes, open, margins membranous. Ligules a fringe of hairs. Blades short, flat or conduplicate, usually covered with a cottony secretion. Prophylls developed, equal to or longer than the subtending sheath and with two nerves that end into awns. Inflorescences included in leafy fascicles with 1-4 or more sub-sessile spikelets, light green or purple tinged. Spikelets sessile or shortly pedicellate, dorsiventrally or laterally compressed, sometimes dimorphic in the same inflorescence in terms of the articulation of the rachilla, number, shape, texture and hairiness of glumes and sexuality. Glumes 1-2 or absent, equal or sub equal, 1-veined, membranous or, coriaceous, thick, often curved, shorter, equal or longer than the spikelet. Florets 2-10, rachilla disarticulating or not. Lemma 3-5-7-veined, membranous or coriaceous, 1-3 awned or mucronate, with awns 0.5-4.5 mm long, 2-4-lobate, lobes separated by awns or mucros, with tufts of hairs or

completely pilose on the base. Palea shorter than the lemma, 2-keeled, wrapping the flower, lanceolate, spatulate or lyre-shaped. Lodicules 2 or absent, membranous when present, free or fused to the margins of the palea. Stamens 2 or 3, exserted, sometimes reduced to staminodes. Ovaries with 2 long styles, and stigmas papillose. Caryopsis dorsiventrally compressed, with the scutellum ca. half of its length; hilum punctiform.

Distribution and habitat. American genus with disjunct distribution in North and South America: *Munroa squarrosa* and *M. pulchella* grow in North America while the remaining four species are from South America, distributed in Argentina, Bolivia, Chile and southern Peru. *Munroa* species inhabit arid and semi-arid regions, in open areas of plains and mountains at middle altitude with dry and sandy-stony modified soils.

Munroa pulchella* (Kunth) Amarilla, **comb. nov.*
Triodia pulchella Kunth, Nov. Gen. Sp.1: 155-156. 1815 [1816]. *Koeleria pulchella* (Kunth) Spreng., Syst. Veg. (ed. 16) [Sprengel] 1: 332. 1824 [dated 1825; publ. in late 1824]. *Uralep-is pulchella* (Kunth) Kunth, Révis. Gramin. 1: 108. 1829. *Tricuspis pulchella* (Kunth) Torr., Pacif. Railr. Rep. 4 (5, no. 4) [Whipple]: 156.

1857. *Dasyochloa pulchella* (Kunth) Willd. ex Rydb., Colorado Exp. Sta. Bull. 100: 18, 37. 1906. *Tridens pulchellus* (Kunth) Hitchc., Fl. Calif. [Jepson] 1: 141. 1912. *Erioneuron pulchellum* (Kunth) Tateoka, Amer. J. Bot. 48: 572. 1961. TYPE: Mexico. In subfrigidis, siccis, aprics reginii Mexicani inter Guanaxuati, Mina de Belgrado et Cubilete, *F. W. H. A. Humboldt & A. J. A. Bonpland s.n.* [holotype P cb 00669430; isotypes B-W 2046, US 91469! (fragment)].

Sieglingia pulchella (Kunth) Kuntze var. *parviflora* Vasey ex L. H. Dewey, Contr. U.S. Natl. Herb. 2(3): 538. 1894. TYPE: United States of America, Western Texas, *V. Havard* 7 (holotype probably US 908213 cb 00141433).

References. A detailed taxonomic description can be found in Valdés-Reyna & Hatch (1997).

Distribution and habitat. *Munroa pulchella* grows in western United States of America (Arizona, California, Nevada, Utah, Colorado, New Mexico, Texas) and north-western Mexico (Aguascalientes, north of Baja California, Chihuahua, Coahuila, Durango, Guanajuato, Hidalgo, Jalisco, Querétaro, México, Nuevo León, San Luis Potosí, Sonora, Tamaulipas and Zacatecas).

ACKNOWLEDGEMENTS

We are grateful to the “Consejo Nacional de Investigaciones Científicas y Técnicas” (CONICET), and the Myndel Botanica Foundation for financial support. We thank the curator and technical staff of BAA, BAB, CORD, LIL, LPB, XAL and UTC herbaria, and Dr. Joseph C. Kuhl (University of Idaho), R. Pozner (IBODA, CONICET-ANCEFN), F. Zuloaga (IBODA, CONICET-ANCEFN) and anonymous reviewers for useful suggestions to improve the manuscript.

BIBLIOGRAPHY

Allred, K. W. 2003. *Aristida*, in M. E. Barkworth, K. M. Capels, S. Long & M. B. Piep (eds.), *Flora of North America* 25: *Magnoliophyta: Commelinidae (in part): Poaceae*, part 2, pp. 315-342. New York: Oxford University Press.

Anton, A. M. & A. T. Hunziker. 1978. El género *Munroa* (Poaceae): Sinopsis morfológica y taxonómica. *Boletín de la Academia Nacional de Ciencias (Córdoba)* 52: 229-252.

Cáceres, M. R. 1950. Los caracteres anatómicos foliares de *Munroa mendocina* y *Blepharidachne benthamiana*. *Revista Argentina de Agronomía* 17: 233-240.

Columbus, J. T.; R. Cerros-Tlatilpa; M. S. Kinney, M. E. Siqueiros-Delgado; H. L. Bell; M. P. Griffith & N. F. Refulio-Rodriguez. 2007. Phylogenetics of Chloridoideae (Gramineae): a preliminary study based on nuclear ribosomal internal transcribed spacer and chloroplast trnL-F sequences. *Aliso* 23: 565-579.

De Gennaro, D. & M. A. Scataglini. 2012. Análisis filogenético del género *Sacciolepis* (Poaceae, Panicoideae, Paniceae) con caracteres morfológicos. *Darwiniana* 50: 81-96.

Farris, J. S. 1989. The retention index and the rescaled consistency index. *Cladistics* 5: 417-419. DOI: <http://dx.doi.org/10.1111/j.1096-0031.1989.tb00573.x>.

Farris, J. S.; V. A. Albert, M. Källersjö, A. Lipscomb & A. G. Kluge. 1996. Parsimony jackknifing outperforms neighbor-joining. *Cladistics* 12: 99-124. DOI: <http://dx.doi.org/10.1111/j.1096-0031.1996.tb00196.x>.

Goloboff, P. A.; J. S. Farris, M. Källersjö, B. Oxelman, M. J. Ramírez & C. A. Szumik. 2003. Improvements to resampling measures of group support. *Cladistics* 19: 324-332. DOI: <http://dx.doi.org/10.1111/j.1096-0031.2003.tb00376.x>.

Goloboff, P. A.; J. S. Farris & K. C. Nixon. 2008. TNT a free program for phylogenetic analysis. *Cladistics* 24: 774-786. DOI: <http://dx.doi.org/10.1111/j.1096-0031.2008.00217.x>.

Hunziker, A. T. & A. M. Anton. 1979. A synoptical revision of *Blepharidachne* (Poaceae). *Brittonia* 31(4): 446-453. <http://dx.doi.org/10.2307/2805993>.

Kluge, A. G. & J. S. Farris. 1969. Quantitative phyletics and the evolution of anurans. *Systematic Biology* 18(1): 1-32. DOI: <http://dx.doi.org/10.1093/sysbio/18.1.1>.

Kunth, K. S. 1815. *Triodia*, in F. W. H. A. von Humboldt, A. J. A. Bonpland & K. S. Kunth (eds.), *Nova Genera et Species Plantarum*, vol. 1, pp. 155-156. París: Librariæ Græco-Latino-Germanicæ.

Kunth, K. S. 1829. *Uralepis*, in F. W. H. A. von Humboldt, A. J. A. Bonpland & K. S. Kunth (eds.), *Révision des Graminées*, vol. 6, pp. 108. París: Botanique, Gide fils, rue Saint-Marc-Feydeau.

Nicora, E. G. & Z. E. Rúgolo de Agrasar. 2009. *Munroa*, in R. Kiesling, *Flora de San Juan*, vol. 4, pp. 218-220. San Juan: Fundación Universidad Nacional de San Juan.

Parodi, L. R. 1934. Contribuciones al estudio de las Gramíneas del género *Munroa*. *Revista del Museo de la Plata* 34: 171-193.

Peterson, P. M. 2003. *Muhlenbergia*, in M. E. Barkworth, K. M. Capels, S. Long & M. B. Piep (eds.), *Flora of North Ame-*

- rica 25: *Magnoliophyta: Commelinidae (in part): Poaceae*, part 2, pp. 145-200. New York: Oxford University Press.
- Peterson, P. M.; K. Romaschenko & G. Johnson. 2010. A classification of the Chloridoideae (Poaceae) based on multi-gene phylogenetic trees. *Molecular Phylogenetics and Evolution* 55: 580-598. DOI: <http://dx.doi.org/10.1016/j.ympev.2010.01.018>.
- Philippi, R. A. 1870. Sertum Mendocinum alternum; ó sea, catálogo de las plantas recogidas cerca de Mendoza y en los caminos que conducen de Chile a esa ciudad. *Anales de la Universidad de Chile* 36: 159-212.
- Rydberg, P. A. 1906. Flora of Colorado. *Colorado Agricultural Experiment Station Bulletin* 100: 18-37.
- Sánchez, E. 1983. *Dasyochloa* Willdenow ex Rydberg (Poaceae), género monotípico de Norteamérica. *Lilloa* 36: 131-138.
- Sánchez, E. 1984. Estudios anatómicos en el género *Munroa* (Poaceae: Chloridoideae: Eragrosidae). *Darwiniana* 25: 43-57.
- Sprengel, K. P. J. [1824] 1825. *Kölera*, in K. P. J. Sprengel, *Systema vegetabilium, editio 16*, part 1, p. 332. Gottingen: Librariae Dieterichianae.
- Stephenson, S. N. & J. M. Saarela. 2007. *Brachyelytrum*, in M. E. Barkworth, K. M. Capels, S. Long, L. K. Anderton & M. B. Piep (eds.), *Flora of North America 25: Magnoliophyta: Commelinidae (in part): Poaceae*, vol. 1, pp. 59-61. New York: Oxford University Press.
- Tateoka, T. 1961. A biosystematics Study of *Tridens* (Gramineae). *American Journal of Botany*. 48: 572.
- Thiers, B. [continuously updated, accessed 2013] Index Herbariorum: a global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium, <http://sweetgum.nybg.org/ih>
- Torrey, J. 1857. Report on the Botany of the expedition: Description of the general botanical collections, in War Department (ed.), *Explorations and surveys for a railroad route from the Mississippi River to the Pacific Ocean [A. W. Whipple]*, vol. 4, part 5, pp. 5[61]-105[161]. Washington, D. C.: Government Printing Office.
- Valdés-Reyna, J. 2003a. *Dasyochloa*, in M. E. Barkworth, K. M. Capels, S. Long & M. B. Piep (eds.), *Flora of North America 25: Magnoliophyta: Commelinidae (in part): Poaceae*, part 2, pp. 45-47. New York: Oxford University Press.
- Valdés-Reyna, J. 2003b. *Erioneuron*, in M. E. Barkworth, K. M. Capels, S. Long & M. B. Piep (eds.), *Flora of North America 25: Magnoliophyta: Commelinidae (in part): Poaceae*, part 2, pp. 44-45. New York: Oxford University Press.
- Valdés-Reyna, J. 2003c. *Munroa*, in M. E. Barkworth, K. M. Capels, S. Long & M. B. Piep (eds.), *Flora of North America 25: Magnoliophyta: Commelinidae (in part): Poaceae*, part 2, p. 51. New York: Oxford University Press.
- Valdés-Reyna, J. & S. L. Hatch. 1995. Anatomical study of *Erioneuron* and *Dasyochloa* (Poaceae: Chloridoideae: Eragrosidae) in North America. *Sida* 16: 413-427.
- Valdés-Reyna, J. & S. L. Hatch. 1997. A revision of *Erioneuron* and *Dasyochloa* (Poaceae: Eragrosidae). *Sida* 17: 645-666.

APPENDIX 1

Representative material examined

Munroa andina Phil.

ARGENTINA. **Catamarca**. Depto. Belén, “El Potrero” Laguna Blanca, 3500 m, 25-II-1934, *Peirano* 299 (LIL 29802). **Jujuy**. Depto. Tilcara, Huacalera, Pampa Corral, 4000 m, 2-I-1955, *Cabrera* 12108 (BAA); Abra Pampa, ruta 71 camino a Cochinoca, 19-II-2011, *Amarilla & Moreno* 17 (CORD). **La Rioja**. Depto. Vinchina, Valle del Río Bonete cercanías de Las Cuevitas, 4-IV-1950, *Hunziker & Caso* 4189 (BAB). **Salta**. Depto. San Antonio de los Cobres, *Cabrera* 8699 (LP). **Tucumán**: Depto. Tañi del Valle, Lara, 3200 m, 16-II-1912, *Rodríguez* 335 (LIL 11190).

CHILE. **I Región Tarapacá**. Prov. El Tamarugal, 6 km SW of Colchane on HWY towards Huará, 27-III-2001, *Peterson* 15666 (CORD). **II Región Antofagasta**. Prov. Antofagasta, km 29,5 East from San Pedro de Atacama, 16-III-2001, *Peterson* 15504 (CORD).

BOLIVIA. **Oruro**. Prov. Ladislao Cabrera, 11-III-2008, *Acho* 53 (LPB). **Potosí**. Prov. Daniel Campos, 15-III-2008, *Acho* 97 (LPB). **La Paz**. Prov. Gualberto Villarroel, 17-II-1992, *Beck* 21006A (LPB).

Munroa argentina Griseb.

ARGENTINA. **Catamarca**. Depto. Andalgalá, Las Minas, 6-III-1916, *Jørgensen* 1696 p.p. (BAA, LIL 45082). Depto. Tinogasta, Vallecito a Agua Negra, 2400 m, 7-II-1930, *Schreiter* 6308 (LIL 1992). **Jujuy**. Depto. Humahuaca, Humahuaca, 3050 m, 16-II-1931, *Parodi* 9703 (BAA). **La Rioja**. Depto. Famatina, Las Gredas, 1700 m, II-1907, *Kurtz* 14387 (CORD). **Mendoza**. Depto. Las Heras, entre Uspallata y Ciénagas de Yalguaraz, 16-XII-1956, *Ruiz Leal & Roig* 18372 (BAA). **Salta**. Depto. Ca-

fayate, Cafayate, Ruta 40, lecho del río Chuscha, 28-III-2010, *Chiapella & Amarilla* 2712 (CORD). Depto. Rosario de Lerma, Puerta Tastil, 24-II-1972, *Cabrera et al.* 22421 (BAA). **San Juan**. Depto. Jáchal, al norte de la escuela Entre Ríos, 22-III-2010, *Chiapella & Amarilla* 2596 (CORD). **Tucumán**. Depto. Tañi del Valle, río Tañi del Valle entre el Molle y Carapunco, 28-III-2010, *Chiapella & Amarilla* 2728 (CORD); Amaicha del Valle, Colalao del Valle, ruta 307 alrededores del museo Pachamama, 16-III-2011, *Amarilla* 11 (CORD).

BOLIVIA. **Chuquisaca**. Prov. Nor Sinti, 1 km al sur de Camargo, 26-II-1995, *Wood* 9498 (LPB). **Potosí**. Prov. Nor Chichas, 2-IV-1999, *López* 522 (LPB).

Munroa decumbens Phil.

ARGENTINA. **Catamarca**. Depto. Tinogasta, Entre Cazadero Grande y El Quemado, 3550 m, 21-III-1951, *Vervoorst* 3188 (BAA, LIL 364811). **Jujuy**. Depto. Rinconada, Mina Pirquitas, 4300 m, 1-III-1964, *Schwabe* 816 (BAA). **La Rioja**. Depto. Vinchina, Las Cortaderas, entre El Peñón y El Jagüel, 27-II-1879, *Hieronymus & Niederlein* 237 (CORD); Rosario Vera Peñaloza, Chepes, frente al cementerio, 20-III-2010, *Chiapella & Amarilla* 2576b (CORD). **Salta**. Depto. Los Andes, entre San Antonio y Cangrejillos, 3850 m, 25-II-1972, *Cabrera et al.* 22432 (BAA). **San Juan**. Depto. Iglesia, Punta del Cerro de Los Médanos, 3040 m, 4-III-1962, *Ruiz Leal* 22154 (BAA). **Tucumán**. Depto. Tañi del Valle, El Molle, 3000 m, 3-II-1933, *Parodi* 10998 (BAA).

BOLIVIA. **Oruro**. Prov. Avaroa. S. Pagador, 8-III-1993, *Peterson* 12764 (LPB). **Potosí**. Prov. Quijarro, 26-III-1992, *Peterson* 13092 (LPB).

PERU. **Arequipa**. Prov. Arequipa, 28-III-1949, *Vargas* 1981 (LIL). *Munroa mendocina* Phil.

ARGENTINA. **Catamarca**. Depto. Belén, Belén, lecho del río Belén, 25-III-2010, *Chiapella & Amarilla* 2656 (CORD). Depto. Tinogasta, Tinogasta, ruta provincial 3, a 1 km del límite con La Rioja, 24-III-2010, *Chiapella & Amarilla* 2618

(CORD). **La Rioja**. Depto. Chilecito, ruta 40 entre Soñogasta y Villa Unión, 11-IV-2012, *Nagahama* 128 (CORD); General Lavalle, ruta 40 entre Soñogasta y Villa Unión, 11-IV-2012, *Nagahama* 129 (CORD). **Mendoza**. Depto. San Rafael, Los Pocitos, desde escuela Parlamentos hacia Sosneado (ruta 144), 9-III-2011, *Fortunato* 10001 (BAB). **San Juan**. Depto. Jáchal, Ruta 40 entre Jáchal y Tucunuco, 8-III-2011, *Fortunato* 9960 (BAB).

Munroa pulchella (Kunth) Amarilla

MEXICO. **Coahuila**. Munic. Sacramento, Ruta 30 entre Sacramento y Cuatro Ciénagas, 7-IX-2011, *Angulo & Amarilla* 380 (XAL). **Chihuahua**. Munic. Ciudad Jiménez, Ruta 80D a 10 km de ruta 16, 10-IX-2011, *Angulo & Amarilla* 391 (XAL); Munic. Ciudad Jiménez, Ruta 45, llegando a Ciudad Jiménez, 10-IX-2011, *Angulo & Amarilla* 389 (XAL). **Durango**. Munic. Gómez Palacio, Ruta 45 hacia el norte, 40 km antes de Torreón, 6-IX-2011, *Angulo & Amarilla* 369 (XAL). **Tamaulipas**. Munic. Ciudad de Maíz, Ruta 101 llegando a la frontera con San Luis Potosí, 13-IX-2011, *Angulo & Amarilla* 400 (XAL).

Munroa squarrosa (Nutt.) Torr.

MEXICO. **Chihuahua**. Munic. Juan Aldama, Ruta 16 pasando Juan Aldama camino a Ojinaga, 10-IX-2011, *Angulo & Amarilla* 392 (XAL).

UNITED STATES OF AMERICA. **Colorado**. Nathrop co., Chaffe Country, 30-VIII-1913, *Hitchcock s.n.* (LIL 412388). **New Mexico**. Quay, 10 millas SE Logan, 13-VIII-1939, *E. Williams s.n.* (LIL 140306). **Texas**. Sine locus, 10-VII-1950, *Thornton s.n.* (LIL 388355). **Utah**. Kane co., Colorado plateau, 27-VIII-2008, *Fertig* 24313 (UTC); White Cliffs, 2-IX-2005, *Fertig* 22317 (UTC). **Wyoming**. Johnson co., Fork Powder River, 1901, *Leslie* 256 (CORD); Thermopolis co., 28-VII-1937, *Weatherwax* 2212 (LIL).

APPENDIX 2

Morphological quantitative (1—20) and qualitative (21—64) characters. Quantitative characters

were measured in centimetre (cm) and millimetre (mm). See matrix in Appendix 3.

Fertile branch length (1): cm. First internode length (2): cm. Stolon length (3): cm. Sheath length (4): cm. Blade length (5): cm. Blade width (6): cm. Ligule length (7): mm. Inflorescence length (8): cm. Inflorescence width (9): mm. Spikelet length (10): mm. First glume length (11): mm. Second glume length (12): mm. Lemma length (13): mm. Palea length (14): mm. Lemma central awn length (15): mm. Anther length (16): mm. Caryopsis length (17): mm. Caryopsis width (18): mm. Scutellum length (19): mm. Lodicule length (20): mm. Habit (21): perennial 0, annual 1. Rhizomes (22): absent 0, present 1. Special branching type (23): absent 0, present 1. Raceme concealed in the sheath (24): absent 0, present 1. Prophyll (25): absent 0, present 1. Sheaths with adaxial surface pilose (26): absent 0, present 1. Sheaths with hairs at the throat (27): absent 0, present 1. Blades with the adaxial surface pilose (28): absent 0, present 1. Ligules (29): absent 0, present 1. Ligule hairy (30): absent 0, present 1. Bicellular micro hairs in the abaxial epidermis of leaves (31): absent 0, present 1. Bicellular micro hairs with hemispherical distal cell in the abaxial epidermis of leaves (32): absent 0, present 1. Morphological dimorphism in spikelets (33): absent 0, present 1. Number of spikelets (34): less than four 1, more than four 0. Spikelets pedicellate or sub sessile (35): pedicellate 0, sub sessile 1. Number of florets per spikelets (36) one 0, more than one 1. Rhachilla disarticulating above the glumes (37): yes 0, no 1. Rhachilla disarticulating between the florets (38): yes 0, no 1. Number of glumes on lateral spikelets (39) zero 0, two 1. Glumes awns (40) without awn 0, with awn 1.

Glume relative length (41): equal to sub-equal 0, different 1. Number of glumes on terminal spikelets (42): two 0, one 1. Glumes fused or non-fused (43): non-fused 0, fused 1. Consistency of lemma on lateral spikelets (44): coriaceous 0, membranous 1, papyraceous 2. Consistency of lemma on terminal spikelets (45) coriaceous 0, membranous 1, papyraceous 2. Central awn of lemma (46): absent 0, present 1. Lateral awn of lemma (47): absent 0, present 1. Insertion of awns on the lemma (48): terminal 0, dorsal 1. Lemma veined (49): 5-veined 0, 3-veined 1, 7-veined 2. Lemma lobes (50): without lobes 0, 2-lobes 1, 3-lobes 2, 4-lobes 3. Strands of hairs on the lemma of lateral spikelets (51): absent 0, present 1. Strands of hairs on the lemma of terminal spikelets (52): absent 0, present 1. Lemma with adaxial surface pilose (53): absent 0, present 1. Lodicules (54): absent 0, present 1. Lodicules vascularization (55): absent 0, present 1. Lodicules fused or non-fused (56): non-fused 0, fused 1. Sex expression (57): monoclinal 0, gynomonocious 1, andromonocious 2. Number of stamens (58): three 0, two 1. Staminodes (59): absent 0, present 1. Stigma papillose (60): absent 0, present 1. Caryopsis dorsiventrally compressed (61): absent 0, present 1. Hilum form (62): punctuate 0, linear 1. Transversal cut of leaf (63): "U" form 0, "V" form 1.

APPENDIX 3

Data matrix used in the phylogenetic analysis. Numbers indicate the morphological characters listed in Appendix 2; ?, indicates missing data; "--", indicates non-applicable data.

	1	2	3	4	5	6	7	8	9	10
<i>M. squarrosa</i>	7.0-15	7.0-13	5.5-9.0	3-6	0.8-1.0	1.5-2	0.5-1	1-2.5	10-15	6-10
<i>M. argentina</i>	8.0-13	5.5-8.0	5.0-9.0	3-5	0.8-1.1	1-1.5	0.5-2	1-1.4	5-10	5-8
<i>M. mendocina</i>	6.5-18	4.0-8.0	4.0-9.0	3-7	0.7-1.1	1.5-2	0.5-0.8	1-1.5	10-13	8-10
<i>M. decumbens</i>	4.0-15	3.5-5.0	3.0-8.8	3-5	0.3-0.4	1.5-2	0.5-1	0.9-1.2	9-11	5.5-8
<i>M. andina</i>	4.0-9.5	2.5-5.0	3.5-7.0	4-7	0.4-0.8	1-1.7	0.4-0.8	0.9-1.2	8-11	6-7
<i>D. pulchella</i>	4.5-9.5	3.5-8.5	3.0-8.0	3-5	1.0-5.0	0.5-1	3-5	1.2-2.4	11-15	6-9.2
<i>E. pilosum</i>	10-30	4.0-9.0	-	3-5.5	1.5-5.0	1-1.5	2-3.5	1-2.0	10-15	6-12
<i>B. bigelovii</i>	9-20	5.5-8.5	-	3-5	1.0-4.5	1-1.5	0.5-0.9	1.5-3.0	10-15	5-7
<i>M. montana</i>	10-62	8.0-60	-	20-25	5.0-12	1-2.5	4-14	4-24	20-60	3-7
<i>A. adscensionis</i>	10-60	4.0-6.0	-	6-36	1.0-12	1-2.5	0.4-1	5-15	5-30	10-40
<i>B. erectum</i>	34-65	3.0-5.0	-	30-65	7.0-16	11-17	2-3.5	9-14	?	29-36

	11	12	13	14	15	16	17	18	19	20
<i>M. squarrosa</i>	3-4.2	2.5-4	2.5-3	3-4.5	0.8-1.2	1-1.5	1.2-1.5	1-1.1	0.3-0.5	0.2-0.3
<i>M. argentina</i>	4.8-5.2	4.7-5.2	2.5-5	3-4.5	0.8-1.3	1-1.5	1-1.6	0.9-1.3	0.6-1	0.2-0.3
<i>M. mendocina</i>	-	-	3.2-6	2.5-3.5	0.4-0.9	1-1.5	1.1-1.5	1-1.2	0.6-0.8	-
<i>M. decumbens</i>	4.5-6.5	8-10	2.9-3.8	2.5-3	0.8-1	0.7-0.9	1.4-1.7	0.8-1	0.6-0.8	-
<i>M. andina</i>	4-5.2	4.5-5.6	3-5	2-2.8	3-4.5	0.4-0.7	1-1.3	0.5-1	0.7-0.8	0.2-0.2
<i>D. pulchella</i>	6-8.5	6.8-9	3-5.5	2.3-3.5	2-4	0.2-0.4	1-1.5	0.9-1	0.5-0.7	0.2-0.3
<i>E. pilosum</i>	4-7	4-7	3-6	3-6	0.5-2.5	0.3-1	1-1.5	0.8-1.1	0.5-0.7	0.2-0.3
<i>B. bigelovii</i>	5.0-6	5.5-6.0	5-6	5-6	2.5-3	0.5-2.3	1.5-2.0	1-1.5	0.9-1.3	-
<i>M. montana</i>	1-3.2	1.5-4	2.8-3.5	2.8-3.5	6-25	1.5-2.3	1.8-2	?	0.5-1.3	0.1-0.2
<i>A. adscensionis</i>	4-8	3-11	6-9	6-9	7-15	0.3-0.7	4-7	1.2-1.8	?	?
<i>B. erectum</i>	0.3-1.1	0.9-7	9-13	7-12	13-17	3.5-6	5.5-7.5	1.5-2	?	?

APPENDIX 3. Continued.

	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
<i>M. squarrosa</i>	1	0	1	1	1	1	1	0	1	1	1	1	1	0	1
<i>M. argentina</i>	1	0	1	1	1	1	1	1	1	1	1	1	1	0	1
<i>M. mendocina</i>	1	0	1	1	1	1	1	0	1	1	1	1	0	0	1
<i>M. decumbens</i>	1	0	1	1	1	1	0	0	1	1	1	1	0	0	0
<i>M. andina</i>	1	0	1	1	1	1	0	0	1	1	1	1	1	0	0
<i>D. pulchella</i>	0	0	1	1	1	1	1	1	1	1	1	1	0	0	1
<i>E. pilosum</i>	0	1	0	0	1	1	1	1	1	1	1	0	0	1	0
<i>B. bigelovii</i>	0	0	0	0	1	0	1	1	0	-	1	0	0	0	[0,1]
<i>M. montana</i>	0	0	0	0	0	0	0	1	1	0	1	0	0	0	0
<i>A. adscensionis</i>	1	0	0	0	0	1	0	0	1	0	0	0	0	1	1
<i>B. erectum</i>	0	1	0	0	0	0	0	1	1	0	0	0	0	0	0

	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50
<i>M. squarrosa</i>	1	[0,1]	1	1	0	0	1	0	0	1	0	0	0	1	1
<i>M. argentina</i>	1	1	0	1	0	0	0	1	0	0	0	0	1	1	1
<i>M. mendocina</i>	1	1	0	0	-	-	-	-	0	0	0	0	1	1	1
<i>M. decumbens</i>	1	0	1	1	0	1	0	0	1	1	1	0	0	0	1
<i>M. andina</i>	1	0	1	1	0	1	1	0	1	1	1	1	1	2	3
<i>D. pulchella</i>	1	0	0	1	0	0	0	0	1	1	1	0	0	1	1
<i>E. pilosum</i>	1	0	1	1	0	0	0	0	1	1	0	0	0	1	1
<i>B. bigelovii</i>	1	0	0	1	1	0	0	0	2	2	1	1	0	1	2
<i>M. montana</i>	1	0	0	1	0	0	0	0	1	1	1	0	0	1	0
<i>A. adscensionis</i>	0	0	0	1	0	1	0	0	0	0	1	1	0	1	0
<i>B. erectum</i>	0	0	0	1	1	1	0	0	0	0	1	0	0	0	0

APPENDIX 3. Continued.

	51	52	53	54	55	56	57	58	59	60	61	62	63
<i>M. squarrosa</i>	0	1	[0,1]	0	1	1	[1,2]	0	0	1	1	0	0
<i>M. argentina</i>	1	1	0	0	1	1	1	1	0	1	1	0	1
<i>M. mendocina</i>	1	1	0	1	-	-	2	1	1	1	1	0	1
<i>M. decumbens</i>	1	1	0	1	-	-	2	1	1	1	1	0	1
<i>M. andina</i>	1	1	0	0	1	0	2	1	1	1	1	0	1
<i>D. pulchella</i>	0	0	1	0	1	0	0	0	0	1	1	0	0
<i>E. pilosum</i>	0	0	1	0	1	1	1	0	0	0	0	0	1
<i>B. bigelovii</i>	0	0	1	1	-	-	1	1	1	0	0	1	1
<i>M. montana</i>	0	0	1	0	1	0	0	0	0	0	0	0	1
<i>A. adscensionis</i>	0	0	1	0	0	0	0	0	0	0	0	1	?
<i>B. erectum</i>	0	0	1	0	0	0	0	0	0	0	0	1	?