



Salud Uninorte

ISSN: 0120-5552

saluduninorte@uninorte.edu.co

Universidad del Norte

Colombia

Martínez, Emilio G.; Gutiérrez, Ana M.

Metabolic Syndrome and Body Composition measurements in Blue-Collar Workers from a
Metal-Mechanic Factory in Soledad. Preliminary Report

Salud Uninorte, vol. 33, núm. 1, 2017, pp. 1-6

Universidad del Norte
Barranquilla, Colombia

Disponible en: <http://www.redalyc.org/articulo.oa?id=81751405002>

- Cómo citar el artículo
- Número completo
- Más información del artículo
- Página de la revista en redalyc.org

redalyc.org

Sistema de Información Científica

Red de Revistas Científicas de América Latina, el Caribe, España y Portugal

Proyecto académico sin fines de lucro, desarrollado bajo la iniciativa de acceso abierto

Metabolic Syndrome and Body Composition measurements in Blue-Collar Workers from a Metal-Mechanic Factory in Soledad. Preliminary Report

Síndrome Metabólico y Mediciones de la Composición Corporal en Trabajadores de una Planta Metal-Mecánica en Soledad. Reporte Preliminar

Emilio G. Martínez¹, Ana M. Gutiérrez²

Abstract

Introduction: Data regarding the incidence of MetS and the relationship with some body composition measurements in blue-collar workers from Soledad were obtained, as part of a wider research.

Methods: Data on weight (W), height (H), BMI, waist, hip, WHR, percentage of body fat (%BF), fat mass, blood pressure (BP), fasting blood glucose (FBG), total cholesterol (TC), high density lipoprotein HDL, low density lipoprotein (LDL), very-low density lipoprotein (VLDL) and triglycerides (TG) were obtained from 236 male blue-collar workers in a metal-mechanic factory at Soledad, a municipality of Departamento del Atlántico in the Caribbean coast of Colombia. The Adult Treatment Panel III (2001) definition for metabolic syndrome was used. Metabolic Syndrome components (MetS) and correlations with body composition measurements were calculated.

Results: The overall prevalence of metabolic syndrome was 20.7% (95%CI=15%, 26%). Waist circumference above 102 cm. was found in 11.8% (95%CI=7.5%, 16.2%); 43.2% (95%CI=36.6%, 49.7%) had triglycerides levels greater than 150 mg/dl., blood pressure $\geq 130/85$ was found in 10.2% (95%CI=6.1%, 14.2%) and FBG >100 mg/dl in 10.5% (95%CI=6.4%, 14.7%). BMI and %BF correlates well with waist, BP, FBG and TG.

Conclusions: Prevalence of MetS in this blue-collar workers' sample at Soledad is in agreement with other papers published in Colombia, as well as in other countries. Abdominal obesity was below the national data. In this population, BMI and %BF could be a good proxy to evaluate MetS.

Key words: Metabolic syndrome; blue-collar workers; body composition; Colombia.

¹ Department of Medicine, Universidad del Norte, Barranquilla, Colombia

² Department of Public Health, Universidad del Norte, Barranquilla, Colombia

Correspondence: Emilio G. Martinez, Department of Medicine, Universidad del Norte, Km 5 Via Puerto Colombia, Barranquilla, Colombia. Telephone (57-5) 3 509 509. egmartinez@uninorte.edu.co

Resumen

Introducción: Se obtuvo un conjunto de datos, como parte de una investigación más amplia, en relación a la incidencia de Síndrome Metabólico (SM) y su relación con algunas mediciones de composición corporal en trabajadores en Soledad.

Métodos: Se registraron los datos del peso (P), talla (T), Índice de Masa Corporal (IMC), cintura, cadera, Índice Cintura-Cadera (ICC), porcentaje de grasa corporal (%GC), masa grasa, presión arterial (PA), glucosa sanguínea en ayunas (GSA), colesterol total (CT), lipoproteínas de alta densidad (LAD), lipoproteínas de baja densidad (LBD), lipoproteínas de muy baja densidad (LMBD) y triglicéridos (TG) en 236 obreros de una fábrica metalmecánica en Soledad, municipio del Departamento del Atlántico en la costa caribe Colombiana. Se utilizó para definir el Síndrome Metabólico (SM) los criterios del Adult Treatment Panel III (2001). Se calcularon los componentes del SM y las correlaciones con las mediciones de la composición corporal.

Resultados: La prevalencia global del Síndrome Metabólico fue de 20.7% (95%CI=15%, 26%). La circunferencia de la cintura por encima de 102cm. se encontró en 11.8% (95%CI=7.5%, 16.2%); 43.2% (95%CI=36.6%, 49.7%) tenían niveles de triglicéridos mayores a 150mg/dl; la presión sanguínea $\geq 130/85$ se encontró en 10.2% de los casos (95%CI=6.1%, 14.2%) y la GSA >100 mg/dl en 10.5% (95%CI=6.4%, 14.7%). El IMC y el %GC se correlacionaron bien con la cintura, la presión arterial, la glucosa sanguínea en ayunas y los triglicéridos.

Conclusiones: La prevalencia de SM entre los obreros en esta muestra en Soledad, está de acuerdo con otros trabajos publicados en Colombia, así como en otros países. La obesidad abdominal estuvo por debajo de los datos nacionales. En esta población, el IMC y el %GC pudieran ser un buen aproximado para evaluar el SM

Palabras clave: Síndrome Metabólico, trabajadores, Composición Corporal, Colombia.

INTRODUCTION

Metabolic Syndrome is defined by a constellation of an interconnected physiological, biochemical, clinical and metabolic factors that directly increase the risk of atherosclerotic cardiovascular disease, diabetes and all cause mortality (1). In Colombia, several papers have been published assessing the prevalence of metabolic syndrome (MetS) in specific subgroups of the population (2-9) but to the best of our knowledge there are no reports published in the country regarding the relationship between MetS and body composition in blue-collar workers.

The goal of this preliminary report was to describe values and relationships between

some anthropometric and metabolic variables in a metal-mechanic factory in the Colombian Caribbean coast.

MATERIAL AND METHODS

Two hundred thirty-six male blue-collar workers in a metal-mechanic factory at Soledad (a municipality of Departamento del Atlántico in the Caribbean coast of Colombia, South America) participated in this study. All the individuals who participated did so voluntarily and a written consent was read and signed by each of them and delivered to the research staff. Ethical approval was obtained from the Universidad del Norte Health Sciences Division Committee.

Metabolic syndrome was defined according to the Adult Treatment Panel III (10). Blood pressure (BP) was measured following recommended procedures (11). Blood samples after 12 hours overnight fasting period was drawn for lipid testing by registered nurses at the factory infirmary and delivered to a certified laboratory for analysis, where the enzymatic determination of total serum cholesterol (TC), High Density Lipoprotein (HDL), Low Density Lipoprotein (LDL), Very Low Density Lipoprotein (VLDL), Triglycerides (TG) and Fasting Blood Glucose (FBG) were carried out according to the laboratory standard procedure.

Height was measured in centimeters with a portable stadiometer (Holtain Ltd. UK); waist circumference was measured in centimeters with a non-stretchable measuring tape 1 cm above the iliac crest and the reading was taken at the end of a normal expiration and hip circumference was measured with the tape around the point with the maximum circumference over the buttocks.

Waist and hip measurements were taken by duplicate, in light underwear, and the average was used as definitive. Waist-Hip Ratio (WHR) was calculated from these two measurements.

Weight, Body Mass Index (BMI), fat mass (FM), % body fat (%BF) and Basal Metabolic Rate (BMR) were obtained using Tanita's BIA (Bioelectrical Impedance Analyzer) model 310 A (Tanita Corporation, Tokyo, Japan T174). Participants with metabolic disturbances were excluded. A strip of paper and a code number with each participant's data was recorded.

Data analysis was performed using SPSS statistical package, version 21 (IBM Software Group, Chicago, IL). Mean and standard deviation of all measurements were obtained. Correlations between measurements of body composition and components of metabolic syndrome were analyzed using linear regression.

RESULTS

Mean, standard deviation, minimum and maximum values of all measurements are listed in Tables 1 and 2 summarize the prevalence of metabolic syndrome among blue-collar workers.

Waist circumference above 102 cm. was found in 11.8% (95%CI=7.5%, 16.2%); 43.2% (95%CI=36.6%, 49.7%) had triglycerides levels higher than 150 mg/dl, blood pressure $\geq 130/85$ was found in 10.2% (95%CI=6.1%, 14.2%) and FBG >100 mg/dl in 10.5% (95%CI=6.4%, 14.7%). The overall prevalence of metabolic syndrome was 20.7% (95%CI=15%, 26%). Body composition measurements and metabolic syndrome components correlations are listed in Table 3. BMI and %BF correlates well with waist, BP, FBG and TG. There were no associations between HDL and body composition measurements and all subjects had HDL above 40 mg/dl, which is intended as the effect of intensive physical activity.

DISCUSSION

Obesity in the developing world can no longer be considered solely as a disease of groups of higher socioeconomic status groups (12,13) and in both sexes, lower socioeconomic position across life was associated with higher fat mass and higher android to gynoid ratio

in early old age (12). Abdominal obesity is one of the clustering factors, which define metabolic syndrome (14). Studies from Europe and the USA found that blue-collar workers had three-times the risk of CVD (15) and ischemic heart disease (16,17) compared with white-collar workers.

Several papers have been published in Colombia providing data about the prevalence of metabolic syndrome (MetS) (2-5,8,18-21). The results are controversial due to differences among regions in the country and definition criteria: (1) In a survey of cardiovascular risks applied to 99 adults in the Caribbean coast of the country, 49.5% of the subjects had MetS according to the International Diabetes Foundation, 41.4% according to the American Heart Association, and 20.2% according to the American Treatment Panel III (22); (2) in Antioquia (a northwest Department) in two different cross-sectional studies conducted in 285 medical staff volunteers (29.1% male) aged 20-61 years and 3000 adults aged 25-64 years from Medellín and surrounding municipalities, global prevalence of MetS was 17.5% (19) and 41% (21) respectively and; (3) in the southwest, a cross-sectional study in 89 working women aged 25-64 years from the public and private sectors, 23% had MetS (8). The reported prevalence of MetS varies depending on the definition used, age, sex, socioeconomic status, and the ethnic background of study cohorts (23). In this preliminary report, 20.7% of the blue-collar workers had MetS, which is in agreement with data reported in the same area (22) as well as in the general population in the United States (23.7%) using Adult Treatment Panel-III criteria (24) and in Korean male blue-collar workers (25,26).

Metabolic syndrome alone cannot predict global cardiovascular disease risk, but abdominal obesity - the most prevalent manifestation of metabolic syndrome - is a marker of 'dysfunctional adipose tissue', and is of central importance in clinical diagnosis (27,28). In this population, BMI and %BF correlate well with waist, BP, FBG and TG and could be a good proxy to evaluate MetS. Visceral fat, which correlates well with waist circumference, is associated with higher insulin resistance and smaller LDL and HDL particle size (29). Abdominal obesity (AO) was 11.8% in this population, which is lower than it was reported in a similar environment, a metal-mechanic factory in Cali (28%) and national data, which accounts for 39% in men aged 18-64 years (20). It may be hypothesized that the socio-economic differences among regions in Colombia explain these findings (30). The scarcity of national data on the metabolic syndrome in blue-collar workers makes difficult to compare the results of this study locally.

LIMITATIONS

This was a pilot study whose main objective was to establish an initial assessment of the problem of metabolic syndrome in blue-collar workers in the Caribbean region. The research aims to cover in a second phase a wider sample in order to obtain regional reference values and educate workers on lifestyles and healthy eating.

CONCLUSIONS

Prevalence of MetS in blue-collar workers at Soledad, Colombia, is in agreement with other reports published on the general population in the country. Abdominal obesity, according with the cut-off values used in this report,

was below the national data. The correlations between MetS components and a set of body composition variables were in general strong and positive.

Acknowledgements: To all the people who participated in this research and the administrative officers from the metal-mechanic factory involved in this project.

Conflicts of interest: Neither of the authors had any conflicts of interest.

Financial disclosure: Emilio Martinez and Ana Maria Gutierrez have no financial disclosures.

REFERENCES

1. Kaur J. A comprehensive review on metabolic syndrome. *Cardiol Res Pract*. 2014;943162. <http://dx.doi.org/10.1155/2014/943162>.
2. Arteaga-Arredondo LF, Fajardo-Rodríguez H. Cardiovascular risk factor prevalence in civil aviation pilots in Colombia during 2005. *Rev Salud Pública*. 2010;12:250-256.
3. Dosman VA, Trivino LP, Uribe YL, Agredo RA, et al. Association between physical aptitude and metabolic syndrome risk factors in workers of a university institution. *Rev Col Cardiol*. 2009;16:153-158.
4. Feliciano-Alonso JE, Olimpo Mendivil C, Sierra Ariza ID, Pérez CE. Cardiovascular risk factors and metabolic syndrome in a population of young students from the National University of Colombia. *Rev Assoc Med Bras*. 2010;56:293-298. <http://dx.doi.org/10.1590/S0104-42302010000300012>.
5. Mesa A, Suarez MS, Arbelaez A, Mosquera M, et al. Lack of relationship of physical activity level with cardiovascular risk factors and metabolic syndrome in apparently healthy men. *Endocrinol Nutr*. 2011;58:68-74.
6. Pinzón JB, Serrano NC, Díaz LA, Mantilla G, et al. Impacto de las nuevas definiciones en la prevalencia del síndrome metabólico en una población adulta de Bucaramanga, Colombia. *Biomedica*. 2007;27:172-179.
7. Navarro E, Vargas RF. Riesgo coronario según ecuación de Framingham en adultos con síndrome metabólico de la ciudad de Soledad, Atlántico. *Rev Colomb Cardiol*. 2012;19:109-118.
8. Davila EP, Quintero MA, Orrego ML, et al. Prevalence and risk factors for metabolic syndrome in Medellín and surrounding municipalities, Colombia, 2008–2010. *Preventive Medicine*. 2012;56:30–34. <http://dx.doi.org/10.1016/j.ypmed.2012.10.027>.
9. González-Zapata LI, Deossa GC, Monsalve-Álvarez J, et al. Metabolic syndrome in healthcare personnel at the University of Antioquia-Colombia; LATINMETS study. *Nutr Hosp*. 2013;28:522-531. <http://dx.doi.org/10.3305/nh.2013.28.2.6315>.
10. National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation*. 2002;106(25):3143-421.
11. Pickering TG, Hall JE, Appel LJ, et al. Recommendations for blood pressure measurements in humans and experimental animals Part1: Blood pressure measurements in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Hypertension*. 2005; 45:142-161. <http://dx.doi.org/10.1161/01.HYP.0000150859.47929.8e>.
12. Bann D, Cooper R, Wills AK, Adams J, Kuh D. Socioeconomic position across life and body composition in early old age: findings from a British birth cohort study. *J Epidemiol Community Health*. 2014;68:516-523. <http://dx.doi.org/10.1136/jech-2013-203373>.

13. Monteiro CA, Moura EC, Conde WL, Popkin BM. Socioeconomic status and obesity in adult populations of developing countries: a review. *Bulletin of the World Health Organization*. 2004;82:940-946.
14. Samson SL, Garber AJ. Metabolic syndrome. *Endocrinol Metab Clin N Am*. 2014;43:1-23. <http://dx.doi.org/10.1016/j.ecl.2013.09.009>.
15. Netterstrom B, Nielsen FE, Kristensen TS, Bach E, Moller L. Relation between job strain and myocardial infarction: A case control study. *Occup Environ Med*. 1999;56:339-342.
16. Chen EY, Lee YW, Kim HS. Job categories and acute ischemic heart disease: A hospital-based, case-control study in Taiwan. *Am J Ind Med*. 2007;50:409-414. <http://dx.doi.org/10.1002/ajim.20462>.
17. Netterstrom B, Kristensen TS, Sjol A. Psychological job demands increase the risk of ischaemic heart disease: a 14-year cohort study of employed Danish men. *Eur J Cardiovasc Prev Rehabil*. 2006;13:414-420.
18. <http://dx.doi.org/10.1097/01.hjr.0000201512.05720.87>.
19. Villa-Roel C, Buitrago A, Rodríguez DC, et al. Prevalence of metabolic syndrome in scholars from Bucaramanga, Colombia: a population-based study. *BMC Pediatrics*. 2009;9:28. <http://dx.doi.org/10.1186/1471-2431-9-28>.
20. González-Zapata LI, Deossa GC, Monsalve-Álvarez J, et al. Metabolic syndrome in healthcare personnel at the University of Antioquia-Colombia; LATINMETS study. *Nutr Hosp*. 2013;28(2):522-531. <http://dx.doi.org/10.3305/nh.2013.28.2.6315>.
21. Agredo-Zúñiga RA, García-Ordoñez ES, Osorio C, et al. Obesidad abdominal y ausentismo por causa médica en una empresa de la industria metalmecánica en Cali, Colombia. *Rev Peru Med Exp Salud Publica*. 2013;30(2):251-5.
22. Suárez-Ortegón MF, Arbeláez A, Mosquera M, Ramírez-Vélez R, Aguilar-De Plata C. Evaluation of the relationship between self-reported physical activity and metabolic syndrome and its components in apparently healthy women. *Biomédica*. 2014;34:60-66. <http://dx.doi.org/10.7705/biomedica.v34i1.1442>.
23. Navarro E, Vargas RF. 2012. Riesgo coronario según ecuación de Framingham en adultos con síndrome metabólico de la ciudad de Soledad, Atlántico 2010.
24. *Rev Colomb Cardiol*. 19: 109-118.
25. Samson SL, Garber AJ. Metabolic syndrome. *Endocrinol Metab Clin N Am*. 2014;43:1-23. <http://dx.doi.org/10.1016/j.ecl.2013.09.009>.
26. Ford E, Giles W, Dietz W. Prevalence of the metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA*. 2002;287:356-9. <http://dx.doi.org/10.1001/jama.287.3.356>.
27. Hwang WJ, Lee CY. Effect of psychosocial factors on metabolic syndrome in male and female blue-collar workers. *Japan Journal of Nursing Science*. 2014;11:23-34. <http://dx.doi.org/10.1111/j.1742-7924.2012.00226.x>.
28. Wong JU, Hong OS, Hwang WJ. Actual Cardiovascular Disease Risk and Related Factors. *Workplace Health Saf*. 2013;61(4):163-171. <http://dx.doi.org/10.3928/21650799-20130327-17>.
29. Després JP1, Lemieux I. Abdominal obesity and metabolic syndrome. *Nature*. 2006;444(7121):881-887. <http://dx.doi.org/10.1038/nature05488>.
30. Phillips LK, Prins JB. The link between abdominal obesity and the metabolic syndrome. *Curr Hypertens Rep*. 2008;10:156-64.
31. Oda E. The metabolic syndrome as a concept of adipose tissue disease. *Hypertens Res*. 2008;31:1283-1291. <http://dx.doi.org/10.1291/hypres.31.1283>.
32. Galvis LA, Meisel-Roca L. Persistencia de las desigualdades regionales en Colombia: un analisis espacial. Banco de la República. Documento No. 120. 2010.