



Salud Uninorte

ISSN: 0120-5552

carvica@gmail.com

Universidad del Norte

Colombia

López-Soto, Lina Maria; López-Soto, Olga Patricia; Osorio-Forero, Alejandro; Restrepo, Francia; Tamayo-Orrego, Lukas

Muscle Activity and Muscle Strength in Atypical Swallowing

Salud Uninorte, vol. 33, núm. 3, septiembre-diciembre, 2017, pp. 273-284

Universidad del Norte

Barranquilla, Colombia

Available in: <http://www.redalyc.org/articulo.oa?id=81753881002>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Muscle Activity and Muscle Strength in Atypical Swallowing

Actividad Muscular y Fuerza Muscular en la Deglución Atípica

Lina Maria López-Soto¹, Olga Patricia López-Soto², Alejandro Osorio-Forero³, Francia Restrepo⁴, Lukas Tamayo-Orrego⁵

Abstract

Objective: To characterize muscle activity and muscle strength in patients with atypical swallowing (AS; $n = 88$) and competent lips (CL) or incompetent lips (IL) versus a control group (Ctrl; $n = 90$)

Methods and materials: an analytical case-control study was conducted using surface electromyography (sEMG) of the orbicularis oris, mentalis and masseter muscles. Compression forces of the orbicularis oris, right and left masseters muscle (Cfrmm, Cflmm), tongue tensile strength (Tts) via myoscan analysis and resistance of the orbicularis oris (Roo) via dynamometry were determined.

Comparisons were made with the Mann-Whitney U test under a 95% confidence interval.

Results: The amplitude of the sEMG record of the orbicularis oris muscle, in maximal contraction, was lower ($p < 0.01$) in the atypical swallowing group (596.40 ± 175.83) compared to controls (659.83 ± 203.79). The muscle strength studied in the experimental group was lower ($p < 0.01$) than in controls (Cfo: AS: 0.31 ± 0.13 ; Ctrl: 0.38 ± 0.11 ; Cfrmm: AS: 0.40 ± 0.08 ; Ctrl: 0.50 ± 0.11 ; Cflmm: AS: 0.41 ± 0.08 ; Ctrl: 0.58 ± 0.59 ; Tts, AS: 0.52 ± 0.14 , Ctrl: 0.65 ± 0.14 , and Roo: AS: 2.47 ± 0.61 ; Ctrl: 2.73 ± 0.60). Patients with incompetent lips had a greater muscle activity of the orbicularis oris in swallowing (AS of IL: 197.01 ± 85.84 ; AS of CL: 160.54 ± 97.03 ; Ctrl: 147.18 ± 80.10).

Conclusion: Patients with atypical swallowing showed differences in the strength of studied muscles and the muscle activity of the orbicular oris muscle compared to controls.

Keywords: Swallowing, Lips, Surface Electromyography, Muscle Strength.

¹ Oral Health Research Group, Autonomía University of Manizales (UAM), Antigua Estación del Ferrocarril, Manizales, Caldas; e-mail: linalopez@autonomia.edu.co, fax 8727670

² Oral Health Research Group, UAM, Antigua Estación del Ferrocarril, Manizales, Caldas; e-mail: sonrie@autonomia.edu.co, fax 8727670

³ Neurolearning Research Group, UAM, Antigua Estación del Ferrocarril, Manizales, Caldas; e-mail: alejosorio1@gmail.com, fax 8727670

⁴ Neurolearning Research Group, UAM, Antigua Estación del Ferrocarril, Manizales, Caldas; e-mail: franci46@gmail.com, fax 8727670

⁵ Neurolearning Research Group, UAM, Antigua Estación del Ferrocarril, Manizales, Caldas; e-mail: lukas-tamayo@gmail.com, fax 8727670

Correspondence: sonrie@autonomia.edu.co; Phone: (576) 8727272-162; Fax: fax 8727670

Fecha de recepción: 11 de junio de 2017
Fecha de aceptación: 18 de septiembre de 2017

Resumen

Objetivo: caracterizar la actividad y fuerza muscular de pacientes con deglución atípica (DA; N=88) con competencia labial (CL) o incompetencia labial (IL) vs un grupo control (Ctrl; N=90).

Materiales y métodos: estudio analítico de casos y controles se realizó una electromiografía de superficie (sEMG) de los músculos orbicular, mental y maseteros; se determinaron las fuerzas compresiva del orbicular de los labios (FCo), contráctil del masetero derecho (FCmd) e izquierdo (FCmi), extensora del músculo lingual (FEl) por mioescanografía y la fuerza de resistencia del orbicular de los labios (FRo) por dinamometría.

Resultados: La amplitud del registro sEMG del músculo orbicular de los labios, en contracción máxima, fue menor ($p < 0.01$) en el grupo DA ($596,40 \pm 175,83$) con respecto a los controles ($659,83 \pm 203,79$); la fuerza de los músculos estudiados en el grupo experimental fue menor ($p < 0,01$) que en los controles (FCo: DA: $0,31 \pm 0,13$; Ctrl: $0,38 \pm 0,11$; FCmd: DA: $0,40 \pm 0,08$; Ctrl: $0,50 \pm 0,11$; FCmi: DA: $0,41 \pm 0,08$; Ctrl: $0,58 \pm 0,59$; FEl: DA: $0,52 \pm 0,14$; Ctrl: $0,65 \pm 0,14$; y FRo: DA: $2,47 \pm 0,61$; Ctrl: $2,73 \pm 0,60$). Los pacientes con DA-IL presentaron mayor actividad muscular del orbicular en deglución (DA-IL: $197,01 \pm 85,84$; DA-CL: $160,54 \pm 97,03$; Ctrl: $147,18 \pm 80,10$).

Conclusiones: Los pacientes con DA difieren en la fuerza de los músculos estudiados y en la actividad muscular del orbicular con respecto a los controles. Los pacientes con DA-CL y DA-IL difieren en la actividad y fuerza muscular del orbicular.

Keywords: Deglución, labios, electromiografía de superficie, Fuerza muscular.

INTRODUCTION

Disorders in oral functions such as mastication, breathing and swallowing and their physiological manifestations such as incompetent lips can determine the occurrence or worsening of occlusal disorders. These can also be seen in the facial appearance of patients. (1-3). Muscle management of disorders in oral functions emphasizes the need for non-invasive methods to qualify and quantify these disorders. Non-invasive methods may help give objective information about the muscle activity and muscle strength in patients with normal swallowing and atypical swallowing (AS) both with incompetent lips (IL) and with competent lips (CL). Surface electrode electromyography (sEMG), myoscan analysis and dynamometry (1) are some measurements of choice for these cases because of their low cost and easy implementation.

Swallowing is an automatic motor action, in which respiratory muscles and gastrointestinal tract intervene. The purpose of swallowing is

the transport of food bolus and the cleansing of the respiratory tract. Swallowing involves up to 30 muscles and 6 encephalic pairs (4-6).

Atypical swallowing is defined as the anterior or lateral pressure of the tongue against the dental arches, or placed in an anterior or lateral position during swallowing. Atypical swallowing can be diagnosed by visual observation of perioral muscles, and using Payne's technique (7). Incompetent lips are diagnosed by clinical observation of lip position at rest. These diagnoses do not allow for quantitative control to guarantee greater accuracy in the assessment of muscle activity and muscle strength involved in the function. Electromyography, myoscan analysis, and dynamometry could be the solution to this problem.

Techniques to record the muscle activity via EMG are well defined and tested. Potential errors in methodology such as electrode placement have been well described (8,9).

Clinical use of surface electromyography has been proposed in the fields of diagnosis, treatment and control of oral disorders at the muscular and functional levels(10-13). This is based on the concept that multiple diseases or dysfunctional conditions can be differentiated by means of surface electromyography (sEMG) records of muscle activity⁽¹⁴⁻¹⁸⁾, abnormal occlusal positions(19-21), functional hypoactivity and hyperactivity (22,23), muscle spasms (24-25), fatigue(26-27) and muscle imbalance(28). Surface electromyography has been suggested as a useful means to record changes in muscle function prior and posterior to therapeutic interventions as an evidence of treatment success(10, 12). For more than a decade, surface electromyography (sEMG) was considered as an alternative to previous studies in order to assess muscle activity associated with swallowing(29); particularly, sEMG studies showed promising results in both adults(30, 31) and children(5) in this knowledge area.

In the 1970s, myoscanners and dynamometers were described as measurement instruments of orofacial muscle activity. These devices were designed by Dr. Richard Shepard and Dr. Daniel Garliner, who have a reference scale of values for the U.S. children and adult population(32).

The purpose of this study was to characterize the values of muscle activity using sEMG and muscle strength using myoscan and dynamometry analyses in patients with atypical swallowing and competent lips or incompetent lips versus a control group.

MATERIALS AND METHODS

This analytical case-control study used the Statcalc calculator in the Epi Info 6 software for the calculation of the sample. Sample was calculated considering an 80% power (1-Beta), a ratio of 1:1 and an expected prevalence of

factor in the non-exposed group of 30% (presence of an abnormal sEMG in normal patients), an OR of 2.5 (72% of probability to show an abnormal electromyography in the exposed group) and a prevalence of factor in the exposed group of 51.72 %. The sample would include 88 patients in the exposed group and 88 in the control group. Finally, 90 school students (51 girls and 39 boys) aged between 7 and 13 (10.55 ± 1.21) in the control group and 88 school students (55 girls and 33 boys) aged between 7 and 13 (10.07 ± 1.23) diagnosed with atypical swallowing were examined.

School students were selected from three high schools in the city of Manizales (Caldas, Colombia). This study requested an authorization from the Ethics Committee of the Autonomía University of Manizales (Minutes No. 12 of February 2, 2011) and met all the requirements of the 8430 rule, governing health research applied to patients. Parents or authorized guardians of students received the necessary information about the experience in which they would participate and they previously signed the informed consent. Similarly, students were requested their authorization to participate in the research project.

Diagnosis of atypical swallowing was performed with the traditional protocol, using a Payne black light, model UVL-56, Black Ray lamp, long wave UV (nm), 115 volts, 60 Hz, 0.16 Amps, UVP, Upland, CA91786, USA. This was used to know the specific location of the tongue during swallowing. During the examination, the patient was sitting with the back straight, head against the wall, and feet on the floor. The patient should have pulled the tongue forward to place a 10% fluorescein solution in the anterior part, apex and lateral sides. Patients were asked

to swallow in order to observe, then, the place where the solution was recorded intra-orally using the black light.

SEMG PROCEDURE

sEMG records were performed with the electrophysiograph and Cadwell Sierra Waves software. Acquisition parameters were: sampling frequency of 76.8 kHz, the bandpass filter 10 to 10k Hz and gain 200 μ V. For acquisition and storage of records, the EMG protocol was selected. 302139-200 stainless steel disk electrodes (Cadwell) of a 10 mm diameter were used. To perform the sEMG record, school students were sitting against the back of a chair with the back straight, arms relaxed, feet on the floor, eyes open and staring at a point in front of them. Prior to the positioning of the electrodes, their faces were cleaned with a piece of cotton with 95% ethanol, Cadwell conductivity gel model 202153-000 was used in order to reduce the artifact occurrence due to the low conductance. The ground electrode, common to all electrode pairs, was placed 2 cm above the nasion. This study assessed the muscle activity of the right and left masseter muscles and left orbicularis oris and mentalis muscles; these were assessed unilaterally because of their symmetric function; these muscles work predominantly in the first phase of swallowing. A bipolar configuration was selected for masseter, orbicularis oris and mentalis muscles. These configurations are sensitive to changes in the strength exerted by a muscle (33). Electrodes were placed as follows: *Left masseter*: subjects were asked to close teeth with maximal force. Masseter muscle was then examined. An active electrode was placed on the motor point, reference electrode was placed 1 cm under the ear lobe. *Left orbicularis oris*: Active electrode was placed 1 cm above the upper lip vermillion and a reference electrode was placed on the zygomatic

bone. Finally, *Mentalis muscle*: active electrode was placed on the chin, 1 cm from the midline and a reference electrode was placed 2 cm above the distal direction.

Every muscle was assessed independently. Maximal voluntary contraction (MVC) of the orbicularis oris muscle (10 seconds of pursed lips) was determined. Nine swallows of 25 ml of water (three by each muscle) were requested. Subjects were requested to keep the liquid in the mouth until staying at rest. Subjects were instructed to “swallow as usual” to promote the normal way of doing it by each subject. Elapsed time during the sEMG process did not exceed 10 minutes.

This study examined the amplitude of muscle activity while swallowing and the MVC of the orbicularis oris muscle. Amplitude corresponds to the peak to peak voltage of the interference pattern of the muscle activity at the time of swallowing.

MYOSCAN PROCEDURE

The myoscan used was the Pounds Myoscanner, Neilco Technology Inc., D-926119T8 Hoxt, 1volt=FS., pt 2500. The physiological force of muscles was estimated by means of a plug. This allowed the measurement of the lip compression strength, strength of tongue projection and contraction of masseter muscles. To do the examination, patients were asked to sit comfortably with their head against a wall and feet on the floor. First, the right masseter muscle contraction was assessed. Patients were asked to close their teeth tightly. The examiner touched the muscle with the tips of the index and middle fingers; the same action was repeated three times in order to determine its location. Then, the test began by calibrating the myoscan in order for its needle to be exactly 0.0 pounds. A plate was

set on the masseter muscle and patients were asked again to have a maximal occlusion. The same procedure was made on the left masseter (Reference values: the result was from 0.4 to 0.6 pounds for students aged between 4 and 10 years old and the result was from 0.6 to 0.8 pounds for students older than 10 years old).

The measurement of the compression strength of the orbicularis oris muscle began with the myoscan calibration set at 0.0 pounds. Patients were asked to close the teeth, stretch lips forward and press the plate strongly. (Reference values: the result was from 0.2 to 0.4 pounds for students aged between 4 and 10 years old and the result was from 0.4 to 0.6 pounds for students older than 10 years old). Finally, the tongue tensile strength was assessed. Patients were instructed to protrude the tongue through the hole in the plate held by the examiner. (Reference values: the result was from 0.6 to 0.8 pounds for students aged between 4 and 10 years old and the result was from 0.8 to 1.2 pounds for students older than 10 years old). All measurements of the myoscan procedure were repeated three times and the average was written for each of them.

DYNAMOMETRY PROCEDURE

The dynamometer used was Chatillon Inc. (New York, NY, USA), model CHA IN-10. Results were measured in pounds of force and were taken with a dynamometer with a hook attached to a six-inch cord, ending in a button of 1 inch in diameter. The button was placed in front of the incisors and the back of the lips. The researcher pulled the dynamometer until the button was expelled and then the result was read. The normal range for lip force, listed by Garliner, is of 3 to 5 pounds(7). To do the examination, patients were sitting, head against the wall, arms at their sides, and feet on the floor.

STATISTICAL ANALYSIS

Statistical analysis was performed with the SPSS statistical package (SPSS® IBM® v.20). Comparisons were made with the Mann-Whitney U test because most variables did not meet the normality tests (Kolmogorov-Smirnov) nor homogeneity of variance (Levene's test). These variables were analyzed under a 95% confidence interval.

RESULTS

The amplitude of the sEMG record of orbicularis oris muscle when swallowing was significantly higher in the IL group than in the CL group and the control group (Table 2). This difference among groups was only significant for the right masseter. The amplitude of the maximal voluntary contraction of the orbicularis oris muscle was significantly lower for patients with atypical swallowing than for the control group. In the mentalis muscle, when swallowing, the amplitude of the sEMG showed no significant differences among the groups (Table 1).

The student group with AS was divided into patients with competent lips (CL) and incompetent lips (IL). The CL group showed an amplitude of the sEMG record of the right masseter muscle significantly higher than that of the control group. The amplitude of the sEMG record of orbicularis oris muscle when swallowing was significantly in the IL group than in the CL group and the control group (Table 2).

In maximal contraction of the orbicularis oris, the amplitude of the sEMG record was significantly lower in the CL group than in the IL group and the control group.

Table 1. Amplitudes of the EMG record and strength of the muscles studied in patients with AS versus controls

Variable			Mean +/- AS	Confidence interval for the mean at 95%
Amplitude of SEMG (µV)				
Right masseter	Swallowing*	Ctrl	94,96 ± 68,10	74,68 - 108,79
		AS	79,26 ± 48,03	68,91 - 92,15
Left masseter	Swallowing	Ctrl	84,61 ± 49,34	72,98 - 98,23
		AS	80,79 ± 56,91	65,81 - 94,06
Orbicularis oris	Swallowing	Ctrl	169,39 ± 85,22	150,82 - 187,01
		AS	149,35 ± 72,27	127,95 - 162,91
	MVC**	Ctrl	166,04 ± 5,67	154,76 - 177,33
		AS	142,26 ± 6,02	130,28 - 154,23
Mentalis	Swallowing	Ctrl	210,55 ± 113,18	192,23 - 244,13
		AS	215,78 ± 99,91	182,48 - 230,17
Strength (pounds)				
Compression strength of orbicularis oris**		Ctrl	0,38 ± 0,11	0,35 - 0,40
		AS	0,31 ± 0,13	0,28 - 0,34
Right masseter contractile**		Ctrl	0,50 ± 0,11	0,48 - 0,53
		AS	0,40 ± 0,08	0,39 - 0,42
Left masseter contractile**		Ctrl	0,58 ± 0,59	0,45 - 0,70
		AS	0,41 ± 0,08	0,39 - 0,43
Tongue tensile muscle**		Ctrl	0,65 ± 0,14	0,63 - 0,68
		AS	0,52 ± 0,14	0,49 - 0,55
Resistance of the orbicularis oris**		Ctrl	2,73 ± 0,60	2,61 - 2,86
		AS	2,47 ± 0,61	2,34 - 2,60

Table 1. Amplitude of sEMG waveforms and strength measurements of studied muscles. Comparison of measurements of patients with atypical swallowing and the control group. (Ctrl) Control group (n =90). (AS) Atypical swallowing group (n =88). (a) Calculation made with Mann-Whitney U test for two independent samples. (*) p<0.05. (**) p<0.01.

Table 2. Amplitudes of the EMG record and strength of the muscles studied in patients with SA and CL and IL versus controls

Variable	Class	Mean \pm DE	Confidence interval for the mean at 95%
Amplitude SEMG (μ V)			
Right Masseter Swallowing ¥	Ctrl	74.88 \pm 46.4	65.16 - 84.60
	IL	68.82 \pm 25.64	58.68 - 78.97
	CL**	89.24 \pm 52.34	75.83 - 102.64
Left Masseter Swallowing	Ctrl	74.15 \pm 39.04	65.97 - 82.33
	IL	79.17 \pm 37.35	64.40 - 93.94
	CL	86.84 \pm 57.02	72.24 - 101.45
Swallowing Orbicularis oris ¥	Ctrl	147.18 \pm 80.10	130.41 - 163.96
	IL**	197.01 \pm 85.84	163.06 - 230.97
	CL*	160.54 \pm 97.03	135.69 - 185.39
MVC Orbicularis oris¥	Ctrl	659.83 \pm 203.79	617.15 - 702.51
	IL**	677.35 \pm 164.41	612.31 - 742.39
	CL††	560.56 \pm 169.89	517.05 - 604.07
Mentalis Swallowing	Ctrl	205.06 \pm 101.64	183.78 - 226.35
	IL	207.92 \pm 103.54	166.97 - 248.88
	CL	230.04 \pm 136.75	195.01 - 265.06
Strength (pound)			
Compression of the orbicularis oris¥	Ctrl	0.376 \pm 0.11	0.35 - 0.40
	IL**	0.25 \pm 0.17	0.19 - 0.32
	CL††	0.34 \pm 0.10	0.31 - 0.36
Resistance of the orbicularis oris¥	Ctrl	2.73 \pm 0.60	2.61 - 2.86
	IL**	2.04 \pm 0.67	1.77 - 2.30
	CL††	2.66 \pm 0.48	2.54 - 2.77
Right Masseter Contractile ¥	Ctrl	0.50 \pm 0.11	0.48 - .53
	IL**	0.40 \pm 0.08	0.37 - 0.43
	CL**	0.40 \pm 0.08	0.38 - 0.42
Left Masseter Contractile ¥	Ctrl	0.58 \pm 0.59	0.45 - 0.70
	IL**	0.41 \pm 0.07	0.38 - 0.44
	CL**	0.41 \pm 0.09	0.39 - 0.43
Tongue tensile muscle ¥	Ctrl	0.65 \pm 0.14	0.63 - 0.68
	IL**	0.50 \pm 0.20	0.42 - 0.57
	CL**	0.53 \pm 0.11	0.51 - 0.56

Table 2. Amplitude of the sEMG waveforms and measurements of strength of the muscles studied. Comparison of measurements of patients with atypical swallowing and the control group. Ctrl: Control group (n =90). IL: Group of atypical swallowing with incompetent lips (n =27). CL: Group of atypical swallowing with competent lips (n =61).

(¥) χ^2 of the Kruskal-Wallis test with $p < 0.05$.

(†) Calculation made with Mann-Whitney U test for the two independent samples.

(*) Differences in respect to the control group $p < 0.05$.

(**) Differences in respect to the control group $p \leq 0.01$.

(††) Differences among the AS-CL and AS-IL groups $p \leq 0.01$.

Contractile forces of the masseter muscles, the compression strength of the orbicularis oris muscle and tongue tensile strength were measured by means of the myoscan analysis and the resistance force of the orbicularis oris muscle via dynamometry. Patients diagnosed with atypical swallowing showed a decrease in the compression strength and resistance strength of the orbicularis oris, contractile force of masseter muscles, and tongue tensile muscle (Table 1).

When analyzing the strength of these muscles in the CL and IL groups, the findings were as follows. The control group showed higher contractile forces of the masseter muscles, tongue tensile muscle, and resistance of the orbicularis oris compared to both IL and CL groups ($p < 0.05$). There were no differences between these two groups. The IL group showed a compression force and resistance of the orbicularis oris lower than the control group and CL group (Table 2).

DISCUSSION

In this research, school students with atypical swallowing (AS) showed greater electromyographic activity than patients in the control group. The contraction of mentalis and orbicularis oris muscles during atypical swallowing has been considered a clinical sign of abnormality. Stormer et al. (34) reported that patients with AS, during saliva or water swallowing, may have an increased electromyographic activity of perioral muscles, contrary to the findings of this research work. They also found a decreased activity of the masticatory muscles, which was similar to the findings of this research work.

Patients with AS and IL showed higher electrical amplitude than LC and controls in the orbicularis oris. These patients with AS would

show higher amplitude in the waveforms of these muscles due to a harder effort when swallowing. The more motor units activated to produce the same force, the higher the reduction of the electromechanical efficiency. When more effort is exerted on these muscles, there is higher amplitude recorded in the activity. Thus, the increase in amplitude would be caused by a higher amount of recruited fibers, not necessarily by higher strength, which is similar to Archer's records.

In this work, the average contractile force of the tongue, registered with the myoscan, was lower in the AS group. The role of the tongue in swallowing is well known, however, as Lear Moores (36), the tongue produces four times more pressure during swallowing than the perioral musculature. In normal swallowing, most of this pressure is exerted on the palate, but not on the teeth, unlike to what happens in atypical swallowing, where these forces are directly applied on the teeth, causing protrusion of the incisors and an increase in the horizontal dimension. Rogers (37) studied swallowing patterns and concluded that the majority of the children who presented tongue pressure in swallowing had overjet with 'short' lips, and the tongue moved itself freely through the anterior teeth without touching the lips. In his study, Stormer (34) found a lower sEMG activity in the masticatory muscles of patients with AS than in controls. This author stated these patients seem to have perioral muscle activity without the intervention of the temporalis muscle and higher activity in the orbicularis oris and mentalis muscles; these findings are different from the ones in this research work.

Electromyographic studies done by Ahlgren (38) and Tosello et al. (39) have shown that the orbicularis oris muscle has a greater activity in patients class II/I with incompetent

lips during swallowing than in patients with competent lips. Findings in these studies show that patients with increased lower facial third proportions and/or maxillomandibular anteroposterior discrepancies are more susceptible to an increased perioral muscle activity and incompetent lips. Gustafsson and Ahlgren (8) found greater activity of the orbicularis oris (upper and lower) in the resting posture of the lips, during swallowing. They state that perioral contraction during atypical swallowing is a common characteristic for individuals with incompetent lips, independent of tongue function.

The pressures exerted by the lips and cheeks in the buccal stage of swallowing are associated mainly with labial sealing and with the expulsion of liquids from the buccal vestibule rather than with the labial musculature contraction in the elevation of the larynx. In this first stage, larger potentials are generated by the orbicularis oris (40).

Lowe *et al.* (41) claim that patients with IL have a reduced activity of the orbicularis oris muscle, which is opposite to the findings in this study. Patients with IL have difficulties to chew with labial contact; this explains the higher activity of the perioral muscle, which would be reflected on the results of this research. Pallú *et al.* also observed a great activity of the orbicularis oris in patients with IL (42). This would mean that these patients should make a greater effort to obtain good closure during swallowing.

In the study of Dutra *et al.* (43) patients with incompetent lips showed masseter muscle hyperactivity during swallowing, suggesting a visceral type of swallowing. In this study, patients with AS and incompetent lips showed no masseter muscle hyperactivity, but there

was a muscle hyperactivity of the orbicularis oris. Balata (44) found that patients with incompetent lips showed a higher amplitude and a lower coefficient of variation of muscle activity, which are consistent with the records available for this study. During closure of the lips, it has been found that patients with incompetent lips show a muscle activity of the upper orbicularis oris and mentalis muscles greater than subjects with competent lips, who show, in most cases, a slight increase in mentalis muscle activity (8).

CONCLUSIONS

Amplitude of the sEMG record of left and right masseter, orbicularis oris and mentalis muscles in school students diagnosed with AS was lower than that of the control group; that amplitude was statistically significant in the orbicularis oris.

Patients with AS diagnosis showed a decrease in the compression and resistance forces of the orbicularis oris, contractile of masseter muscles and tongue tensile strength. The difference with respect to the control group was statistically significant.

The IL group recorded a higher activity in the orbicularis oris than LC and control groups; MVC of the orbicularis oris was statistically significant.

Acknowledgements

The authors are grateful to the Autonomía University of Manizales for providing the necessary financial resources to carry out this research project. Special thanks to the Translation Center of the Autonomía University of Manizales for the translation of this manuscript into English, especially to the translator: Carlos A. Muñoz Torres and reviser: Chris Cadman.

Recommendation

The use of the support materials described in this study (EMG, myoscan and dynamometry) gives the possibility to go beyond the subjective clinical criteria of the clinical diagnosis and to achieve more objective parameters to attain a reliable diagnosis.

Disclosure

The authors report no conflict of interests related to materials and methods used in this study or the findings reported in this project.

Financial: Autonoma University of Manizales (UAM).

REFERENCES

1. Farret S, Vitti M, Farret M. Electromyographic analysis of the mentalis and depressor labii inferior muscles in the production of speech. *Electromyography and clinical neurophysiology*. 1982;22(1-2):137.
2. Lowe AA, Johnston WD. Tongue and jaw muscle activity in response to mandibular rotations in a sample of normal and anterior open-bite subjects. *American Journal of Orthodontics*. 1979;76(5):565-76.
3. Proffit W. Equilibrium theory revisited: factors influencing position of the teeth. *The Angle Orthodontist*. 1978;48(3):175-86.
4. Logemann JA. Evaluation and treatment of swallowing disorders. 1983.
5. Vaiman M, Eviatar E, Segal S. Surface electromyographic studies of swallowing in normal subjects: a review of 440 adults. Report 1. Quantitative data: timing measures. *Otolaryngology-Head and Neck Surgery*. 2004;131(4):548-55.
6. Miller AJ. Deglutition. *Physiological Reviews*. 1982;62(1):129-84.
7. Hanson M. Orofacial myofunctional disorders: guidelines for assessment and treatment. *The International journal of orofacial myology: official publication of the International Association of Orofacial Myology*. 1988;14(1):27.
8. Gustafsson M, Ahlgren J. Mentalis and orbicularis oris activity in children with incompetent lips: an electromyographic and cephalometric study. *Acta Odontologica*. 1975;33(6):355-63.
9. O'Dwyer NJ, Quinn PT, Guitart BE, Andrews G, Neilson PD. Procedures for verification of electrode placement in EMG studies of orofacial and mandibular muscles. *Journal of Speech, Language and Hearing Research*. 1981;24(2):273.
10. Cooper BC, Cooper DL, Lucente FE. Electromyography of masticatory muscles in craniomandibular disorders. *The Laryngoscope*. 1991;101(2):150-7.
11. Goldstein L. The use of surface electromyography in objective measurement of the muscle function in facial pain/temporomandibular dysfunction patients. *The Functional orthodontist*. 2000;17(3):26.
12. Cooper BC. The role of bioelectronic instrumentation in the documentation and management of temporomandibular disorders. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*. 1997;83(1):91-100.
13. Cooper BC. The role of bioelectronic instruments in documenting and managing temporomandibular disorders. *The Journal of the American Dental Association*. 1996;127(11):1611.
14. Lous I, Sheik-Ol-Eslam A, Möller E. Postural activity in subjects with functional disorders of the chewing apparatus. *European Journal of Oral Sciences*. 1970;78(1-4):404-10.
15. Sheikholeslam A, Möller E, Lous I. Postural and maximal activity in elevators of mandible before and after treatment of functional disorders. *European Journal of Oral Sciences*. 1982;90(1):37-46.
16. Cooper B, Rabuzzi D. Myofascial pain dysfunction syndrome: a clinical study of asymptomatic subjects. *The Laryngoscope*. 1984;94(1):68-75.
17. Dolan E, Keefe F. Muscle activity in myofascial pain-dysfunction syndrome patients:

- a structured clinical evaluation. *Journal of craniomandibular disorders: facial & oral pain*. 1988;2(2):101.
18. Jankelson R. Scientific rationale for surface electromyography to measure postural tonicity in dental patients. *Cranio: the journal of craniomandibular practice*. 1990;8(3):207.
 19. Franks AS. Masticatory muscle hyperactivity and temporomandibular joint dysfunction. *The Journal of prosthetic dentistry*. 1965;15(6):1122-31.
 20. Funakoshi M, Fujita N, Takehana S. Relations between occlusal interference and jaw muscle activities in response to changes in head position. *Journal of dental research*. 1976;55(4):684-90.
 21. Michler L, Möller E, Bakke M, Andreassen S, Henningsen E. On-line analysis of natural activity in muscles of mastication. *Journal of craniomandibular disorders: facial & oral pain*. 1988;2(2):65.
 22. Möller E, Sheikholeslam A, Lous I. Response of elevator activity during mastication to treatment of functional disorders. *European Journal of Oral Sciences*. 1984;92(1):64-83.
 23. Yemm R. A neurophysiological approach to the pathology and aetiology of temporomandibular dysfunction*. *Journal of oral rehabilitation*. 1985;12(4):343-53.
 24. Ramfjord S. Bruxism, a clinical and electromyographic study. *J Am Dent Assoc*. 1961;62:36-58.
 25. Gordon Jr T. The influence of the herpes simplex virus on jaw muscle function. *The Journal of cranio-mandibular practice*. 1983;2(1):31.
 26. Bostel Av, Goudswaard P. Changes in EMG power spectra during fatigue in muscle contraction and migraine headache patients. *Headache: The Journal of Head and Face Pain*. 1983;23(5):223-8.
 27. Naeije M, Hansson T. Electromyographic screening of myogenous and arthrogenous TMJ dysfunction patients. *Journal of oral rehabilitation*. 1986;13(5):433-41.
 28. Festa F. Joint distraction and condyle advancement with a modified functional distraction appliance. *Cranio: the journal of craniomandibular practice*. 1985;3(4):343.
 29. Perlman A, Palmer P, McCulloch T, Vandaele D. Electromyographic activity from human laryngeal, pharyngeal, and submental muscles during swallowing. *Journal of Applied Physiology*. 1999;86(5):1663-9.
 30. Crary MA, Groher ME. Biomechanical correlates of surface electromyography signals obtained during swallowing by healthy adults. *Journal of Speech, Language and Hearing Research*. 2006;49(1):186.
 31. Crary M, Carnaby G, Groher M. Identification of Swallowing Events from sEMG Signals Obtained from Healthy Adults. *Dysphagia*. 2007;22(2):94-9.
 32. Garliner D. Myofunctional therapy. *General dentistry*. 1976;24(1):30.
 33. Gabriel DA. Effects of monopolar and bipolar electrode configurations on surface EMG spike analysis. *Medical engineering & physics*. 2011;33(9):1079-85.
 34. Störmer K, Panherz H. Electromyography of the perioral and masticatory muscles in orthodontic patients with atypical swallowing. *Journal of Orofacial Orthopedics/Fortschritte der Kieferorthopädie*. 1999;60(1):13-23.
 35. Archer SK, Garrod R, Hart N, Miller S. Dysphagia in Duchenne Muscular Dystrophy Assessed Objectively by Surface Electromyography. *Dysphagia*. 2012;1-11.
 36. Lear CS, Moorrees CF. Buccolingual muscle force and dental arch form. *American Journal of Orthodontics*. 1969;56(4):379-93.
 37. Rogers JH. Swallowing patterns of a normal-population sample compared to those of patients from an orthodontic practice. *American Journal of Orthodontics*. 1961;47(9):674-89.
 38. Tosello D, Vitti M, Berzin F. EMG activity of the orbicularis oris and mentalis muscles in children with malocclusion, incompetent lips and atypical swallowing—part I. *Journal of oral rehabilitation*. 1998;25(11):838-46.
 39. Ahlgren J. EMG studies of lip and cheek activity in sucking habits. *Swedish dental journal*. 1995;19(3):95.

40. Perkins RE, Blanton PL, Biggs NL. Electromyographic analysis of the " buccinator mechanism" in human beings. *Journal of dental Research*. 1977;56(7):783-94.
41. Lowe AA, Takada K, Taylor LM. Muscle activity during function and its correlation with craniofacial morphology in a sample of subjects with Class II, Division 1 malocclusions. *American Journal of Orthodontics*. 1983;84(3):204-11.
42. Pallú V, Magnani M, Berzin F, Bevilacqua D. Muscle Alterations in mouth breathing subjects. *Publicatio UEPG*. 1996;2:73-89.
43. Dutra EH, Maruo H, Vianna-Lara MS. Electromyographic activity evaluation and comparison of the orbicularis oris (lower fascicle) and mentalis muscles in predominantly nose-or mouth-breathing subjects. *American Journal of orthodontics and dentofacial orthopedics*. 2006;129(6):722. e1-. e9.
44. Balata PMM, Silva HJ, Nascimento GKO, Moraes KJR, Pernambuco LA, Freitas MCR, et al. Incomplete Swallowing and Retracted Tongue Maneuvers for Electromyographic Signal Normalization of the Extrinsic Muscles of the Larynx. *Journal of Voice*. 2012; 26(6):813e1-813e7. <http://dx.doi.org/10.1016/j.jvoice.2012.03.006>