



Biomédica

ISSN: 0120-4157

biomedica@ins.gov.co

Instituto Nacional de Salud

Colombia

de Vries, Esther; Meneses, María Ximena; Piñeros, Marion
Years of life lost as a measure of cancer burden in Colombia, 1997-2012
Biomédica, vol. 36, núm. 4, 2016, pp. 547-556
Instituto Nacional de Salud
Bogotá, Colombia

Available in: <http://www.redalyc.org/articulo.oa?id=84348515008>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

ORIGINAL ARTICLE

Years of life lost as a measure of cancer burden in Colombia, 1997-2012

Esther de Vries^{1,2}, María Ximena Meneses¹, Marion Piñeros³

¹ Subdirección de Investigaciones, Vigilancia Epidemiológica, Promoción y Prevención, Instituto Nacional de Cancerología, Bogotá, D.C., Colombia

² Departamento de Epidemiología Clínica y Bioestadística, Pontificia Universidad Javeriana, Bogotá, D.C., Colombia

³ Section of Cancer Surveillance, International Agency for Research on Cancer, Lyon, France

Introduction: There is hardly any information regarding years of life lost due to cancer in Colombia.

Objective: To quantify total and average years of life lost due to cancer in Colombia, and to investigate changes in this burden between 1997 and 2012.

Materials and methods: We obtained sex-specific data on age distribution, remaining life expectancy, deaths due to specific cancers and total number of deaths from the Colombian *Departamento Administrativo Nacional de Estadísticas*. We calculated years of life lost based on each individual's age at death and the remaining life expectancy at that age; as for average years of life, we divided these by the number of deaths due to specific cancers.

Results: The total number of cancer deaths increased by 33% between 1997 and 2012, comprising 15% of male and 20% of female deaths in the period 2010-2012. Total years of life lost due to cancer increased by 25.1% for males and 31.1% for females over the study period. The highest increases (>40%) were observed for colorectal, pancreas and kidney cancers in both sexes, for melanomas and bladder cancer in men, and for breast and ovarian cancer in women. Trends in average years of life lost were stable. Almost half (48-50%) of the years of life lost due to childhood cancers were due to leukemia.

Conclusion: Cancer is an increasing health burden in Colombia. The high average years of life lost illustrate the poor prognosis of the disease compared to other countries. Primary prevention, early detection, and adequate and timely treatments are needed to change this situation.

Key words: Neoplasms; potential years of life lost/trends; cost of illness; Colombia.

doi: <http://dx.doi.org/10.7705/biomedica.v36i4.3207>

Años de vida perdidos como medida de la carga de cáncer en Colombia, 1997-2012

Introducción. En Colombia existe escasa información sobre los años de vida perdidos por cáncer.

Objetivo. Determinar los años de vida perdidos por cáncer en Colombia y sus cambios entre 1997 y 2012.

Materiales y métodos. La información por sexo y edad de la mortalidad por cáncer y la expectativa de vida en Colombia se obtuvo del Departamento Administrativo Nacional de Estadística, DANE. El cálculo de los años de vida perdidos por cáncer se basó en la edad de cada individuo al momento de morir y el número de años de vida esperados en esta edad. El promedio de años de vida perdidos se obtuvo dividiendo los años de vida perdidos por el número de muertes a causa de cánceres específicos.

Resultados. Entre 1997 y 2012, el número de muertes por cáncer aumentó 33 %: 15 % en hombres y 20 % en mujeres. En el mismo periodo, el total de años de vida perdidos por cáncer se incrementó en 25,1 % en hombres y 31,1 % en mujeres. Los mayores incrementos (>40 %) se dieron en los cánceres de colon y recto, páncreas y riñón en ambos sexos, en los hombres, en el melanoma y el cáncer de vejiga, y en mujeres, en los cánceres de mama y ovario. El promedio de años de vida perdidos fue estable en el tiempo; 40 a 50 % de los años de vida perdidos por cáncer en niños se debió a leucemias.

Conclusiones. La carga de cáncer está aumentando en Colombia. El elevado promedio de años de vida perdidos demuestra que el pronóstico de la enfermedad es malo. La prevención primaria, la detección temprana, y el tratamiento adecuado y oportuno son necesarios para mejorar esta situación.

Palabras clave: neoplasias; años potenciales de vida perdidos/tendencias; costo de enfermedad; Colombia.

doi: <http://dx.doi.org/10.7705/biomedica.v36i4.3207>

Author's contributions:

Esther de Vries: study idea, design and writing of the manuscript

Marion Piñeros: writing and critical review of manuscript

María Ximena Meneses: critical review of manuscript

All authors analyzed and drafted the section of results and the figures.

From 1984 to 2008, overall cancer mortality rates decreased slightly in both sexes in Colombia (1). Around this same period (1985-2005), substantial demographic changes occurred in the country, including population growth (from 30.8 million to 42.9 million inhabitants) and a rapidly increasing life expectancy: 4.3 years for men and 4.8 years for women (2). Many societal changes also occurred: a large healthcare reform led to greater health insurance coverage, which passed from 59% around year 2000 to 92% in 2012 (3), and security levels increased as violent deaths substantially reduced (4).

This combination of population growth and ageing of the population usually leads to a growing cancer burden, even with relatively stable or slightly decreasing cancer mortality rates (1,5). Although age-standardized mortality rates for cancer as a whole decreased slightly, the total annual number of cancer deaths among males increased between 1984 and 1988 and between 2004 and 2008, from 8,791 to 15,417 (175% increase), and among females from 9,624 to 15,847 (165% increase) (1). This apparent discrepancy between stable or even decreasing mortality rates with increasing number of cancer deaths can be explained by population growth and ageing. Considering only the total number of cancer deaths, irrespective of age and cancer type, overlooks the fact that some cancers have a much heavier burden on society compared to others, because they occur more frequently among younger people.

Years of life lost (YLL) is a useful measure to take the age at death as well as the deaths at each age into account. The measure results from the sum of years of potential life lost by individuals in the population because of death by a certain disease, given their remaining life expectancy estimated by age and sex. The average years of life lost (AYLL) is the average number of years of life lost by each person dying from the disease under study.

Approximately 25-30% of YLL in western countries are attributed to cancers (6). While YLL depend strongly on absolute frequency of deaths, AYLL are mostly determined by the age at which the deaths occur. Together, these measures provide meaningful

indicators for decisions regarding resource allocations and design of prevention programs (7-9). They are also a useful base for comparison with other countries, and show the relative importance of different causes of death, taking into account the life expectancies of each population (10). Previous national and local Colombian studies used external life expectancy data from Japan or the Global Burden of Disease reports, considered as the "potential" life expectancy for other countries (11-13). While using a standard "ideal life expectancy" provides a useful comparison with other countries, a major limitation is that it does not reflect the actual situation of years of life lost in a given country.

The purpose of this paper was to quantify YLL and AYLL due to cancer in general and to the most important cancer types in Colombia for the most recent periods with available data (up to 2012), and to evaluate how this burden changed between 1997 and 2012 using the country's life expectancy data.

Materials and methods

Data and corrections to the data

We used Colombia's official mortality data and population denominators retrieved from the *Departamento Administrativo Nacional de Estadística*, DANE, and we analyzed the following cancer types as listed in the International Classification of Diseases, ICD-10: All cancers (C00-C96), oral cavity and pharynx (C00-C14), oesophagus (C15), stomach (C16), colorectum (C18-C20), pancreas (C25), bronchus and lung (C32-C34), breast (C50), cervix uteri (C53), ovary (C56), prostate (C61), brain and central nervous system (CNS) (C70-C72), leukemia (C91-C95), lymphomas (C81-C85, C96), melanoma (C43), kidney (C64-C65) and bladder and urinary tract cancer (C66-68). We also considered all deaths combined, regardless of the cause of death.

We redistributed deaths without information on age proportionally by sex based on the age distribution of the other cases in the same cause of death category during the corresponding period. Similarly, we redistributed deaths from unspecified uterine cancer (C55) according to the proportions of deaths for cervical cancer (C53) and cancer of the corpus uteri (C54) (14).

Analysis

We calculated annual YLL for the general population (all ages) by period and sex summing up the number of cancer deaths in a certain age group with

Corresponding author:

Esther de Vries, Carrera 21 N° 106B-82, Bogotá, D.C., Colombia
Telefax: (571) 334 1844
estherdevries77@gmail.com

Received: 26/01/16; accepted: 22/04/16

the remaining life expectancy in that age for all age groups according to the method previously explained by Brustugun, *et al.* (6), using the following formula:

$$\sum_{i=0}^{80+} (\# \text{deaths at age } a_i) * (\text{expected remaining life years at age } a_i)$$

To provide an estimate of the YLL due to cancer as a fraction of total YLL, we also calculated a hypothetical total YLL, based on the sum of the products of the total number of deaths per age group, multiplied by the estimated length of life on each age level. We applied this same method for childhood cancers (0- 14 years).

Life expectancy among Colombian population was calculated by age group, sex and time period according to DANE's five-calendar year periods from 1997 to 2005, and from 2005 to 2015 (2). We calculated the number of deaths by cancer type, sex and age groups for the same periods based on the national mortality databases for the 1997-2012 period as provided by DANE (1997-1999; 2000-2004; 2005-2009; 2010-2012).

We calculated the AYLL by dividing the total YLL per cause by the total number of deaths due to this cause, and for each sex and period separately. We determined the changes in YLL and AYLL percentages over time as the percentage increase or decrease between 2010 and 2012 compared to the reference period (1997-1999).

Results

Overall number of cancer deaths

The total annual number of cancer deaths increased for both sexes in about 33%, from 12,292 in 1997-1999 to 16,376 in 2010-2012 in males, and from 12,687 to 17,001 in females. In the first period of the series (1997-1999), cancer comprised 12% of all male and 10% of all female deaths, while in the last period (2010-2012) it accounted for 15% and 20% of male and female deaths, respectively.

Around 3-4% of all deaths occurring in children up to 14 years of age, was due to cancer, and this proportion was stable over time. The total number of cancer deaths in children (0 to 14 years) was 1,749 in the period 1997-1999, which decreased to 1,510 in 2010-2012.

Overall years of life lost and average years of life lost

In 1997-1999, a hypothetical 534,960 (239,626 males; 295,334 females) years of life were lost due to cancer annually, and this figure increased

to 687,540 (299,874 males; 387,666 females) in 2010-2012 representing an overall increase of 28.5%: 25.1% in males and 31.1% in females. In the same period, the Colombian population grew by 24,8% (from 38.6 million in 1997 to 48.2 million in 2012) (2).

Years of life lost and average years of life lost by cancer types

Whereas YLL for any cause of death decreased markedly between 1997 and 2012 (-12% for males, and -54% for females), in this same period they increased for cancer, with 25% in males and 31% in females. For the 2010-2012 period the three most common cancer causes of YLL in men were stomach cancer (49,907), lung cancer (41,871), and leukemia (29,908), while in women they were breast cancer (61,398), cervical cancer (40,165), and stomach cancer (30,210) (figure 1). In this same period, one cancer death in Colombia represented on average 25.7 years of life lost for men and 22.8 for women, though these estimates vary significantly between cancer types.

Virtually, all cancer types showed AYLL of 15 or more years, with the exception of prostate and bladder cancer in males. The highest AYLL in both sexes corresponded to cancers occurring predominantly at young ages: Leukemia (males: 33.8, females: 36.2), cancers of the brain and central nervous system (males: 29.0, females: 30.8) and lymphomas (2010-2012: males 24.5, females 24.6) (figure 2).

We observed the highest increase in YLL for both sexes in colorectal, pancreas and kidney cancers, all by more than 40%. For males, the most substantial increases corresponded to pancreas, melanoma and bladder cancer, while in females, breast, ovarian and colorectal cancers had the largest increases (table 1).

AYLL trends remained fairly stable, except for very rare cancers such as male breast cancer; the few changes observed were generally towards lower AYLL. The largest reductions in AYLL were observed for cancers of the brain and central nervous system, melanomas, lymphomas and leukemia, where two or more years of life expectancy were gained (table 1, figure 2).

Leukemia was the main cause of YLL for children (48-50% of YLL), followed by cancers of the central nervous system (15-16%), lymphomas (6-10%) and kidney cancers (3-4%), which were stable over time; these four cancer types explained 75%




Figure 1. Years of life lost by sex and cancer types, Colombia, 1997-2012.




Figure 2. Average years of life lost to cancer by sex and cancer types, Colombia, 1997-2012

or more of childhood cancer deaths during the study period (figure 3). During this period, YLL for childhood cancers dropped substantially, mainly due to decreases in lymphoma- and leukemia-related YLL. YLL for childhood cancers of the central nervous system and kidney remained rather stable.

The AYLL were very similar for childhood cancer, regardless of the cancer type, varying from 67.0 for colorectal cancer to 72.9 years for kidney cancer (results not shown).

Discussion

According to our results, the contribution of cancer to the total YLL in Colombia (15-20%) was lower than the one reported for western countries (25-

30%) (6). Our estimates were also lower compared to the 2011 and 2015 Colombian studies produced by the *Observatorio Nacional de Salud* (ONS) (12,13) (table 2). Such differences are most likely due to the life expectancy tables of the Global Burden of Disease study used in ONS reports (12,13,15), which show higher life expectancy estimates, and, therefore, a higher number of YLL due to cancer. Although widely used and recommended, the disadvantage of Global Burden of Disease estimates is that they do not reflect the real number of years of life lost in local settings. Using optimal life expectancy gives a hypothetical number of life years lost, whereas in this study we were interested in the reality of the Colombian

Table 1. Change in years of life lost (YLL) and average years of life lost (AYLL) between 1997 and 2012 by cancer type and sex

Cancer types	Percentage of change in YLL, 1997-2012		Change in AYLL, 1997-2012 (years)	
	Males (%)	Females (%)	Males	Females
C00-C14 Head and neck	+ 27	+ 33	- 0.4	+ 1.6
C15 Oesophagus	+ 4	- 6	- 0.5	- 0.5
C16 Gastric	+ 20	+ 22	+ 0.2	+ 1.8
C18-C20 Colorectal	+ 84	+ 76	- 0.9	- 0.2
C25 Pancreas	+ 63	+ 57	- 0.4	- 0.2
C32-C34 Lung	+ 25	+ 44	- 1.1	- 0.9
C43 Melanoma	+ 56	+ 39	- 3.4	- 3.9
C50 Breast	- 52	+ 80	- 4.7	+ 0
C53 Cervix*	-	+ 7		+ 0.7
C56 Ovarian	-	+ 77		+ 0.1
C61 Prostate	+ 39	-	- 0.6	
C64-C65 Kidney	+ 62	+ 48	- 2.1	- 0.2
C67 Bladder	+ 56	+ 38	- 0.5	- 0.5
C70-72 CNS Tumours	+ 11	+ 16	- 1.8	- 2.4
C81-C85 + C96 Lymphoma	+ 28	+ 35	- 4.4	- 4.1
C91-C95 Leukemia	+ 10	+ 16	- 4.9	- 4.8
C 00-96 All cancers	+ 25	+ 31	- 1.2	- 0.5
Total deaths	- 12	- 54	- 5.3	- 12.3




Figure 3. Years of life lost for cancers occurring during childhood (under 15 years of age), 1997-2012

Table 2. Years of life lost by cancer types according to our estimates and estimates by the *Observatorio Nacional de Salud* (ONS) (13)

Cancer types	Our estimates	ONS estimates
Stomach	86,763	100,377
Lung and bronchus	73,446	81,055
Leukemia	58,226	74,152
Breast	61,398	67,425
Colorectal	42,866	61,133
Cervix uteri	51,340	49,296
Prostate	27,507	33,485

population. In addition, the ONS reports used 1998-2011 data, and did not apply redistribution of cases with unknown age or unspecified uterine cancer cases (14) as we did following international recommendations, which could account for the higher YLL in our study.

YLL among the Colombian female population decreased, but the AYLL by cancer type was higher, largely because of women's longer life expectancy (78.5 years at birth in 2010-2012 versus 72.1 for males) (2). This is in line with the latest ONS study reporting neoplasms among the top three causes contributing to most YLL in females alone (12).

Breast and cervical cancer were very important contributors to female cancer mortality. Despite efforts to improve early detection and treatment of these cancers, YLL due to breast cancer increased substantially, with no reductions in AYLL; this indicates increasing number of breast cancer deaths with minimal improvements in age of death, as female life expectancy at birth increased by 3.6 years (from 74.95 in 1997-99 to 78.54 in 2010-12).

Virtually all cancer types showed AYLL of 15 or more years, with the exception of prostate and bladder cancer, which for 2010-12 contributed 11.3 and 13.7 years to the AYLL, respectively. In general, our results for AYLL were high when compared to international reports, which inform of less than 7 years AYLL for colorectal and breast cancer (USA), 1.8 for prostate cancer, and 13.8 for lung cancer (6,7,16). In a Polish study, only male brain cancer and cervical cancer showed more than 10 years of life lost per death (AYLL 10.7 and 10.2, respectively) (16).

Our figures, although not comparable with those from other countries due to differences in overall life expectancy and in the calculating methodologies, evidence the great disparity between estimates (less than 13.5 AYLL in the US compared to more

than 15 years in Colombia, except for prostate and male bladder cancer), and confirm that cancer prognosis in our country is relatively poor. Breast cancer in the USA was responsible for 6.3 years of life lost per death, in Colombia this figure was 25.6 years, illustrating the combined effect of a usually very late diagnosis of breast cancer and/or its more frequent occurrence. Indeed, there is evidence that in Bogotá only 12-30% of women with breast cancer were diagnosed at the early stages, and that delays are substantial both for diagnosis and initiation of treatment (17,18).

Despite a high prostate cancer incidence in Colombia (19), the AYLL due to this cancer was relatively low compared to other cancer types, though high when compared to other countries, which could be attributed to the advanced age at diagnosis, and to its relatively good prognosis.

Leukemia frequently occurs amongst children and has a poor prognosis in Colombia, explaining the very high AYLL encountered for this cancer type. Not surprisingly, all childhood cancer deaths (all cancer types) had an AYLL of around 70 years, because of the very similar remaining life expectancy during childhood.

The YLL of leukemia increased among children and among all age groups, while AYLL of all cancers decreased substantially, indicating more deaths or later ages of death. However, AYLL in childhood leukemia slightly increased (64.6 to 67.8 among girls, 72.0 to 73.9 among boys), possibly due to increases in life expectancy, which indicates that no much progress has been achieved despite the special efforts displayed by the government since 2008 to reduce administrative delays (20-22). On the other hand, it could be attributed to an increase in the number of cases, as population-based survival estimates for childhood cancer show improvements in survival even if it still remains poor in Colombia (20,22,23). Indeed, long-term data from Cali show slight increases in childhood incidence of leukemia and central nervous system cancers (24). We observed no progress at all regarding childhood cancers of the brain and central nervous system, or for kidney cancers. YLL for childhood leukemia were higher than for brain and central nervous system cancers, although this difference is decreasing; it is conceivable that if the actions to improve leukemia prognosis in Colombia are successful, these two other type of cancers will be the main cause of cancer deaths amongst children, similar to the situation observed in the USA (10).

Despite the fact that good methods exist for early detection of breast, cervical, prostate and colorectal cancer, all of these cancer types showed stable or increasing YLL and AYLL, reflecting the great potential for improvements in their early detection in Colombia.

Our AYLL for breast and cervical cancer were much higher (around 25 years) than previous national estimates for 2007 (with 14.4 YLL for breast and 14.6 for cervical cancer), probably due to differences in the life expectancy information used (25). In a study estimating the burden of disease for the Colombian department of Santander, stomach cancer appeared as the first cause of “healthy” life years lost in men, followed by leukemia and colorectal cancer, while for females the first causes were cervical, breast and stomach cancers (12). In contrast, in our study lung cancer was among the first causes in males, while in females the top causes were the same as in Santander. As smoking habits and lung cancer mortality vary heavily within Colombia (smoking prevalence was lower in Santander than the Colombian average) (26,27), regional YLL estimates will vary substantially as well. Moreover, the Santander study reported healthy life expectancy using life tables from Japan as source for remaining life expectancy, which are not comparable to those we used, and this resulted in different life expectancies and, therefore, different YLL.

The observed YLL trends, with more than 40% increase in certain cancer types, are in line with the increasing life expectancy, and, perhaps, the recent efforts to diminish poverty rates in Colombia. This is in line with the fact that cancer types with increasing YLL (breast, colorectal, pancreas, kidney, bladder cancer and melanomas) are mostly related to a more “western” lifestyle and slowly replace the traditionally high burden of infection- and poverty-related cancers, such as gastric and cervical cancers. The increasing proportion of cancer deaths in the total death toll in Colombia is due to population ageing, but also to diminishing number of deaths due to violence (from 0.1% to 0.03% over the study period).

Even though gastric cancer mortality rates have significantly decreased (1), YLL and AYLL have increased, implying that their diagnosis is very late. Anti-tobacco measures should be strengthened to further reduce the burden of lung- and other tobacco-related cancers. Early detection and screening efforts, with a close and direct access to diagnosis and treatment of abnormal results, should be able to reduce breast, cervical and colorectal cancer

mortality substantially. Even though interventions to improve survival in children with leukemia seem to be successful, survival rates are still very poor. In Cali, for example, population-based data from the cancer registry showed rates of only 55% in the most recent diagnostic period (23).

We used national mortality data, generally considered to have acceptable quality and to be among the best within Central and South America (28), particularly after redistributing the cases of unknown age and “uterus, not specified” cases. The quality of the coding in death certificates has improved over time and, therefore, misclassification may have been a larger problem in earlier periods. Indeed, the proportion of deaths with unknown age reduced over time, supporting this hypothesis. However, stable estimates or trends allow us to conclude that misclassification has been relatively minor, and it does not hinder deriving general conclusions such as those arrived at in our study.

Contrary to several previous studies, we did not use an external population to provide the expected remaining years of life at death (life expectancy from life tables), but rather the period and sex-specific life tables provided for Colombia by DANE. The advantage of this method is that it provides the most realistic YLL and AYLL estimates, as they are based on the local real life tables for the corresponding periods. In this respect, we took into account the substantial increases in life expectancy among Colombian population in calculating the trends. However, these improvements in life expectancy may hide other improvements also achieved. If deaths in recent periods occurred at the same ages as before, more potential years of life would be lost and, therefore, the trend would be negative.

Consequently, the stable trends observed may suggest an improvement, which is not apparent because of overall life expectancy improvement. Similarly, YLL increases in this study may not be so much a deterioration as a stabilization of the status quo, due to this increased general life expectancy, while AYLL decreases, mainly in leukemia, lymphomas, cancers of the central nervous system and melanomas (cancers ideally occurring at relatively young ages), actually mean even larger improvements regarding the average age at death. Indeed, in sensitivity analyses using 2010-2012 life expectancy for all periods, AYLL increased in earlier periods (i.e., more years lost given higher life expectancy). Tendencies, however, remained the same (supplementary table, available in: doi: <http://dx.doi.org/10.7705/biomedica.v36i4.3207>)

Together with cancer incidence, mortality and survival statistics, YLL and AYLL provide information on the burden of disease in a population. With increasing life expectancy and population size, the burden of cancer in Colombia is unsurprisingly also increasing. The AYLL was very high for many cancer types, and although comparisons with other countries are difficult given their different population structures, it is clear that primary or secondary prevention, early detection and more timely and adequate treatment can prevent many cancer deaths. The marked increases in YLL observed for colorectal, breast, ovarian and pancreatic cancer are worrisome and call for urgent preventive measures. Melanoma is another largely avoidable cause of death, while the lack of progress in AYLL for childhood leukemia also gives large room for improvement. Given population growth and aging in the country, there is need of more health workers specialized in oncology and preventive medicine.

Conflicts of interest

The authors have no conflicts of interest to declare with respect to this study.

Funding

This project did not receive any external funding. Salaries of the authors were paid for by their respective employers and data was provided free of charge by DANE.

References

1. Piñeros M, Gamboa O, Hernández-Suárez G, Pardo C, Bray F. Patterns and trends in cancer mortality in Colombia 1984-2008. *Cancer Epidemiol*. 2013;37:233-9. <http://dx.doi.org/10.1016/j.canep.2013.02.003>
2. Departamento Administrativo Nacional de Estadística, DANE. Proyecciones de población 2005-2020. Colombia. Tablas de mortalidad nacionales y departamentales, 1985-2020. Date of entry: February 10, 2015. Available from: http://www.dane.gov.co/files/investigaciones/poblacion/proyepobla06_20/8Tablasvida1985_2020.pdf
3. Así vamos en salud. Gráfica - Aseguramiento - Georreferenciado - 2012. Date of entry: November 11, 2015. Available from: <http://www.asivamosensalud.org/inidicadores/aseguramiento/grafica.ver/15>.
4. Dávila-Cervantes C, Pardo-Montañón AM. Magnitud y tendencia de la mortalidad por homicidios en Colombia y México, 2000-2011. *Rev Panam Salud Pública*. 2014;36:10-6.
5. Piñeros M, Hernández G, Bray F. Increasing mortality rates of common malignancies in Colombia: An emerging problem. *Cancer*. 2004;101:2285-92. <http://dx.doi.org/10.1002/cncr.20607>
6. Brustugun OT, Moller B, Helland A. Years of life lost as a measure of cancer burden on a national level. *Br J Cancer*. 2014;111:1014-20. <http://dx.doi.org/10.1038/bjc.2014.364>
7. Carter AJ, Nguyen CN. A comparison of cancer burden and research spending reveals discrepancies in the distribution of research funding. *BMC Public Health*. 2012;12:526. <http://dx.doi.org/10.1186/1471-2458-12-526>
8. Thun MJ, DeLancey JO, Center MM, Jemal A, Ward EM. The global burden of cancer: Priorities for prevention. *Carcinogenesis*. 2010;31:100-10. <http://dx.doi.org/10.1093/carcin/bgp263>
9. De Blank PM, Ostrom QT, Rouse C, Wolinsky Y, Kruchko C, Salcido J, et al. Years of life lived with disease and years of potential life lost in children who die of cancer in the United States, 2009. *Cancer Med*. 2015;4:608-19. <http://dx.doi.org/10.1002/cam4.410>
10. Yang BH, Bray FI, Parkin DM, Sellors JW, Zhang ZF. Cervical cancer as a priority for prevention in different world regions: An evaluation using years of life lost. *Int J Cancer*. 2004;109:418-24. <http://dx.doi.org/10.1002/ijc.11719>
11. Esquiaqui-Felipe R, Posso-Valencia H, Peñaloza RE, Rodríguez-García J. Carga de enfermedad por cáncer en Santander, Colombia, 2005. *Rev Salud Pública*. 2012;14:213-25. <http://dx.doi.org/10.1590/S0124-00642012000200003>
12. Castillo-Rodríguez L, Díaz-Jiménez D, Castañeda-Orjuela C, De la Hoz-Restrepo F. Year of life lost (YLL) in Colombia 1998-2011: Overall and avoidable causes of death analysis. *PLoS One*. 2015;10:1-11. <http://dx.doi.org/10.1371/journal.pone.0125456>
13. Instituto Nacional de Salud, Observatorio Nacional de Salud. Primer Informe ONS, aspectos relacionados con la frecuencia de uso de los servicios de salud, mortalidad y discapacidad en Colombia, 2011. Date of entry: November 3, 2015. Available from: <http://www.ins.gov.co/lineas-de-accion/ons/SiteAssets/Paginas/publicaciones/informe-final%20ONS.pdf>.
14. Loos AH, Bray F, McCarron P, Weiderpass E, Hakama M, Parkin DM. Sheep and goats: Separating cervix and corpus uteri from imprecisely coded uterine cancer deaths, for studies of geographical and temporal variations in mortality. *Eur J Cancer*. 2004;40:2794-803. <http://dx.doi.org/10.1016/j.ejca.2004.09.007>
15. Murray CJ, Ezzati M, Flaxman AD, Lim S, Lozano R, Michaud C, et al. GBD 2010: Design, definitions, and metrics. *Lancet*. 2012;380:2063-6. [http://dx.doi.org/10.1016/S0140-6736\(12\)61899-6](http://dx.doi.org/10.1016/S0140-6736(12)61899-6)
16. Maniecka-Bryla I, Brylo M, Bryla P, Pikala M. The burden of premature mortality in Poland analysed with the use of standard expected years of life lost. *BMC Public Health*. 2015;15:1-8. <http://dx.doi.org/10.18294/sc.2015.725>
17. Piñeros M, Sánchez R, Cendales R, Perry F, Ocampo R. Patient delay among Colombian women with breast cancer. *Salud Pública Mex*. 2009;51:372-80. <http://dx.doi.org/10.1590/S0036-36342009000500004>
18. Piñeros M, Sánchez R, Perry F, García OA, Ocampo R, Cendales R. Delay for diagnosis and treatment of breast cancer in Bogotá, Colombia. *Salud Pública Mex*. 2011;53:478-85. <http://dx.doi.org/10.1590/S0036-36342011000600003>
19. Restrepo JA, Bravo LE, García-Perdomo HA, García LS, Collazos P, Carbonell J. Prostate cancer in Cali, Colombia, 1962-2011: Incidence, mortality and survival. *Salud Pública Mex*. 2014;56:440-7.

20. **Piñeros M, Gamboa O, Suárez A.** Child mortality from cancer in Colombia, 1985-2008. *Rev Panam Salud Pública.* 2011;30:15-21. <http://dx.doi.org/10.1590/S1020-49892011000700003>
21. **Piñeros M, Pardo C, Otero J, Suárez A, Vizcaíno M, Álvarez V, et al.** Protocolo de vigilancia en salud pública de las leucemias agudas pediátricas. *Rev Colomb Cancerol.* 2007;11:219-27.
22. **Ministerio de Salud y Protección Social, Instituto Nacional de Salud.** Protocolo de Vigilancia en Salud Pública. Leucemias - 2014. Date of entry: November 3, 2015. Available from: <http://www.ins.gov.co/lineas-de-accion/Subdireccion-Vigilancia/sivigila/Protocolos%20SIVIGILA/PRO%20Leucemias.pdf>
23. **Allemani C, Weir HK, Carreira H, Harewood R, Spika D, Wang XS, et al.** Global surveillance of cancer survival 1995-2009: Analysis of individual data for 25,676,887 patients from 279 population-based registries in 67 countries (CONCORD-2). *Lancet.* 2015;385:977-1010. [http://dx.doi.org/10.1016/S0140-6736\(14\)62038-9](http://dx.doi.org/10.1016/S0140-6736(14)62038-9)
24. **Bravo LE, García LS, Collazos P, Aristizábal P, Ramírez O.** Descriptive epidemiology of childhood cancer in Cali, Colombia, 1977-2011. *Colomb Med.* 2013;44:155-64.
25. **Agüedo-Botero M, Dávila-Cervantes CA, Atehortúa-Becerra SC.** Diferencias socioeconómicas en los años de vida perdidos por cáncer de mama y cáncer cérvico-uterino en Colombia, 1997 y 2007. *Rev Gerenc Polit Salud.* 2011;10:138-51.
26. **Piñeros M, Pardo C, Gamboa O, Hernández-Suárez G.** Atlas de mortalidad por cáncer en Colombia. Bogotá: Instituto Nacional de Cancerología E.S.E., Instituto Geográfico Agustín Codazzi; 2010.
27. **Ministerio de Salud y Protección Social.** Socialización del informe final de evaluación de necesidades para la ampliación del Convenio Marco de Control del Tabaco. Cifras oficiales para Colombia. 2007. Date of entry: November 3, 2015. Available from: <https://www.minsalud.gov.co/Documents/General/Cifras-tabaco-Colombia.pdf>.
28. **World Health Organization Global Health Observatory data repository.** Census and civil registration coverage. Data by country. Date of entry: April 1, 2016. Available from: <http://apps.who.int/gho/data/node.main>.

Supplementary table 1. Comparison of life expectation in women and men

Women's life expectation in the various periods																		
		C00-14 Head and neck	C15 Oesophagus	C16 Gastric	C18-20 Colorectal	C25 Pancreas	C32-34 Lung	C43 Melanoma	C50 Breast	C53 Cervix	C56 Ovarian	C64-65 Kidney	C67 Bladder	C70-72CNS Tumours	C81-85 + C96 Lymphoma	C91-95 Leukaemia	C 00-96 All cancers	Total deaths
YLL	1997-2000	2952,410	3681,08	30210,42	15804,114	8467,487	21994,758	1632,287	34022,757	40165,177	10700,113	2742,388	1597,357	11373,863	9061,263	24549,691	295334,250	4202427,085
	2000-2005	3560,419	3782,712	33587,034	20168,376	9404,606	25557,096	2175,394	44323,528	44548,978	13617,96	3247,560	1912,842	13545,972	10807,444	26017,197	335795,327	2017080,434
	2005-2010	4034,877	3692,056	36336,364	24742,259	11443,972	29674,828	2470,23	51994,238	44152,482	17115,136	3975,426	2180,66	14653,781	11932,644	26689,107	365498,150	1943224,322
	2010-2015	3912,648	3461,647	36855,47	27838,215	13270	31574,650	2274,747	61398,457	42866,933	18987,063	4058,493	2204,867	13237,183	12271,323	28356,347	387665,861	1917656,786
AYLL	1997-2000	17,511	17,175	19,173	19,818	18,421	19,293	18,907	25,568	26,747	24,981	24,215	16,189	33,138	28,622	41,014	23,279	34,673
	2000-2005	19,309	16,390	19,193	19,938	18,261	18,963	18,916	25,960	27,284	25,426	25,734	15,994	32,563	27,841	39,686	23,216	26,407
	2005-2010	19,308	16,409	20,000	20,093	17,932	18,737	16,082	25,537	27,084	25,110	25,648	15,444	31,898	26,214	37,761	22,899	23,723
	2010-2015	19,087	16,643	20,937	19,660	18,253	18,397	14,998	25,597	27,461	25,071	24,015	15,712	30,760	24,559		36,231	22,342
Women's life expection for 2010-2012 applied to all periods																		
		C00-14 Head and neck	C15 Oesophagus	C16 Gastric	C18-20 Colorectal	C25 Pancreas	C32-34 Lung	C43 Melanoma	C50 Breast	C53 Cervix	C56 Ovarian	C64-65 Kidney	C67 Bladder	C70-72CNS Tumours	C81-85 + C96 Lymphoma	C91-95 Leukaemia	C 00-96 All cancers	Total deaths
YLL	1997-2000	3080,257	3846,503	31626,007	16541,087	8884,89	23110,026	1695,413	35839,79	42302,44	11268,727	2876,706	1667,35	11920,244	9496,522	25598,005	309890,672	4383842,989
	2000-2005	3680,023	3912,95	34720,294	20847,937	9732,628	26451,748	2242,858	45841	46044,092	14078,836	3350,923	1977,26	13955,316	11145,955	26730,212	346920,725	2075219,184
	2005-2010	4098,529	3751,354	36918,446	25140,198	11633,12	30163,248	2506,522	52849,258	44861,006	17395,238	4035,254	2215,2	14867,507	12114,66	27046,902	371290,920	1970501,142
	2010-2015	3912,648	3461,647	36855,47	27838,215	13270	31574,650	2274,747	61398,457	42866,933	18987,063	4058,493	2204,867	13237,183	12271,323	28356,347	387665,861	1917656,786
AYLL	1997-2000	18,269	17,946	20,072	20,743	19,329	20,271	19,638	26,934	28,170	26,308	25,401	16,899	34,730	29,997	42,766	24,427	36,170
	2000-2005	19,958	16,954	19,840	20,610	18,898	19,626	19,503	26,848	28,199	26,286	26,553	16,532	33,547	28,714	40,774	23,985	27,168
	2005-2010	19,613	16,673	20,321	20,416	18,228	19,045	16,319	25,957	27,519	25,521	26,034	15,688	32,363	26,614	38,267	23,262	24,056
	2010-2015	19,087	16,643	20,937	19,660	18,253	18,397	14,998	25,597	27,461	25,071	24,015	15,712	30,760	24,559	36,231	22,803	22,342
Men's life expection for the corresponding periods																		
		C00-14 Head and neck	C15 Oesophagus	C16 Gastric	C18-20 Colorectal	C25 Pancreas	C32-34 Lung	C43 Melanoma	C50 Breast	C61 Prostate	C64-65 Kidney	C67 Bladder	C70-72CNS Tumours	C81-85 + C96 Lymphoma	C91-95 Leukaemia	C 00-96 All cancers	Total deaths	
YLL	1995-1999	4273,537	6145,291	41533,14	12766,285	6761,707	33628,896	1628,82	410,09	19804,58	3274,700	2447,13	13129,057	12565,992	27096,819	239625,691	3288627,668	
	2000-2004	5358,453	6744,064	46183,334	15031,143	7615,678	38456,362	2183,68	253,814	24217,064	3925,463	3103,94	15270,297	14240,310	28653,857	267075,215	3458992,683	
	2005-2009	5251,704	7058,63	48600,182	14050,306	9344,976	43189,548	2609,982	239,884	26354,504	3162,453	3679,796	15907,979	15487,092	29566,173	289478,641	3014243,484	
	2010-2012	5440,443	6408,437	49907,29	23501,472	11015,177	41871,474	2546,567	198,387	27507,463	5295,720	3810,347	14567,503	16045,427	29908,938	299873,804	2894751,430	
AYLL	1995-1999	18,172	15,575	17,656	19,080	17,609	16,907	18,509	21,212	11,824	23,243	14,228	30,834	28,935	38,775	19,495	30,969	
	2000-2004	18,415	15,532	17,949	18,851	17,613	16,756	17,414	17,874	11,588	23,228	14,767	31,190	28,188	37,893	19,280	30,497	
	2005-2009	17,553	15,259	17,719	18,106	17,553	16,297	15,371	16,893	11,241	21,083	14,197	29,657	26,348	35,622	18,598	27,077	
	2010-2012	17,779	15,114	17,830	18,171	17,238	15,832	15,158	16,532	11,267	21,183	13,739	29,019	24,547	33,834	18,312	25,653	
Men's life expectation for 2010-2012 applied to all periods																		
		C00-14 Head and neck	C15 Oesophagus	C16 Gastric	C18-20 Colorectal	C25 Pancreas	C32-34 Lung	C43 Melanoma	C50 Breast	C61 Prostate	C64-65 Kidney	C67 Bladder	C70-72CNS Tumours	C81-85 + C96 Lymphoma	C91-95 Leukaemia	C 00-96 All cancers	Total deaths	
YLL	1995-1999	4383,802	6262,414	42534,93	13123,928	6922,48	34399,786	1674,427	422,213	19952,477	3389,472	2486,983	13709,541	13115,598	28538,670	246811,079	3444225,497	
	2000-2004	5399,167	6772,516	46483,356	15157,023	7659,994	38640,749	2206,52	255,546	24264,466	3975,702	3118,454	15583,659	14511,861	29454,937	270007,268	3541895,992	
	2005-2009	5268,441	7071,77	48743,652	14100,883	9368,844	43292,317	2619,484	240,37	26380,744	3177,432	3686,558	16052,662	15614,165	29936,417	290883,677	3044662,081	
	2010-2012	5440,443	6408,437	49907,29	23501,472	11015,177	41871,474	2546,567	198,387	27507,463	5295,720	3810,347	14567,503	16045,427	29908,938	299873,804	2894751,430	
AYLL	1995-1999	18,640	15,872	18,082	19,615	18,027	17,294	19,028	21,839	11,912	24,058	14,459	32,198	30,201	40,838	20,080	32,435	
	2000-2004	18,555	15,598	18,066	19,009	17,715	16,836	17,596	17,996	11,611	23,525	14,836	31,830	28,725	38,952	19,492	31,228	
	2005-2009	17,609	15,287	17,771	18,171	17,597	16,336	15,427	16,927	11,253	21,183	14,223	29,927	26,564	36,068	18,688	27,350	
	2010-2012	17,779	15,114	17,830	18,171	17,238	15,832	15,158	16,532	11,267	21,183	13,739	29,019	24,547	33,834	18,312	25,653	