



Boletín Latinoamericano y del Caribe de
Plantas Medicinales y Aromáticas

ISSN: 0717-7917

editor.blacpma@usach.cl

Universidad de Santiago de Chile
Chile

HUSSAIN, Mohammad M.; DASTAGIR, Mohammad G.; Masum BILLAH, A.H.M; ISMAIL, Md
Alpinum isoflavone from *Erythrina stricta* Roxb
Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas, vol. 10, núm. 1, enero,
2011, pp. 88-90
Universidad de Santiago de Chile
Santiago, Chile

Available in: <http://www.redalyc.org/articulo.oa?id=85618182011>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative



Alpinum isoflavone from *Erythrina stricta* Roxb

[Alpinum isoflavona de *Erythrina stricta* Roxb]

Mohammad M. HUSSAIN^{1*}, Mohammad G. DASTAGIR², A.H.M Masum BILLAH² & Md ISMAIL²

¹ Department of Pharmacy, Noakhali Science and Technology University, Sonapur, Noakhali-3802, Bangladesh. ² Department of Chemistry, University of Dhaka, Dhaka-1000, Bangladesh.

Contactos / Contacts: E-mail address: Mohammad M. Hussain moshopharma@yahoo.com

Abstract

Alpinum isoflavone was isolated from the n-hexane extract of stem bark of *Erythrina stricta*. The structure of the compound was elucidated by extensive spectroscopic studies and comparison with published spectroscopic data.

Keywords: *Erythrina stricta*, Fabaceae, Alpinum isoflavone

Resumen

A partir de un extracto, obtenido con n-hexano de corteza de *Erythrina stricta*, se aisló la alpinum isoflavona. La estructura del compuesto fue determinada por métodos espectroscópicos y comparación con datos espectroscópicos publicados

Palabras Clave: *Erythrina stricta*, Fabaceae, Alpinum isoflavona

Recibido | Received: December 25, 2010.

Aceptado en versión corregida | Accepted in revised form: January 20, 2011.

Publicado en línea | Published online: January 30, 2011.

Este artículo puede ser citado como / This article must be cited as: Mohammad M. HUSSAIN, Mohammad G. DASTAGIR, A.H.M Masum BILLAH, Md ISMAIL 2011. Alpinum isoflavone from *Erythrina stricta* Roxb. Bol Latinoam Caribe Plant Med Aromat 10(1): 88 – 90.

INTRODUCTION

A medium size deciduous tree with deeply cracked cork bark. Branches armed with white or pale yellow prickles. The genus *Erythrina* belongs to the family Leguminosae and comprises over 100 species of which 8 are known to occur in Bangladesh (Ghani, 2003, Kirtikar *et al*, 1980). Previous investigation of *Erythrina* species have led to the isolation of several phenolic metabolites, such as pterocarpanes, isoflavones, flavonones and chalcones. *Erythrina stricta* has been reported to contain alkaloids, erythline, erysodine, erythrine and other known compounds. *E. stricta* roxb is used for various ailments. Crude extract and elucidated compounds showed antiplasmodial activity, antimicrobial activity and cytotoxic property (Rukachaisirikul *et al*, 2007). As part of our research project on bioactive compounds from Bangladeshi medicinal plants for the treatment of tropical diseases, we have investigated this plant species.

METHODS AND MATERIALS

Collection and preparation of plant material

Plant sample of *Erythrina stricta* was collected from Brahramanbaria in April 2008. A voucher specimen has been deposited in University of Dhaka Herbarium (Herbarium No: 20250). The plant(s) part usually collected in fresh condition. Therefore it should be washed well. It is cut in small pieces and then sun dried followed oven dried at reduced temperature and powdered after drying.

Column chromatography of n-hexane crude extract

The extract (550 mg) was fractionated by column chromatography (Kieselgel 60, mesh 70-230) with a step gradient of *n*-hexane, *n*-hexane-ethyl acetate, ethyl acetate and ethyl acetate-methanol.

Analysis of the column fraction

The fractions were screened by TLC under UV light and by spraying with vanillin-sulphuric acid

reagent. A number of compounds were detected, which were purified from the different sub-fractions employing various techniques. After evaporation of solvent from fraction 66-70, it was subjected to solvent treatment. It was washed with *n*-hexane and then with ethyl acetate in a sample vial. White needle shaped crystals were obtained. It was proved to be pure by TLC.

RESULT AND DISCUSSION

Chromatographic separation and purification of the *n*-hexane soluble fraction of the bark of *Erythrina stricta* afforded one flavones (compound 1). Compound 1 was obtained as needle shaped crystals. It melted at 213 °C, which was identical to that reported for alpinum isoflavone (Martinez *et al*, 1982, Rahman *et al*, 2007, Hernandez *et al*, 2000). It was evident as a yellow spot on TLC (silica gel PF₂₅₄) when the developed plate was sprayed with vanillin-sulphuric acid followed by heating at 110 °C for 5-10 minutes. The R_f value of the compound was 0.533 in toluene-ethyl acetate (90:10) over silica gel PF₂₅₄ plate. It was found to be soluble in ethyl acetate, chloroform, acetone and methanol. The ¹H NMR spectrum of compound-1 (400 MHz, CDCl₃) revealed well resolved signals typical of an isoflavone nucleus having a pyran ring. Thus the ¹H NMR spectrum showed a pair of doublets (*J* = 10.6 Hz) centered at δ 5.53 and 6.60 and a sharp singlet of six proton intensity at δ 1.48. These were assigned to a 2,2-dimethylchromene ring system. The characteristic C-2 proton of the isoflavone skeleton was evident as a singlet at δ 7.83 (1H). The ¹H NMR spectrum also displayed a pair of doublets (*J*=8.5 Hz), each integrating for two protons, at δ 6.96 and 7.27, which were assigned to the H-3' & H-5' and H-2' & H-6' of the *para*-disubstituted aromatic nucleus. The relatively upfield resonance (δ 6.90) of H-3' and H-5' suggested the presence of an oxygenated substituent at C-4'. This was substantiated by the presence of a broad singlet at δ 5.63 (1H), due to a hydroxyl group proton. The remaining two

signals at δ 13.14 and 6.34 (each 1H) could be attributed to the chelated hydroxyl group proton at C-5 and H-18, respectively.

On the basis of the above spectral data and by comparison of these values with those reported for Alpinum isoflavone (Martinez *et al*, 1982, Rahman *et al*, 2007, Hernandez *et al*, 2000). The identity of compound-1 was confirmed as Alpinum isoflavone and this is the first report of its isolation from *Erythrina stricta*.




Figure 1: Compound 1

CONCLUSIONS

The chemical study of the stem bark of *Erythrina stricta* afforded Alpinum isoflavone. The structure was established by extensive spectroscopic studies and comparison with published spectroscopic data (Martinez *et al*, 1982, Rahman *et al*, 2007, Hernandez *et al*, 2000).

ACKNOWLEDGEMENT

The author is highly acknowledged to BCSIR for NMR data for elucidation the structure of the isolated compound.

REFERENCES

- Ghani A. 2003. Medicinal plants of Bangladesh with chemical constituents and uses; Asiatic Society of Bangladesh, Second Edition, p. 222.
- Hernandez NE, Tereschuk ML, Abdala LR. 2000. Antimicrobial activity of flavonoids in medicinal plants from Tadalal Valle (Tucuman, Argentina). J Ethnopharmacol 73: 317 - 322.
- Kirtikar KR, Basu BD. 1980. Indian Medicinal Plants, Published by Singh B and Singh MP, India; 1980, Second edition, Vol-1, pp. 781 - 784.
- Martinez E, Lwande W, Delle Monache F, Marini Bettolo GB. 1982. A Pyranoisoflavone from seeds of *Milletia thonningii*. Phytochemistry 21: 1763 - 1765.
- Rahman MZ, Sultana SJ, Faroque CF, Ferdous F, Rahman MS, Islam MS, Rashid MA. 2007. Phytochemical and biological investigation of *Erythrina variegata*. Saudi Pharm J 15: 140 - 145.
- Rukachaisirikul T, Saekee A, Tharibun S, Watkuolham S, Suksamrarn A. 2007. Biological activities of the chemical constituents of *Erythrina stricta* and *Erythrina subumbrans*; Arch Pharmacol Res 30: 1398 - 1403.