



Boletín Latinoamericano y del Caribe de
Plantas Medicinales y Aromáticas

ISSN: 0717-7917

editor.blacpma@usach.cl

Universidad de Santiago de Chile
Chile

Aponte-Buitrago, Ricardo; Mayorga-Wandurraga, Humberto; Moreno-Murillo, Bárbara
Flavonols and sesquiterpenoids from outer bark and leaves of *Croton polycarpus* Benth.
(Euphorbiaceae)

Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas, vol. 16, núm.
5, septiembre, 2017, pp. 471-485
Universidad de Santiago de Chile
Santiago, Chile

Available in: <http://www.redalyc.org/articulo.oa?id=85652864003>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Artículo Original | Original Article

Flavonols and sesquiterpenoids from outer bark and leaves of *Croton polycarpus* Benth. (Euphorbiaceae)

[Flavonoles y sesquiterpenoides de la corteza y hojas de *Croton polycarpus* Benth. (Euphorbiaceae)]Ricardo Aponte-Buitrago², Humberto Mayorga-Wandurraga¹ & Bárbara Moreno-Murillo¹¹Departamento de Química, Facultad de Ciencias, Universidad Nacional de Colombia, Bogotá, Colombia²Departamento de Química, Facultad de Ciencias, Universidad Antonio Nariño, Bogotá, ColombiaContactos | Contacts: Bárbara MORENO-MURILLO - E-mail address: bmorenom@unal.edu.coContactos | Contacts: Ricardo APONTE-BUITRAGO - E-mail address: ricardoaponte@uan.edu.co

Abstract: Some known flavonols from the outer bark polar fraction of the native species *Croton polycarpus* Benth. (Euphorbiaceae) were isolated and quercetin, quercitrin and rutin were identified as the most abundant constituents. From this sample, a clean TLC non polar sub-fraction (A-1) obtained by CC, afforded several isomeric compounds, which were characterized by GC-MS/EI and its results conduct to a group of eighteen sesquiterpenoids with a unique R_f value in TLC; among them, one sesquiterpene hydrocarbon with molecular weight (MW) of 200 g/mol (C₁₅H₂₀), two with MW 202 g/mol (C₁₅H₂₂), thirteen with a MW of 204 g/mol (C₁₅H₂₄) and two sesquiterpenols with MW 220 g/mol (C₁₅H₂₄O) were determined. From its fresh leaves essential oil obtained by two methods, hydro-distillation and supercritical CO₂ extraction, other terpenoids were identified, where the first one process was more efficient (1.5% DM yield) than the last one, and it was selected for a pilot study. This is the first report about chemical composition of the native species *C. polycarpus*, and the analysis of sesquiterpene isomerism is very important to recognize its structural diversity and similar chemical behavior.

Keywords: Flavonoids, sesquiterpenoids, isomerism, GC-MS, *Croton polycarpus*, Euphorbiaceae.

Resumen: De la fracción polar de la corteza de la especie nativa *Croton polycarpus*, (Euphorbiaceae) se separaron algunos flavonoles y quercetina, quercitrina y rutina se identificaron como los constituyentes más abundantes. Desde una fracción apolar de esta muestra, se obtuvo por CC una fracción limpia en CCD (C-2), cuyo fraccionamiento por análisis por CGAR-EM/IE permitió determinar un grupo de 18 sesquiterpenoides mayoritarios los cuales presentan un valor único de R_f; entre estos se detectó un hidrocarburo de PM 200 (C₁₅H₂₀), dos derivados de PM 202 (C₁₅H₂₂), trece son hidrocarburos isómeros con PM 204 g/mol (C₁₅H₂₄) y dos terpenoles de PM 220 una (C₁₅H₂₄O). De igual forma, de las hojas frescas se extrajo el aceite esencial, por dos métodos: hidrodestilación y extracción con CO₂ supercrítico (SFE), y el primero de ellos fue el más eficiente (1,5% rendimiento en peso seco) por lo cual se seleccionó para el desarrollo de experimentos a escala de planta piloto. Este informe es la primera contribución al conocimiento de la composición química y la bioactividad de las hojas y corteza de la especie arbórea nativa *Croton polycarpus* (Euphorbiaceae) y el análisis de mezclas de sesquiterpenos isómeros es muy importante para reconocer la diversidad estructural y su similitud en polaridad y comportamiento químico.

Palabras clave: Flavonoles, sesquiterpenoides, GC-MS/EI, *Croton polycarpus*, Euphorbiaceae.

Recibido | Received: December 20, 2016

Aceptado | Accepted: February 25, 2017

Aceptado en versión corregida | Accepted in revised form: March 1, 2017

Publicado en línea | Published online: September 30, 2017

Este artículo puede ser citado como / This article must be cited as: R Aponte-Buitrago, H Mayorga-Wandurraga, B Moreno-Murillo. 2017. Flavonols and sesquiterpenoids from outer bark and leaves of *Croton polycarpus* Benth. (Euphorbiaceae). *Bol Latinoam Caribe Plant Med Aromat* 16 (5): 471 – 485.

INTRODUCTION

The Euphorbiaceae family is worldwide distributed and contains trees, shrubbery and herbs. In Colombia, it is represented by 78 genera, 390 species, 12 subspecies and 9 varieties, which has been increased along the last years with the introduction of new species belonging to genera *Mabea*, *Croizatia* and *Pausandra*. Within the most representative genera, it is *Euphorbia* with 43 species (spp.), *Croton* (80 spp.), *Phyllanthus* (36 spp.), *Acalypha* (25 spp.), *Alchornea* (19 spp.), *Mabea* (18 spp.) and others that present less than 11 species (Murillo, 1999; Murillo, 2004).

Chemical composition and biological activity in Euphorbiaceae have gotten some advances related mainly with *Euphorbia*, *Alchornea* and *Croton* taxa.

Regarding *Croton* genus, there are extensive reports about bioactivity, chemical composition, their biological properties and bioactive constituents; in this respect, it is worth mentioning several studies about the product known as “Sangre de Drago” which is a red resin with anti-cancer, anti-inflammatory, anti-oxidant and bactericidal activities, and it is obtained from numerous *Croton* species widely distributed in South America (Ramírez, 2003). This genus offers a great amount and variety of constituents like phorbol esters, alkaloids, di and triterpenoids as clerodanes, euphol derivatives, and flavonoids and their glycosides with notable medicinal applications, as summary in Table 1.

Table 1
Chemical composition and biological activity review of *Croton* genus.

Species	Place	Part	Compound isolated	Bioactivity	Reference
<i>C. brasiliensis</i>	Alagoas, Brasil	Stem and leaves	Dit. as crotobrasilin A and B	Not known reports.	Palmeira Jr. <i>et al.</i> , 2005
			Flav. as casticin, penduletin & crisospenol-D		
<i>C bogotanus</i>	Bogotá, Colombia	Leaves	Dit. as cassipourol		Vegas <i>et al.</i> , 2011
<i>C. cajucara</i>	Jacundá, Brasil	Bark	Cler. as <i>t</i> -cajucarine B. sacacarine. <i>t</i> -crotonine, cajucarinolide.	Antimicrobial, antiviral.	Aparecida <i>et al.</i> , 2002
			Trit. as acetilaleuritolic acid.		
		Leaves	Flav. as methylkaempferol.	Diabetes. cholesterolemia	Campos <i>et al.</i> , 2002
			Ster. as β -sitosterol & stigmasterol.		
	Bark and stem	Cler. as <i>trans</i> -crotonine		Perazzo <i>et al.</i> , 2007	
	Benfica, Brasil	Bark	Dit. as dehydrocrotonine	Antiulcer.	Silva <i>et al.</i> , 2003
<i>C. californicus</i>	Arizona, USA	Bark	Dit. as methyl-barbascoate. Éster 12-deoxiphorbol.	Tumor promoter.	Chavez <i>et al.</i> , 1982
<i>C. campestris</i>	Sao Paulo, Brasil	Bark and roots	Cler. as velamone & velamolone.	hepatic diseases & syphilis	Babili <i>et al.</i> , 1998
				Leishmaniasis	Babili <i>et al.</i> , 2006
<i>C. caudatus</i>	Yunnan, China	Stem	Sesq. as crocaudatol & oplopanone	Anticancer	Wang & Zou, 2008
			Flav. as Crotoncaudatin, tangeretin, nobiletin & sinensetin	Antimalaric and fever.	Zou <i>et al.</i> , 2010
<i>C. celtidifolius</i>	Santa Catarina, Brasil	Bark	Proantocianidin 63SF	Pain and wounds.	DalBó <i>et al.</i> , 2006
			galocatequin & catequin	Leukemia	Nardi <i>et al.</i> , 2003
	Orleans City, Brasil			Ster. as β -sitosterol.	Antiinflammatory
<i>C. cuneatus</i>	Barinas, Venezuela	Bark and stem	Terp. as α -11 eudesmen, julocrotonine & julocrotol	Analgesic and antiinflammatory	Suárez <i>et al.</i> , 2006
<i>C. hovarum</i>	Ankazobe, Madagascar	Bark	Trit. as β -amirine	Toxic compounds.	Krebs & Ramiarantsoa, 1996
			Trit. as 4-hydroxyhigrinic acid		

<i>C. kongensis</i>	Yunnan-Tibet, China	Leaves	Dit. as kongensin & rabdoubrosain 4-hidroxi benzoic acid	Antimalaric	Yang et al., 2009
<i>C. lechleri</i>	Cerro de Pasco, Perú	Latex	Lign. as 3'4-O-dimetilcedrusin. catequin, epicatequin & galocatequin. Proantocianidin SP-303.	Antiinflammatory antiviral and antitumor.	Coussio et al., 1997
			taspin & apomorfina	Antioxidant and anticancer	Lopes e Lopes et al., 2004
<i>C. malambo</i>	Santa Bárbara del Zulia, Venezuela	Bark	β -pinene, α -bergamotene, β -metil-isoeugenol, linalool	Analgesic and antiinflammatory	Suárez et al., 2003
	Puerto Colombia, Atlántico			Citotóxic and antiinflammatory	Jaramillo et al., 2007
<i>C. megalocarpus</i>	Arusha, Tanzania	Essential oil	Fatty acids C16 and C18	Biodiesel.	Kafuku et al., 2010
<i>C. membranaceus</i>	Krobo-Gyakiti, Ghana	Roots	Alk as julocrotin	Not known reports	Aboagye et al., 2000
<i>C. micradenus</i>	Guantánamo, Cuba	Stem and leaves	Alk as sinoacutin, cobotrin & soboldin	Antimalaric	Payo et al., 2001
<i>C. oblongifolius</i>	Petchaboon, Tailandia	Bark	Cemb. as crotoembraneic acid.	Dyspepsia treatment.	Roengsumran et al., 1998
<i>C. pullei</i>	Pará, Brasil	Stem and bark	Alk as julocrotin, crotonimide A and B	Not known reports	Bárbara et al., 2007
<i>C. schiedeana</i>	Tocaima, Colombia	Stem and leaves	Phenbut. as acetoxi & diacetoxirododendrol		Puebla et al., 2005
<i>C. sonderianus</i>	Ceara, Brasil	Roots	Cler. as sonderianin	Antifungal, antibacterial.	McChesney et al., 1991
<i>C. sublyratus</i>	Bangkok, Tailandia	Leaves	geranylgeraniol. & plaunotol.	Antibacterial against <i>H. pylori</i> .	Tansakul & De Eknankul, 1998
<i>C. tonkinensis</i>	Vietnam	Leaves	Flav. as vitexine, isovitexine & tiliroside	Antioxidant, antitumor and antiinflammatory.	Giang et al., 2004
<i>C. zambesicus</i>	Yaounde, Camerún	Bark	Dit. as crotonadiol. & crotochryliferane.	Laxative.	Ngadjuli & Folefoc, 1999
			Trit. as lupeol. betusinol.		
			Ster. as 3 β -glucopiranosil-sitosterol		
	Benin, África	Leaves	Dit. as ent-18-hidroxitraquilobane	Hypertension	Martinsen et al., 2010
<i>C. zehntneri</i>	Pernambuco, Brasil	Roots	crototropone	Antioxidant & antiinflammatory.	Bracher et al., 2008
	Ceará, Brasil	Leaves	Sesq. as trans-anetol & trans-cariofililene	Sedative.	Rodrigues et al., 2009

Dit: Diterpenoid; **Trit:** Triterpenoid; **Sesq:** Sesquiterpenoid; **Alk:** Alkaloid; **Ster:** Steroids; **Flav:** Flavonoids; **Cemb:** Cembranoids; **Terp:** Terpenoid; **Lign:** Lignane; **Phenbut:** Phenilbutanoid; **Cler:** Clerodane.

In Colombia, previous works include a report of two constituents of non-polar fractions from *Alchornea glandulosa*, where the presence of 3-hidroxi-friedelin and 3-oxo-friedelin were determined, whose

bioactivity in Brine Shrimp Test (BST) was EC₅₀= 31 μ g/mL for the first compound and EC₅₀= 43 μ g/mL for the last one (Tello-Camacho, 2005). Also for this

plant the biological activity of phenolic compounds isolated, was assessed (Urrea-Bulla et al., 2004).

From genus *Croton* some terpenoids have been obtained like linalool, germacrene D, *trans*- β -caryophyllene, and others from volatile fractions of *C. malambo* (Jaramillo et al., 2010) where eugenol and methyl-eugenol were the main constituents with potential cytotoxic and repellent activities (Jaramillo et al., 2007; Muñoz-Acevedo et al., 2014; Mendoza-Meza et al., 2014). Likewise, it was found *trans*-calamenene, dihydrocurcumene and β -caryophyllene as more abundant constituents of essential oil (EO) of *C. trinitatis* (Jaramillo-Colorado et al., 2016) and cassipourol, which is a diterpenic alcohol with potential larvicidal activity, isolated from a non-polar fraction of *C. funckianus* senescent leaves. From the polar fraction of *C. funckianus*, some phenolic compounds were isolated and they were elucidated as quercetin and quercitrin, were the first one gave a promissory larvicidal activity [$CE_{50} = 47 \mu\text{g/mL}$] against mosquito larvae of *Culex quinquefasciatus* Say (Diptera: Culicidae), and the last one was characterized as a flavonol glycoside without relevant biological activity (Vegas-Mendoza, 2010; Vegas et al., 2011).

At the best of our knowledge, there are not previous reports about *Croton polycarpus* Benth. [Basiosym. *Oxidectes polycarpa* (Benth.) Kuntze] chemical composition or bioactivity, so this is the first contribution to its chemical analysis. The main goal of this research was to study the compounds of native species *C. polycarpus* grown on the Bogotá highlands. From its outer bark and fresh leaves by chromatographic methods as TLC, HRGC-MS/EI the essential oil (EO) chemical composition was analyzed; the EO was obtained by two methods: hydro-distillation and CO_2 supercritical extraction, with subsequent application of analytical chromatographic methods in order to select the best for larger scale experiments.

MATERIALS AND METHODS

Plant material

Bark trunks and fresh leaves of *C. polycarpus* were collected in August 2010 at the Universidad Nacional de Colombia campus located in Bogotá (UNC-SB), at 2600 AMSL, where a voucher specimen under the code COL 520458 is deposited at the Herbario Nacional Colombiano (HNC) in the Instituto de Ciencias Naturales (ICN) – UNC- Bogotá.

Extraction and Isolation

A *C. polycarpus* sample was separated manually in outer bark, wood and senescent leaves and then were dried and powered for subsequent solvent extraction. Firstly, a bark sample (720 g) was extracted with EtOH (96%) at RT by percolation, changing the solvent thoroughly. Filtered extract (F-1) was concentrated at reduced pressure (RP) at maximum 40°C in a rotatory evaporator (Heidolph WB2000). A portion of the extract (10 g) was fractionated by L-L partition between water and petroleum ether (F-2), CHCl_3 (F-3), AcOEt (F-4) *n*-BuOH (F-5) and aqueous residue (F-6) were obtained. After application of general BST with *Artemia salina nauplii* and specific larvicidal activity bioassay (LAB) with mosquito *Culex quinquefasciatus* larvae, a progressive chromatographic analysis by TLC was developed (McLaughlin et al., 1998).

According TLC results, fraction C-2 (1.12 g) was fractionated in a silica gel (70-230 mesh) column (PE/AcOEt 100:0 $\rightarrow$ 0:100 and then AcOEt/MeOH 90:10 $\rightarrow$ 10:90); 25 fractions eluted with PE/AcOEt 95:5 (96 mg) were collected, analyzed by TLC in hex/tol 50:50 and an unique spot ($R_f = 60$) was observed with acid reagent $\text{H}_2\text{SO}_4:\text{H}_2\text{CO}:\text{W}$ (2:1:7) and heating at 120°C (H^+/Δ); a sample of $1 \mu\text{L}$ solution was subjected to CG-MS/EI with splitless injection method and lineal velocity of 37 cm/sec. The oven heating ramp program was 50°C (2 minutes), increasing until 200°C ($4^\circ\text{C}/\text{min}$) and then up to 250°C ($10^\circ\text{C}/\text{min}$).

Fresh leaves sample (500 g) was subjected to hydrodistillation at ATM (3 L/ H_2O , 3 h) in a Clevenger-type apparatus, where the volatile oil was isolated by L-L partition with Et_2O and then dried over anhydrous Na_2SO_4 (yield 1.5%). After filtration, it was kept in amber-colored bottle at RT. Other leaves sample (25 g) was submitted for supercritical CO_2 extraction (1400 psi, 43°C , $0.3 \text{ g}/\text{cm}^3$ gas density) for 60 minutes (yield 0.1%). According these results a sample of EO obtained by HD was selected for CG-MS/EI with a same heating ramp program previously proposed.

Flavonols and their glycosides were isolated from the raw fraction in AcOEt (1.1 g), by successive CC (silica gel 70-230 mesh) with mixtures of toluene/AcOEt (100:0 $\rightarrow$ 0:100) and AcOEt/MeOH (100:0 $\rightarrow$ 0:100), where it was collected F-7 (10 mg), F-8 (13 mg) and F-9 (15 mg) fractions. The ^1H NMR (400 MHz) spectra afforded enough information to characterize the compounds as the known flavonol

quercetin and their glycosylated derivatives quercitrin (quercetin-3-O- α -L-rhamnoside) and rutin (quercetin-3-rutinoside) by comparison with previously reported values (Fathiazad *et al.*, 2006).

General chromatographic conditions

Analysis by HRGC-MS for C-2 sample was performed on a Shimadzu QP2010 coupled with MS-EI detector QP-2010 S (70 eV) and an ion source temperature of 230°C, using a column packed with 5% phenil-95% dimethylpolysiloxane S-5 (30 m $\times$ 0.25 mm $\times$ 0.25 μ m). HRGC-MS for essential oils was analyzed on a HP6890 Series II with HP-5973N MS detector (e.i. 70 eV) at 230°C, equipped with a fused silica capillary column HP-5MS (30 m $\times$ 0.25 mm $\times$ 0.25 μ m). In the same conditions, a mixture of linear hydrocarbons from C-8 until C-20 was run. The identification of compounds was made with application of two main criteria: the chromatographic one comparing data with retention indexes (KI) calculated based on the Kovats system related with the reference lineal paraffin's (C₈ to C₂₀) mixture, as

well as with the spectroscopic criteria based on the grade of coincidence of the mass spectra registered with the reported in NIST08 and Wiley Registry 8e GC-MS libraries. ¹H NMR and ¹³C spectra were registered in a Bruker MX-400 at 400 MHz for ¹H and 100 MHz for ¹³C, using MeOHd₄ and Me₂COd₆ as dissolvent and TMS as internal standard.

RESULTS AND DISCUSSION

From BST results of EP-outer bark and AcOEt senescent leaves extracts obtained from *C. polycarpus*, under the protocol proposed by McLaughlin *et al.* (1998) it was found that these present promised bioactivity in outer bark, whose values are 58 μ g/mL and 198 μ g/mL (Table 2). Otherwise, senescent leaves present better bioactivity for its AcOiPr, *n*-BuOH and PE extracts with specific larvicidal activity bioassay (LAB) using *Culex quinquefasciatus* larvae. Based on these reports, the study of its chemical composition was conducted and the results agree with data coming from other *Croton* species (Vegas *et al.*, 2014).

Table 2
BST and LAB results as EC₅₀ for *C. polycarpus* of outer bark and senescent leaves extracts

Extracts	BST Bark (μ g/mL) EC ₅₀	LAB
		Senescent leaves (μ g/mL) EC ₅₀
F-1	315	105
F-2	198	549
F-3	229	>1000
F-4	149	---
F-5	---	494
F-6	---	>1000
F-7 (AcOiPr)	---	451
Caffeine	42	450

Fractionation of constituents of outer bark F-2 extract were made over a silica gel column (70-230 mesh) and led to C-2 clean fraction, whose BST EC₅₀ value was 58 μ g/mL, and shows a unique spot with R_f=60. Its chemical composition was studied by HRGC-MS, where more abundant compounds are a set of isomeric sesquiterpenoids according the profile observed in Figure 1.

Figure 1 records the chromatographic profile of sample C-2 coming from HRGC analysis joint with its respective enlargement in the range of 20 to 35 minutes of elution. The chromatogram allows to differentiate the Retention Time (R_T) of at least

eighteen compounds, most of them with MW varying between 200 and 220 uma corresponding to sesquiterpenoid derivatives like sixteen hydrocarbons and two sesquiterpenols that by application of the chromatographic and spectroscopic criteria have conducted to detection of their identities complemented by fragmentography analysis. Sesquiterpene hydrocarbons with two or more rings and exocyclic double bonds exhibit spectra consistent with isomerization to cyclohexene derivatives that can suffer other rearrangements and afford aromatic fragments as seen in Figure 3 (Mc Lafferty & Tureček, 1993). The mass spectra analysis conducted

to the identification of following known terpenoids: α -copaene [1], β -caryophyllene [2], aromadendrene [3], δ -cadinene [4], α -calacorene [5], δ -cadinol [6],

9,10-dehydro-isolongifolene [7] and (+)-cuparene [8] as most abundant constituents (Figure 2).




Figure 1
Total and enlargement chromatograms of C-2 fraction







Figure 2
Structures of main compounds from outer bark C-2 fraction.

Table 3 shows the list of compounds identified by spectroscopic analysis of C-2-fraction which is mainly formed by thirteen isomeric sesquiterpenoid hydrocarbons of MW of 204 g/mol, most of them as decaline derivatives. On the other hand, there were found two other hydrocarbons of 202 g/mol known as (-)-calamenene and (+)-cuparene, two sesquiterpenols with 222 g/mol named δ -cadinol and germacrene-D-4-ol and finally α -calacorene with a MW of 200 g/mol. Application of chromatographic and spectroscopic criteria, to each

component, for example, to germacrene D [4] and α -bergamotene [5] show that both retention index with values of 1474 and 1476 are similar and present differences with literature reports based on different experimental conditions although their mass spectrum present a molecular ion of $[M^+] = 204$ uma; however, by comparative analysis the first one presents a base peak of $m/e = 161$ uma, while the last one has it at $m/e = 93$ uma, this led to confirm their structures.



[4]



[5]

Figure 3
MS spectra of [4] and [5] compounds with respective fragmentography for germacrene D

Table 3
Chemical composition and Area% from *C. polycarpus* C-2 fraction

Peak	<i>I_R</i> lit.	<i>I_R</i> exp.	Compound name	Formula	MW	Area%	Base Peak MS	Id. Criteria	<i>I_R</i> theor. Ref.
1	1364	1447	(+)-cicloisositivene	C ₁₅ H ₂₄	204	1.05	105	MS	a
2	1376	1451	α-copaene	C ₁₅ H ₂₄	204	5.79	105	MS	a
3	1418	1470	β-caryophyllene	C ₁₅ H ₂₄	204	10.79	93	MS	a
4	1480	1474	germacrene-D	C ₁₅ H ₂₄	204	0.56	161	MS	a
5	1436	1476	α-bergamotene	C ₁₅ H ₂₄	204	1.63	93	MS	a
6	1454	1484	α-humulene	C ₁₅ H ₂₄	204	4.56	93	MS	a
7	--	1487	unknown 1	C ₁₅ H ₂₄	204	6.45	91	MS	a
8	1438	1492	aromadendrene	C ₁₅ H ₂₄	204	6.29	161	MS	a
9	1377	1494	isodene	C ₁₅ H ₂₄	204	1.01	105	MS	b
10	1418	1495	(-)-β-caryophyllene	C ₁₅ H ₂₄	204	0.88	69	MS	a
11	1511	1498	germacren-D-4-ol	C ₁₅ H ₂₆ O	222	3.52	161	<i>I_R</i> , MS	c
12	1499	1501	α-murolene	C ₁₅ H ₂₄	204	0.48	105	<i>I_R</i> , MS	a
13	1524	1509	δ-cadinene	C ₁₅ H ₂₄	204	11.26	161	<i>I_R</i> , MS	a
14	1532	1512	cadin-1,4-diene	C ₁₅ H ₂₄	204	0.80	119	<i>I_R</i> , MS	a
15	1542	1518	α-calacorene	C ₁₅ H ₂₀	200	8.02	157	<i>I_R</i> , MS	a
16	1636	1523	δ-cadinol	C ₁₅ H ₂₆ O	222	3.71	161	MS	d
17	1531	1539	(-)-calamenene	C ₁₅ H ₂₂	202	5.14	159	<i>I_R</i> , MS	a
18	1513	1548	(+)-cuparene	C ₁₅ H ₂₂	202	8.99	132	<i>I_R</i> , MS	b

Mass spectra of unknown compound, m/z (int. %): 91(100), 105(98), 161(70), 133(70), 119(68), 79(68), 41(62), 67(52), 147(52), 55(40), 189(25). ^aPino et al., 2006; ^bElías et al., 1997; ^cHamm et al., 2004; ^dHamm et al., 2005.

Fragmentography and ten higher peaks intensity in MS analysis for each compound (Table 3) was applied and allowed to identify eighteen isomeric compounds. However, a detailed comparison between compounds [4] and [5] leads to propose that while its *I_R* has a little difference they present spectroscopic data that justify its structural diversity despite having the same molecular ion at *m/e* = 204 uma (Figure 3).

Analysis by HRCG-MS/EI of the EO led to identification of monoterpenes such as α-thujene, α and β-pinene, β-phellandrene, a monoterpenol known as β-linalool, some esters as bornil acetate, *trans* - pinocarvyl acetate and some sesquiterpenols among which are α-cadinol, α-bourbonenol and caryophyllene oxide; nevertheless, despite the great variety of compounds, EO is constituted by hydrocarbons (56.5%) and alcohols (24.7%) mainly (Table 4).

Table 4
Constituents of *C. polycarpus* leave essential oil by hydrodistillation

R _T	<i>I_R</i> theor.	<i>I_R</i> exp.	Compound name	MW	Area%	Base Peak	Id. criteria	<i>I_R</i> theor. Ref.
3.67	930	902	α-thujene	136	1.59	93	<i>I_R</i> , MS	a
3.81	940	948	α-pinene	136	15.9	93	<i>I_R</i> , MS	a
4.09	953	943	camphene	136	0.27	93	<i>I_R</i> , MS	a
4.56	1053	964	β-phellandrene	136	0.88	93	MS	b
4.64	980	943	β-pinene	136	10.75	93	<i>I_R</i> , MS	a
4.89	994	958	β-myrcene	136	0.49	41	<i>I_R</i> , MS	c
5.51	1017	998	α-terpinene	136	0.35	121	<i>I_R</i> , MS	a
5.71	1010	1042	m-cymene	134	2.82	119	<i>I_R</i> , MS	d
5.80	1020	1018	δ-limonene	136	0.78	68	<i>I_R</i> , MS	b
5.87	1039	1059	eucalyptol	154	1.53	43	<i>I_R</i> , MS	b

6.57	1062	998	γ -terpinene	136	5.01	93	MS	a
6.83	1189	1158	cis- α -terpineol	154	0.73	43	I_R , MS	e
7.73	1097	1081	β -linalool	154	3.20	71	I_R , MS	e
8.37	1141	1109	trans-2-menthenol	154	0.21	43	I_R , MS	f
8.89	1139	1131	L-trans-pinocarveol	152	0.76	92	I_R , MS	a
9.03	1132	1121	camphor	152	0.40	95	I_R , MS	b
9.07	1131	1136	cis-verbenol	152	0.66	94	I_R , MS	b
9.67	1228	1228	nerol	154	0.30	69	I_R , MS	a
10.05	1162	1137	terpinen-4-ol	154	0.72	71	MS	d
10.49	1172	1143	α -terpineol	154	0.70	59	MS	d
10.91	1265	1380	4-(1,2-dimethyl-cyclopent-2-enyl)-butan-2-one	170	0.64	95	MS	g
11.13	1227	1131	trans-3(10)caren-2-ol	152	0.56	109	MS	h
11.82	1234	1231	o-methylthymol	164	0.30	149	I_R , MS	a
11.96	1001	948	2-carene	136	1.13	93	I_R , MS	i
13.43	1275	1277	bornyl acetate	196	1.50	95	I_R , MS	c
14.11	---	1060	unknown 1	140	0.32	55	---	---
15.44	1086	1333	p-mentha-2,4(8)-diene	196	0.30	43	MS	j
15.95	---	1140	unknown 2	204	0.26	105	---	---
16.24	1377	1221	α -copaene	204	3.21	105	MS	e
16.52	1388	1344	β -bourbonene	204	0.47	81	I_R , MS	c, e
16.69	1351	1339	β -cubebene	204	0.33	161	I_R , MS	j
17.58	1418	1494	β -caryophyllene	204	6.44	93	MS	a, e, i
18.63	---	1579	unknown 3	204	0.75	93	---	---
18.84	1460	1386	L-alloaromadendrene	204	0.52	105	MS	a
19.34	1480	1515	germacrene D	204	2.64	161	I_R , MS	a
19.92	1511	1660	germacren-D-4-ol	222	1.32	161	MS	c
20.74	1524	1469	δ -cadinene	204	0.96	161	MS	a, c
21.32	1542	1547	α -calacorene	200	0.49	157	I_R , MS	a
21.93	1466	1410	dehydroaromadendrene	220	0.94	119	MS	k
22.18	1654	1580	α -cadinol	222	3.38	43	MS	a, e
22.36	1576	1569	sphathulenol	220	1.06	43	I_R , MS	a
22.47	1583	1507	caryophyllene oxide	220	2.90	43	MS	h
23.16	1604	1543	cedrenol	222	2.26	95	I_R , MS	l
23.48	1524	1537	L-calamenene	202	0.56	159	I_R , MS	c
23.71	No report	1387	α -vetivone	218	0.79	105	MS	---
24.31	---	1580	unknown 4	222	0.61	161	---	---
24.93	No report	1547	α -calacorenol	216	0.75	157	MS	---
25.21	No report	1646	ledene oxide (II)	220	3.57	43	MS	---
25.43	---	---	unknown 5	220	0.30	91	---	---
26.02	No report	1685	cycloisolongifol-8-ol	220	7.83	159	MS	---
Main constituents of Essential Oil				%				
Total				95.1				
Hydrocarbons				56.5				
Alcohols				24.7				
Esters				1.5				
Ketones				1.8				
Ethers				8.0				
Unknown				2.5				

^aPino et al., 2006; ^bHögnadóttir & Rouseff, 2003; ^cHamm et al., 2004; ^dCavalli et al., 2003; ^eNúñez-Arevalo et al., 2010; ^fVardar-Ünlü et al., 2003; ^gShivashankar et al., 2012; ^hCampeol et al., 2001; ⁱKarioti et al., 2003; ^jTéllez et al., 2001; ^kElías et al., 1997; ^lChoi, 2005.

The essential oil senescent leaves of *C. polycarpus* obtained by hydro distillation afforded 1.5% yield, and regarding to CO₂ supercritical extraction (SFE) was 0.1% with similar chemical composition but lower amount; its comparative analysis shows that HD is more efficient and cheap; the EO was analyzed by HRCG – MS ESI and compared in its constituents with C-2 non-polar fraction obtained by chromatographic methods (Blanco & Caicedo, 2010).

Besides, Tables 3 and 4 show that β -caryophyllene is the most abundant compound in both extracts as well as in other EO *Croton* species grown in Colombia such as *C. trinitatis* (Jaramillo-Colorado et al., 2016), *C. malambo* (Jaramillo et al., 2010), *C. bogotanus* (Núñez-Arévalo et al., 2010) has been analyzed. Several EO reports from other countries such as *C. flavens* (Sylvestre et al., 2006), *C. urucurana* (Simionatto et al., 2007), *C. cajucara* (Lopes et al., 2000), *C. micradenus* and *C. myricaefolius* (Pino et al., 2005; Pino et al., 2006) are reported in Table 5, which shows some common

components besides to β -caryophyllene and its oxide or α and β pinene, more abundant compounds in these species.

HRGC-MS analysis of EO shows that fresh leaves chemical composition of *C. polycarpus* are rich in monoterpenes like α -pinene (15.9%) and β -pinene (10.7%), sesquiterpenes such as β -caryophyllene (6.4%) and germacrene D (2.6%), sesquiterpenols as α -cadinol (3.4%), cedrenol (2.3%), cycloisolongifol-8-ol [9] (7.9%) and oxides such as caryophyllene oxide and ledene oxide II [10], with 2.9% and 3.6% respectively. Other alcohols are the 24.7% of EO. In Table 5, α - and β -pinene were the most abundant monoterpenes, which are common in most of analyzed species; however, their values are larger in *C. polycarpus* than in other species. Among most prevalent sesquiterpenes of the *Croton* EO are: α -copaene [1], β -bourbonene [11], β -caryophyllene [2], germacrene D [12], δ -cadinene [4] sesquiterpenols such as sphathulenol [13] and caryophyllene oxide [14] (Figure 4).

Table 5
Main constituents of essential oils from *Croton* species in comparison with *C. polycarpus*.

Compound name	Cp	Ct	Cm	Cb	Cf	Cu	Cc	Cmc	Cmy
α -thujene	1.59				0.27	0.1	<0.1		
α -pinene	15.9	0.4	0.4	3.2			0.6	5.3	3.9
camphene	0.27			<0.1			0.1	2.0	2.6
β -pinene	10.75	0.2	1.0	0.3			0.1	1.1	2.0
β -myrcene	0.49			0.6		0.2	<0.1	2.1	0.2
α -terpinene	0.35	0.4				0.1		<0.1	<0.1
δ -limonene	0.78	0.3		55.2		0.2	1.1	4.5	
eucalyptol	1.53	2.4				1.8	1.1		4.3
γ -terpinene	5.01				0.54	0.3	<0.1		0.1
β -linalool	3.20		5.6	0.7		0.3	41.2	34.9	1.2
<i>trans</i> -2-menthenol	0.21		0.2			0.1			
camphor	0.40	0.2							
α -terpineol	0.70			0.1		1.2		1.2	3.8
bornyl acetate	1.50		0.5			5.9	<0.1	3.3	
α -copaene	3.21		1.4	0.2	0.50	0.1	0.5		0.2
β -bourbonene	0.47	0.2	0.4	0.2	0.46		2.0		
β -caryophyllene	6.44	15.3	2.5	1.3	4.95		6.9	2.4	0.8
L-alloaromadendrene	0.52		0.6		1.21		0.4		
germacrene D	2.64	0.7	1.6	1.1	2.45		4.0	0.1	
germacren-D-4-ol	1.32				3.97				
δ -cadinene	0.96		0.6	0.3	2.31	0.8		0.2	0.2
α -calacorene	0.49					0.1		0.2	
dehydroaromadendrene	0.94				1.59				

α -cadinol	3.38	1.1		0.4	3.97			0.1	
spathulenol	1.06	1.9	0.1			0.3	2.0		
caryophyllene oxide	2.90	2.5	0.6	0.2	3.55	0.4	1.8	2.8	8.8

Cp: *C. polycarpus*; **Ct:** *C. trinitatis*; **Cm:** *C. malambo*; **Cb:** *C. bogotanus*; **Cf:** *C. flavens*; **Cu:** *C. urucurana*; **Cc:** *C. cajucara*; **Cmc:** *C. micradenus*; **Cmy:** *C. myricaefolius*.




Figure 4
Main compounds in analyzed *Croton* species

From polar fraction of *C. polycarpus* leaves three flavonols derivatives were isolated, known as quercetin and their glycosides quercitrin (quercetin-3-O- α -L-rhamnoside) and rutin (quercetin-3-rutinoside). Quercetin and rutin flavonoids present an EC_{50} of 47 μ g/mL and 64 μ g/mL respectively, against *C. quinquefasciatus* mosquitoes larvae, having a promissory larvicidal activity; nevertheless, the quercitrin doesn't give notable bioactivity in this specific LAB, although its antioxidant activity is widely known (Nuñez-Arevalo, 2005). Moreover, these compounds have showed the highest antioxidant activity, (Zhang *et al.*, 2014) and are useful ingredients in nutraceuticals and this preliminary study of *C. polycarpus* is a contribution to the knowledge of our native species. (Sintayehu *et al.*, 2012; Bose *et al.*, 2013).

CONCLUSIONS

Herein, are reported preliminary results about the presence of a group of sesquiterpenoids and a flavonol and their glycosides characterized by HRGC-MS and NMR from the outer bark and leaves of *C. polycarpus* tree. For compounds derived of

decaline nucleus whose structures depend on different substituent groups or double bond positions, like in δ -cadinene and cadina-1,4-diene, could be seen the functional diversity and stereochemical arrangements joint with the efficiency of the selected chromatographic systems to separate isomers and compounds with similar structures, that have potential as pedagogical strategy and other applications in formative programs. This work is part of a research project that studies the chemical composition and biological activity of native plants and other fractions are under current investigation.

Quercetin: MeOH d_4 . 1H -NMR (δ in ppm, J in Hz) 12.76 (1H, s, C5-OH), 6.24 (1H, d, J =2.0, C6-H), 6.46 (1H, d, J =2.0, C8-H), 7.51 (1H, d, J =2.1, C2'-H), 6.95 (1H, d, J =8.3, C5'-H), 7.39 (1H, dd J = 8.3, 2.0, C6'-H).

Quercitrin: Me $_2$ CO d_6 . 1H -NMR (δ in ppm, J in Hz) 12.64 (1H, s, C5-OH), 6.26 (1H, d, J =2.0, C6-H), 6.47 (1H, d, J =2.0, C8-H), 7.51 (1H, d, J =2.3, C2'-H), 6.95 (1H, d, J =8.2, C5'-H), 7.37 (1H, dd, - J =8.2, 2.3, C6'-H), 5.50 (1H, s broad, C1''-H), 4.23 (1H, d,

$J=1.6$, C2''-H), 3.75 (1H, dd, $J=9.2, 3.2$, C3''-H), 3.35 (1H, t, $J=9.2$, C4''-H), 3.42 (1H, dd, $J=9.4$, C5''-H), 0.91 (3H, d, $J=6.4$, C6''-CH₃); ¹³C -RMN δ in ppm) 158.9 (C-2), 136.8 (C-3), 180.3 (C-4), 165.9 (C-5), 95.5 (C-6), 166.0 (C-7), 100.5 (C-8), 159.3 (C-9), 106.7 (C-10), 123.5 (C-1'), 117.0 (C-2'), 146.8 (C-3'), 150.0 (C-4'), 117.1 (C-5'), 123.8 (C-6'), 103.7 (C-1''), 73.0 (C-2''), 72.4 (C-3''), 72.3 (C-4''), 70.8 (C-5''), 18.8 (C-6'').

Rutin: MeOHd₄. ¹H-NMR (δ in ppm, J in Hz) 12.7 (1H, d, $J=2.33$, C5-H), 6.27 (1H, d, $J=2.0$, C6-H), 6.49 (1H, d, $J=2.0$, C8-H), 7.55 (1H, d, $J=2.0$, C2'-H), 7.00 (1H, d, $J=8.0$, C5'-H), 7.46 (1H, dd, $J=8.0, 2.0$, C6'-H), 5.21 (1H, d, $J=8.2$, C1''-H), 4.91 (1H, d, $J=2.0$, C1'''-H), 1.25 (1H, d, $J=6.0$, C6'''-CH₃).

REFERENCES

- Aboagye F, Sam G, Massiot G, Lavaud C. 2000. Julocrotine, a glutaramide alkaloid from *Croton membranaceus*. **Fitoterapia** 71: 461 - 462.
- Aparecida M, Pinto A, Brabo S. 2002. Terpenoids from *Croton cajucara*. **Phytochemistry** 43: 823 - 828.
- Babili F, Fabre N, Moulis C, Bon M, Respaud M, Fourasté I. 1998. Three furanoditerpenes from the bark of *Croton campestris*. **Phytochemistry** 48: 165 - 169.
- Babili F, Fabre N, Moulis C, Fouraste I. 2006. Molluscicidal activity against *Bulinus truncatus* of *Croton campestris*. **Fitoterapia** 77: 384 - 387.
- Barbosa P, Abreu A, Batista E. 2007. Glutaramide alkaloids and terpenoids from *Croton pullei* var. *glabrior* Lanj. **Biochem Syst Ecol** 35: 887 - 890.
- Blanco O, Caicedo G. 2010. **Ingeniería conceptual de una planta piloto, para la extracción de aceites esenciales de especies nativas, en grupo de investigación de la Universidad Nacional, sede Bogotá** (Tesis de pregrado). Fundación Universidad de América, Bogotá, Colombia.
- Bose S, Maji S, Chakraborty P. 2013. Quercitrin from *Ixora coccinea* Leaves and its Anti-oxidant Activity. **J Pharmaceut Sci Technol** 2: 72 - 74.
- Bracher F, Randau K, Lerche H. 2008. Crototropone, a new tropone derivative from *Croton zehtneri*. **Fitoterapia** 79: 236 - 237.
- Campeol E, Flamini G, Chericoni S, Catalano S. 2001. Volatile Compounds from Three Cultivars of *Olea europaea* from Italy. **J Agric Food Chem** 49: 5409 - 5411.
- Campos A, Albuquerque F, Rao V, Maciel M, Pinto A. 2002. Investigations on the antinociceptive activity of crude extracts from *Croton cajucara* leaves in mice. **Fitoterapia** 73: 116 - 120.
- Carvalho T, Rabello de Moraes A, Buchele C. 2008. Antiplatelet activity of *Croton celtidifolius*. **Braz J Pharmaceut Sci** 44: 127 - 132.
- Cavalli J, Tomi F, Bernardini A, Casanova J. 2003. Composition and chemical variability of the bark oil of *Cedrelopsis grevei* H. Baillon from Madagascar. **Flav Frag J** 18: 532 - 538.
- Chavez P, Jolad S, Hoffmann J, Cole J. 1982. Four new 12-deoxyphorbol diesters from *Croton californicus*. **J Nat Prod** 45: 745 - 748.
- Choi H. 2005. Characteristic Odor Components of Kumquat (*Fortunella japonica* Swingle) Peel Oil. **J Agric Food Chem** 53: 1642 - 1647.
- Coussio J, Desmarchelier C, Witting F, Cicca G. 1997. Effects of Sangre de Drago from *Croton lechleri* Muell.-Arg. on the production of active oxygen radicals. **J Ethnopharmacol** 58: 103 - 108.
- DalBó S, Jürgensen S, Horst H. 2006. Analysis of the antinociceptive effect of the protoanthocyanidin-rich fraction obtained from *Croton celtidifolius* barks. **Pharmacol Biochem Behav** 85: 317 - 323.
- Elías V, Simoneit B, Cardoso J. 1997. Analysis of Volatile sesquiterpenoids in environmental and geological samples. **High Resolution Chromatography** 20: 305 - 309.
- Fathiazad F, Delazar A, Amiri R, Sarker S. 2006. Extraction of Flavonoids and Quantification of Rutin from waste Tobacco Leaves. **Iran J Pharmaceut Res** 3: 222 - 227.
- Giang P, Lee J, Son P, Lee J, Otsuka H. 2004. Flavonoid glucosides from the leaves of *Croton tonkinensis* GAGNEP. **Chem Pharmaceut Bull** 52: 879 - 882.
- Hamm S, Bleton J, Tchaplal A. 2004. Headspace solid phase microextraction for screening for the presence of resins in Egyptian archaeological samples. **J Separation Sci** 27: 235 - 243.

- Hamm S, Bleton J, Connan J, Tchaplal A. 2005. A chemical investigation by headspace SPME and GC-MS of volatile and semi-volatile terpenes in various olibanum samples. **Phytochemistry** 66: 1499 - 1514.
- Högnadóttir A, Rouseff, R. 2003. Identification of aroma active compounds in orange essence oil using gas chromatography-olfactometry and gas chromatography-mass spectrometry. **J Chromatograph A** 998: 201 - 211.
- Jaramillo B, Olivero J, Muñoz K. 2007. Composición química volátil y toxicidad aguda (CL₅₀) frente a *Artemia salina* del aceite esencial de *Croton malambo* colectado en la costa norte colombiana. **Scientia et Technica** 33: 299 - 302.
- Jaramillo B, Duarte E, Muñoz K, Stashenko E. 2010. Composición química volátil del aceite esencial de *Croton malambo* H. Karst. colombiano y determinación de su actividad antioxidante. **Rev Cub Plant Med** 15: 133 - 142.
- Jaramillo-Colorado B, Duarte-Restrepo E, Jaimes L. 2016. Bioactividad del aceite esencial de *Croton trinitatis* Millsp Colombiano. **Bol Latinoam Caribe Plant Med Aromat** 15: 249 - 257.
- Kafuku G, Lam M, Kansedo J, Lee K., Mbarawa M. 2010. *Croton megalocarpus* oil: A feasible nonedible oil source for biodiesel. **Bioresource Technol** 101: 7000 - 7004.
- Karioti A, Skaltsa H, Demetzos C, Perdetzoglou D. 2003. Effect of nitrogen concentration of the nutrient solution on the volatile constituents of leaves of *Salvia fruticosa* Mill. in solution culture. **J Agric Food Chem** 51: 6505 - 6508.
- Krebs H, Ramiarantsoa H. 1996. Clerodane diterpenes and other constituents of *Croton hovarum*. **Phytochemistry** 41: 561 - 563.
- Lopes D, Bizzo H, Sobrinho A, Pereira M. 2000. Linalool-rich essential oil from leaves of *Croton cajucara* Benth. **J Essent Oil Res** 12: 705 - 708.
- Lopes e Lopes M, Saffi J, Echeverrigaray S, Pegas J, Salvador M. 2004. Mutagenic and antioxidant activities of *Croton lechleri* sap in biological systems. **J Ethnopharmacol** 95: 437 - 445.
- Martinsen A, Baccelli C, Navarro I, Abad A, Quetin-Leclercq J, Morel N. 2010. Vascular activity of a natural diterpene isolated from *Croton zambesicus* and of a structurally similar synthetic trachylobane. **Vasc Pharmacol** 52: 63 - 69.
- McChesney J, Clark A, Silveira E. 1991. Antimicrobial diterpenes of *Croton sonderianus*, 1. Hardwickic and 3,4-secotrachylobanoic acids. **J Nat Prod** 54: 1625 - 1633.
- McLafferty F, Turecek F. 1993. **Interpretation of mass spectra**. University Science Books. Fourth Edition. Sausalito, California, USA.
- McLaughlin J, Rogers LL, Anderson J. 1998. The use of biological assays to evaluate botanicals. **Drug Informat J** 32: 513 - 524.
- Mendoza-Meza D, Benavides-Henriquez H, Taborda-Martínez M. 2014. Actividad acaricida del aceite esencial de la corteza de *Croton malambo* H. Karst, metil-eugenol y metil-isoeugenol contra *Dermatophagoides farinae* Hughes, 1961. **Bol Latinoam Caribe Plant Med Aromat** 13: 537 - 544.
- Muñoz-Acevedo A, Puerto C, Rodríguez J, Aristizabal-Cordoba S, Kouznetsov V. 2014. Estudio químico-biológico de los aceites esenciales de *Croton malambo* H. Karst y su componente mayoritario, metileugenol. **Bol Latinoam Caribe Plant Med Aromat** 13: 336 - 343.
- Murillo JA. 1999. Composición y distribución del género *Croton* (Euphorbiaceae) en Colombia con cuatro especies nuevas. **Caldasia** 21: 141 - 166.
- Murillo J. 2004. Las Euphorbiaceae de Colombia. **Biota Colombiana** 5: 183 - 200.
- Nardi G, Felippi R, Dalbó S, Siqueira-Junior J. 2003. Antiinflammatory and antioxidant effects of *Croton celtidifolius* bark. **Phytomedicine** 10: 176 - 184.
- Ngadjui B, Folefoc G. 1999. Crotonadiol, a labdane diterpenoid from the stem bark of *Croton zambesicus*. **Phytochemistry** 51: 171 - 174.
- Núñez-Arévalo L. 2005 **Contribución al estudio químico y de bioactividad de las hojas senescentes de *Croton funckianus* [Euphorbiaceae]** (Tesis de grado). Universidad Nacional de Colombia, Bogotá, Colombia.
- Núñez-Arévalo L, Moreno-Murillo B, Quijano-Celis C, Pino J. 2010. Composition of the essential oil from leaves of *Croton bogotanus* Cuatrec grown in Colombia. **J Essent Oil Res** 22: 486 - 487.

- Palmeira Jr. S, Conserva L, Silveira E. 2005. Two clerodanes diterpenes and flavonoids from *Croton brasiliensis*. **J Braz Chem Soc** 16: 1420 - 1424.
- Payo A, Sandoval D, Vélez H, Oquendo M. 2001. Alcaloide en la especie cubana *Croton micradenus* Urb. **Rev Cub Farm** 35: 61 - 65.
- Perazzo F, Carvalho J, Rodrigues M, Kadja E, Morais L, Aparecida M, Maciel M. 2007. Comparative anti-inflammatory and antinociceptive effects of terpenoids and an aqueous extract obtained from *Croton cajucara* Benth. **Rev Bras Farmacogn** 17: 521 - 528.
- Pino J, Marbot R, Payo A, Herrera P, Marti P. 2005. Chemical composition of the leaf oil of *Croton micradenus* Urb. from Cuba. **J Essent Oil Bear Plant** 8: 1 - 5.
- Pino J, Marbot R, Payo A, Herrera P, Martí P. 2006. Volatile leaf oils from Cuban Euphorbiaceae: *Croton rosmarinoides* Millsp., *Croton litoralis* Urb., *Croton spiralis* Muell. Arg. and *Croton myricifolius* Griseb. **J Essent Oil Res** 18: 256 - 260.
- Puebla P, Correa S, Guerrero M, Carron R, San Feliciano A. 2005. New cysclerodane diterpenoids from *Croton schiedeana*s. **Chem Pharmaceut Bull** 53: 328 - 329.
- Ramírez G. 2003. Sangre de Drago (*Croton lechleri* Muell. Arg). **Natura Medicatrix** 21: 213 - 217.
- Rodrigues F, Costa J, Coutinho H. 2009. Synergy effects of the antibiotics gentamicin and the essential oil of *Croton zehntneri*. **Phytomedicine** 16: 1052 - 1055.
- Roengsumran S, Achayindee S, Petsom A. 1998. Two new cembranoids from *Croton oblongifolius*. **J Nat Prod** 61: 652 - 654.
- Shivashankar S, Roy T, Krishna P. 2012. Headspace Solid Phase Micro Extraction and GC/MS Analysis of the Volatile Components in Seed and Cake of *Azadirachta indica* A. juss. **Chem Bull Politech Univ Timisoara** 57: 1 - 6.
- Simionatto E, Bonani V, Farias Morel A, Ré Poppi N, Raposo Júnior J, Stuker C, Peruzzo G, Peres M, Hess S. 2007. Chemical composition and evaluation of antibacterial and antioxidant activities of the essential oil of *Croton urucurana* Baillon (Euphorbiaceae) stem bark. **J Braz Chem Soc** 18: 879 - 885.
- Silva P, Durán N, Hiruma-Lima C. 2003. Comparison of the gastro protective effect of a diterpene lactone isolated from *Croton cajucara* with its synthetic derivatives. **J Ethnopharmacol** 87: 169 - 174.
- Sintayehu B, Asres K, Raghavendra Y. 2012. Radical scavenging activities of the leaf extracts and a flavonoid glycoside isolated from *Cineraria abyssinica* Sch. Bip. Exa. Rich. **J Applied Pharmaceut Sci** 2: 44 - 49.
- Suárez A, Blanco Z, Compagnone R, Salazar-Bookaman M, Tillett M, Delle Monache F, Di Giulio C, Bruges G. 2003. Antinociceptive and anti-inflammatory effects of *Croton malambo* bark aqueous extract. **J Ethnopharmacol** 88: 11 - 14.
- Suárez A, Compagnone R, Salazar-Bookaman M, Zapata V, Alvarado C. 2006. Anti-inflammatory activity of *Croton cuneatus* aqueous extract. **J Ethnopharmacol** 105: 99 - 101.
- Sylvestre M, Pichette A, Longtin A, Nagaub F, Legault J. 2006. Essential oil analysis and anticancer activity of leaf essential oil of *Croton flavens* L. from Guadeloupe. **J Ethnopharmacol** 103: 99 - 102.
- Tansakul P, De-Eknamkul W. 1998. Geranylgeraniol 18-hydroxylase: The last enzyme on the plaunotol biosynthetic pathway in *Croton sublyratus*. **Phytochemistry** 47: 1241 - 1246.
- Tellez M, Estell R, Fredrickson E, Powell J, Wedge D, Schrader K, Kobaisky M. 2001. Extracts of *Flourensia cernua* (L): Volatile constituents and Antifungal, Antialgal, and Antitermite Bioactivities. **J Chem Ecol** 27: 2263 - 2273.
- Tello-Camacho E. 2005. **Constituyentes mayoritarios del extracto no polar de hojas de *Alchornea glandulosa* End. & Poepp [Euphorbiaceae]** (Tesis de pregrado). Universidad Nacional de Colombia, Bogotá, Colombia.
- Urrea-Bulla A, Suarez M, Moreno-Murillo B. 2004. Biological activity of phenolic compounds from *Alchornea glandulosa*. **Fitoterapia** 75: 392 - 394.
- Vardar-Ünlü G, Candan F, Sökmen A, Daferera D, Polissiou M, Sökmen M, Dömnez E, Tepe, B. 2003. Antimicrobial and antioxidant activity of the essential oil and methanol extracts of *Thymus pectinatus* Fisch. et Mey. Var.

- pectinatus* (Lamiaceae). **J Agric Food Chem** 51: 63 - 67.
- Vegas-Mendoza S. 2010 **Estudio químico de los compuestos bioactivos de hojas senescentes de *Croton funckianus* [Euphorbiaceae]** (Tesis de maestría). Universidad Nacional de Colombia, Bogotá, Colombia.
- Vegas S, Moreno-Murillo B, Quevedo R. 2011. Cassipourol: Un diterpenoide monocíclico con actividad larvica en *Croton funckianus*. **Bol Latinoam Caribe Plant Med Aromat** 10: 228 - 232.
- Wang Y, Zou Z. 2008. Sesquiterpenes from the stems of *Croton caudatus* Geisel. Var. *tormentosus* Hook. **Chin J Nat Med** 6: 339 - 341.
- Yang X, Chen W, Zhao J, Yang L, Zhang H, Li L. 2009. Entkaurane diterpenes and phenolic compounds from *Croton kongensis* (Euphorbiaceae). **Biochem Syst Ecol** 37: 237 - 240.
- Zhang Y, Wang D, Zhou D, Zhang J. 2014. Purification and characterization of flavonoids from the leaves of *Zanthoxylum bungeanum* and correlation between their structure and antioxidant activity. **Plos One** 9: 1 - 11.
- Zou G, Heng-Su Z, Zhang H. 2010. Flavonoids from the stems of *Croton caudatus* Geisel. var. *tormentosus* Hook. **Molecules** 15: 1097 - 1102.