



Revista Colombiana de Estadística

ISSN: 0120-1751

revcoles_fcbo@unal.edu.co

Universidad Nacional de Colombia

Colombia

Kumar, Sunil; Bhogal, Sandeep; Nataraja, N. S.; Viswanathaiah, M.
Estimation of Population Mean in the Presence of Non-Response and Measurement Error
Revista Colombiana de Estadística, vol. 38, núm. 1, 2015, pp. 145-161
Universidad Nacional de Colombia
Bogotá, Colombia

Available in: <http://www.redalyc.org/articulo.oa?id=89938627009>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Estimation of Population Mean in the Presence of Non-Response and Measurement Error

Estimacion de la media poblacional con presencia de errores de medición y no respuesta

SUNIL KUMAR^{1,a}, SANDEEP BHOUGAL^{2,b}, N. S. NATARAJA^{1,c},
M. VISWANATHAIAH^{1,d}

¹ALLIANCE UNIVERSITY, BANGALORE, INDIA

²SCHOOL OF MATHEMATICS, SHRI MATA VAISHNO DEVI UNIVERSITY, KAKRYAL, INDIA

Abstract

Under classical survey sampling theory the errors mainly studied in the estimation are sampling errors. However, often non-sampling errors are more influential to the properties of the estimator than sampling errors. This is recognized by practitioners, researchers and many great works of literature regarding non-sampling errors have been published during last two decades, especially regarding non-response error which is one of the cornerstones of the non-sampling errors. The literature handles one kind of non-sampling error at a time, although in real surveys more than one non-sampling error is usually present. In this paper, two kinds of non-sampling errors are considered at the estimation stage: non-response and measurement error. An exponential ratio type estimator has been developed to estimate the population mean of the response variable in the presence of non-response and measurement errors. Theoretically and empirically, it has been shown that the proposed estimator is more efficient than usual unbiased estimator and other existing estimators.

Key words: Estimation, Mean Squared Error, Measurement Error, Non-response, Ratio Estimator, Sampling Error.

Resumen

En la teoría de muestreo de la encuesta clásica los errores estudiados principalmente en la estimación son el muestreo errores. Sin embargo, a menudo los errores ajenos al muestreo son más influyentes que las propiedades del

^aAssistant Professor. E-mail: sunilbhougal06@gmail.com

^bAssistant Professor. E-mail: sandeep.bhougal@smvdu.ac.in

^cAssistant Professor. E-mail: nataraja.ns@alliance.edu.in

^dAssistant Professor. E-mail: vmatam@gmail.com

estimador de errores de muestreo. Esto es reconocido por los profesionales, los investigadores y muchos grandes obras de la literatura en relación con los errores ajenos al muestreo se ha publicado en los últimos dos decenios, especialmente en relación con el error de falta de respuesta, que es una de las piedras angulares de los errores ajenos al muestreo. La literatura se ocupa de un tipo de error no muestral a la vez, aunque en las encuestas reales más de un error no muestral suele estar presente. En este trabajo, dos tipos de errores ajenos al muestreo son considerados en la etapa de la estimación: la falta de respuesta y el error de medición. Un tipo exponencial estimador de razón ha sido desarrollado para estimar la media poblacional de la variable de respuesta en presencia de errores de falta de respuesta y de medición. Teóricamente y empíricamente, se ha mostrado que el estimador propuesto es más eficiente que estimador insesgado habitual y otros estimadores existentes.

Palabras clave: error cuadrático medio, error de medición, error de muestreo, estimación, estimador de razón.

1. Introduction

Design based estimation methods use the sampling distribution that results when the values for the finite population units are considered to be fixed, and the variation of the estimates arises from the fact that statistics are based on a random sample drawn from the population rather than a census of the entire population, (see, Kish 1954, Sarndal, Swensson & Wretman 1992, Kish 1994, Gregoire 1998, Koch & Gillings 2006, Binder 2008, Dorazio 1999, Shabbir, Haq & Gupta 2014).

The results of a survey are used to make quantitative statements about the population studied i.e. descriptive statements about the aggregate population analytic statements about the relationship among subgroups of the population, or interpretive statements about the nature of social or economic processes. A survey error occurs when there is discrepancy between the statements and reality. These errors are of two types sampling and non-sampling errors. Sampling errors comprise the differences between the sample and the population due solely to the particular units that happen to have been selected. Non sampling errors encompass all other things that contribute to survey error. Non sampling errors are said to arise from wrongly conceived definitions, imperfections in the tabulation plans, failure to obtain response from all sample members, and so on. (see, Ilves 2011, Groves 1989).

In practice the researcher faces the problem of measurement error while collecting information from individuals. Measurement error is the difference between the value that is recorded and the true value of a variable in the study. For example, in surveys regarding household consumption/expenditure where the respondents are asked to report their consumption/expenditure catalog, there is a great likelihood that the respondents may fail to recall precisely how much they spent on various items over the interval. Many researchers have studied measurement errors like Cochran (1963), Cochran (1977), Fuller (1995), Shalabh (1997), Manisha & Singh (2001), Manisha & Singh (2002), Wang (2002), Allen, Singh & Smarandache

(2003), Singh & Karpe (2007), Singh & Karpe (2008), Singh & Karpe (2009), Singh & Karpe (2010), (Gregoire & Salas 2009, Salas & Gregoire 2010), Shukla, Pathak & Thakur (2012), and Sharma & Singh (2013), etc.

Another problem the researcher faces is due to non-response which refers to the failure to collect information from one or more respondents on one or more variables. The reasons non-response occurs include non-availability of the respondents at home, refusal to answer the questionnaire, lack of information, etc. Hansen & Hurwitz (1946) considered the problem of non-response while estimating the population mean by taking a sub sample from the non-respondent group with the help of extra efforts and an estimator was proposed by combining the information available from the response and non-response groups. In estimating population parameters like the mean, total or ratio, sample survey experts sometimes use auxiliary information to estimate improve precision. When the population mean of the auxiliary variable X is known and in presence of non-response, the problem of estimating the population mean of the study variable Y has been discussed by Cochran (1977), Rao (1986), Khare & Srivastava (1997), Kumar, Singh, Bhoulgal & Gupta (2011), Singh & Kumar (2008). In Hansen & Hurwitz (1946) method, questionnaires are mailed to all the respondents included in a sample and a list of non-respondents is prepared after the deadline is over. Then a sub sample is drawn from the set of non-respondents and a direct interview is conducted with the selected respondents and the necessary information is collected.

Researchers who have studied non-response have ignored the presence of possible measurement errors and researchers who have studied measurement errors have neglected non-response. In practice, it is possible for a researcher to face the problem of measurement error and non-response at the same time. Jackman (1999) dealt with both non-response and measurement error simultaneously, in the case of voter turnout, where a reasonably large body of vote validation studies supply auxiliary information, allowing the components of bias in survey estimates of turnout rates to be isolated. Averaging over the auxiliary information provides bounds on the quantity of interest, yielding an estimate corrected for both non-response and measurement error. Further, Dixon (2010) studied the estimation of non-response bias and measurement error on the data from Consumer Expenditure Quarterly Interview Survey (CEQ), Current Population Survey (CPS) and National Health Interview Survey (NHIS), an attempt to measure the differences in employment status of Washington.

In this paper, we have developed new estimators for estimating the population mean of the variable of interest when there is measurement error and non-response error in the study as well as in the auxiliary variable. An empirical study is carried out to show the efficiency of our suggested estimators over some available estimators.

2. Sampling Procedure and Some Well Defined Estimators

A simple random sample of size n is selected from the population of size N by a simple random sampling without replacement (SRSWOR) method. Let Y and X be the study variable and auxiliary variable, respectively. Let $\mu_Y = \frac{1}{N} \sum_{i=1}^N y_i$, $\mu_X = \frac{1}{N} \sum_{i=1}^N x_i$, $\sigma_y^2 = \frac{1}{N} \sum_{i=1}^N (y_i - \mu_Y)^2$ and $\sigma_x^2 = \frac{1}{N} \sum_{i=1}^N (x_i - \mu_X)^2$ denote the population mean and the population variance of the study variable y and auxiliary variables x .

Let (x_i, y_i) be the observed values and (X_i, Y_i) be the true values of two characteristics (x, y) respectively associated with the i^{th} ($i = 1, 2, \dots, n$) sample unit. Let the measurement errors be

$$U_i = y_i - Y_i,$$

and

$$V_i = x_i - X_i.$$

Measurement errors are assumed to be random in nature and they are uncorrelated with mean zero and variances σ_U^2 and σ_V^2 respectively. Let σ_X^2 and σ_Y^2 denote the variances of the auxiliary variable X and the variable of interest Y respectively for the population, C_y and C_x be the coefficient of variations of variable Y and X respectively for the population, and let ρ_{yx} be the coefficient of correlation between the variable Y and X for the population. We further assume that the measurement errors for variable Y and X are independent.

If there is some non-response, (Hansen & Hurwitz 1946) proposed a double sampling scheme for estimating population a mean, simple random sample of size n is selected and the questionnaire is mailed to the sampled units; the number of respondents in the sample is denoted by n_1 and the number of the non-respondents in the sample is denoted by n_2 and a sub-sample of size r ($r = n_2/k; k > 1$) is taken from non-respondents in the sample, where k is the inverse sampling ratio. Let (x_i^*, y_i^*) be the observed values and (X_i^*, Y_i^*) be the true values of two characteristics (x, y) respectively associated with the i^{th} ($i = 1, 2, \dots, n$) sample unit. Let the measurement error associated with the study variable be

$$u_i^* = y_i^* - Y_i^*.$$

If there is complete response on the auxiliary variable, let the measurement error associated with the auxiliary variable be

$$v_i = x_i - X_i.$$

If there is some non-response on the auxiliary variable, let the measurement error associated with the auxiliary variable be

$$v_i^* = x_i^* - X_i^*.$$

The measurement errors are random in nature having mean zero and variances σ_U^2 and σ_V^2 respectively for the responding units and $\sigma_{U(2)}^2$ and $\sigma_{V(2)}^2$ respectively for non-respondents of the population. Let $\sigma_{X(2)}^2$ and $\sigma_{Y(2)}^2$ be the variances for variables X and Y respectively for population non-respondents and $\rho_{yx(2)}$ be the coefficient of correlation between the variable X and Y for non-respondents of the population. Let $C_{x(2)}$ and $C_{y(2)}$ be the coefficient of variations for variables X and Y respectively for the non-respondents in the population. It is further assumed that the measurement errors for variables X and Y are independent.

The usual unbiased estimator for the population mean of the study variable in the presence of measurement error is given as

$$t_0 = \hat{\mu}_y = \frac{1}{n} \sum_{i=1}^n y_i$$

The variance in the presence of *measurement error* of the mean per unit estimator is given as

$$Var(t_0) = \lambda_2 (\sigma_y^2 + \sigma_U^2), \quad (1)$$

where $\lambda_2 = (\frac{1}{n} - \frac{1}{N})$.

Shalabh (1997) developed the following ratio-type estimator in the presence of *measurement error* $t_1 = \hat{\mu}_y \left(\frac{\mu_x}{\hat{\mu}_x} \right)$; where $\hat{\mu}_x = \frac{1}{n} \sum_{i=1}^n x_i$; $\mu_x = \frac{1}{N} \sum_{i=1}^N X_i$. The mean square error of t_1 (using finite population correction factor) is given as:

$$MSE(t_1) \cong \lambda_2 \{ \sigma_y^2 + R^2 \sigma_x^2 - 2R\rho_{yx}\sigma_y\sigma_x + \sigma_U^2 + R^2 \sigma_V^2 \}, \quad (2)$$

Shukla et al. (2012) developed the following estimator

$$t_2 = \alpha \hat{\mu}_y \left(\frac{\hat{\mu}_x^\delta}{\mu_x} \right) + (1 - \alpha) \hat{\mu}_y,$$

where $\hat{\mu}_x^\delta = \frac{N\mu_x - n\hat{\mu}_x}{N-n}$ and α is a suitable constant.

The mean square error of t_2 in the presence of *measurement error* is given as:

$$MSE(t_2) \cong \lambda_2 \{ \sigma_y^2 + \mu_1^2 R^2 \sigma_x^2 - 2\mu_1 R\rho_{yx}\sigma_y\sigma_x + \sigma_U^2 + \mu_1^2 R^2 \sigma_V^2 \}, \quad (3)$$

where $\mu_1 = \alpha \frac{n}{N-n}$.

$$MSE(t_2) = MSE_1(t_2) + MSE_2(t_2),$$

where

$$MSE_1(t_2) \cong \lambda_2 \{ \sigma_y^2 + \mu_1^2 R^2 \sigma_x^2 - 2\mu_1 R\rho_{yx}\sigma_y\sigma_x \}, \quad (4)$$

is the mean square error of t_2 without *measurement error*, and

$$MSE_2(t_2) \cong \lambda_2 \{ \sigma_U^2 + \mu_1^2 R^2 \sigma_V^2 \}, \quad (5)$$

is the contribution of *measurement error* to the mean square error of t_2 .

The optimum value of μ_1 is given as

$$\mu_{1(opt)} = \frac{\sigma_x^2}{\sigma_x^2 + \sigma_y^2} \rho_{yx} \frac{C_y}{C_x} = \mu_0, \quad (\text{say}).$$

The optimum mean square error of t_2 is given as

$$\min .MSE(t_2) = \lambda_2 \{ \sigma_y^2 + \mu_0^2 R^2 \sigma_x^2 - 2\mu_0 R \rho_{yx} \sigma_y \sigma_x + \sigma_U^2 + \mu_0^2 R^2 \sigma_V^2 \}. \quad (6)$$

When there is some non-response, it is assumed that the population of size N is composed of two mutually exclusive groups, the N_1 respondents and the N_2 non-respondents, though their sizes are unknown. Let $\mu_{Y_1} = \frac{1}{N_1} \sum_{i=1}^{N_1} y_i$ and $\sigma_{y(1)}^2 = \frac{1}{N_1-1} \sum_{i=1}^{N_1} (y_i - \mu_{Y_1})^2$ denote the mean and variance of the response group. Similarly, let $\mu_{Y_2} = \frac{1}{N_2} \sum_{i=1}^{N_2} y_i$ and $\sigma_{y(2)}^2 = \frac{1}{N_2-1} \sum_{i=1}^{N_2} (y_i - \mu_{Y_2})^2$ denote the mean and variance of the non-response group. The population mean can be written as $\mu_Y = W_1 \mu_{Y_1} + W_2 \mu_{Y_2}$, where $W_1 = N_1/N$ and $W_2 = N_2/N$. Let $\hat{\mu}_{y_1} = \frac{1}{n_1} \sum_{i=1}^{n_1} y_i$ and $\hat{\mu}_{y_{2r}} = \frac{1}{r} \sum_{i=1}^r y_i$ denote the means of the n_1 responding units and the r sub sampled units. Thus an unbiased estimator of the population mean \bar{Y} due to Hansen and Hurwitz is given by

$$\hat{\mu}_y^* = w_1 \hat{\mu}_{y_1} + w_2 \hat{\mu}_{y_{2r}},$$

where $w_1 = n_1/n$ and $w_2 = n_2/n$ are responding and non-responding proportions in the sample. The variance of $\hat{\mu}_y^*$ to terms of order n^{-1} , is given by

$$Var(\hat{\mu}_y^*) = \lambda_2 \sigma_y^2 + \theta \sigma_{y(2)}^2, \quad (7)$$

where $\lambda_2 = \frac{1-f}{n}$, $\theta = \frac{W_2(k-1)}{n}$, $C_y = \sigma_y/\mu_Y$ and $C_{y(2)} = \sigma_{y(2)}/\mu_Y$, (see, Cochran 1977, p. 371).

Let the information on auxiliary variable x be available and correlated with study variable y . In some situations, information on the auxiliary variable is not fully available i.e. non-response on auxiliary variable. One can define in similar manner to the above the auxiliary variable i.e. $\hat{\mu}_{x_1} = \frac{1}{n_1} \sum_{i=1}^{n_1} x_i$ and $\hat{\mu}_{x_{2r}} = \frac{1}{r} \sum_{i=1}^r x_i$ denotes the means of responding and r sub sampled units. Under such situation, an unbiased estimator for the population mean \bar{X} of the auxiliary variable as

$$\hat{\mu}_x^* = w_1 \hat{\mu}_{x_1} + w_2 \hat{\mu}_{x_{2r}}.$$

The variance of $\hat{\mu}_x^*$ is

$$Var(\hat{\mu}_x^*) = \lambda_2 \sigma_x^2 + \theta \sigma_{x(2)}^2. \quad (8)$$

Cochran (1977) proposed the following ratio-type estimator of population mean

$$t_3 = \hat{\mu}_y^* \left(\frac{\mu_X}{\hat{\mu}_x^*} \right)$$

The mean square error of t_3 is given as

$$MSE(t_3) \cong \lambda_2 (\sigma_y^2 + R^2 \sigma_x^2 - 2R \rho_{yx} \sigma_y \sigma_x) + \theta (\sigma_{y(2)}^2 + R^2 \sigma_{x(2)}^2 - 2R \rho_{y(2)x(2)} \sigma_{y(2)} \sigma_{x(2)}) \quad (9)$$

Singh & Karpe (2008) suggested the following generalized estimator of population mean

$$t_4 = \hat{\mu}_y^* \left(\frac{\mu_X}{\hat{\mu}_x^*} \right)^{\alpha_1} \left(\frac{\mu_X}{\hat{\mu}_x} \right)^{\alpha_2},$$

where α_1 and α_2 are suitably chosen constants.

The optimum mean square error of t_4 is given as

$$\min .MSE(t_4) \cong \lambda_2 \{ \sigma_y^2 (1 - \rho_{yx}) \} + \theta \left\{ \sigma_{y(2)}^2 (1 - \rho_{yx(2)}) \right\}, \quad (10)$$

In the situation where there is measurement and non-response both in the study and auxiliary variables, one can obtain the following estimator

$$t_5 = \hat{\mu}_y^* \frac{\hat{\mu}_x^\delta}{\mu_X} \exp \left(\frac{\hat{\mu}_x^\delta - \hat{\mu}_x^*}{\hat{\mu}_x^\delta + \hat{\mu}_x^*} \right),$$

where $\hat{\mu}_x^\delta = \frac{N\mu_X - n\hat{\mu}_x^*}{N-n}$.

The MSE of the estimator t_5 is given as

$$MSE(t_5) = MSE_1(t_5) + MSE_2(t_5), \quad (11)$$

where

$$\begin{aligned} MSE_1(t_5) \cong \lambda_2 \left\{ \sigma_y^2 + \frac{1}{4} \left(\frac{N+2n}{N-n} \right)^2 R^2 \sigma_x^2 - \left(\frac{N+2n}{N-n} \right) R \rho_{yx} \sigma_y \sigma_x \right\} \\ + \theta \left\{ \sigma_{y(2)}^2 + \frac{1}{4} \left(\frac{N+2n}{N-n} \right)^2 R^2 \sigma_{x(2)}^2 \right. \\ \left. - \left(\frac{N+2n}{N-n} \right) R \rho_{yx(2)} \sigma_{y(2)} \sigma_{x(2)} \right\}, \end{aligned} \quad (12)$$

is the mean square error of t_5 without *measurement error*, and

$$\begin{aligned} MSE_2(t_5) \cong \lambda_2 \left\{ \sigma_U^2 + \frac{1}{4} \left(\frac{N+2n}{N-n} \right)^2 R^2 \sigma_V^2 \right\} \\ + \theta \left\{ \sigma_{U(2)}^2 + \frac{1}{4} \left(\frac{N+2n}{N-n} \right)^2 R^2 \sigma_{V(2)}^2 \right\}, \end{aligned} \quad (13)$$

is the contribution of *measurement error* to the mean square error of t_5 .

In the present study, we have proposed exponential ratio type estimator in the situation where non-response and measurement errors are present in both study variable and auxiliary variable.

3. The Suggested Estimator

Non sampling errors are present in both sample surveys and censuses, and can occur at any stage of the survey process. There are many potential sources of

non-sampling error, for example, businesses not responding to a survey, processing errors, or respondents unintentionally reporting incorrect values. The greater the impact these sources of error, the greater the difference will be between our survey (or census) estimate and the true value. The following is the proposed estimator when there is non-response and measurement error in both study and auxiliary variable

$$t = \hat{\mu}_y^* \frac{\hat{\mu}_x^\delta}{\mu_X} \left\{ \alpha \exp \left(\frac{\hat{\mu}_x^* - \hat{\mu}_x^{\delta}}{\hat{\mu}_x^* + \hat{\mu}_x^{\delta}} \right) + (1 - \alpha) \exp \left(\frac{\hat{\mu}_x^{\delta} - \hat{\mu}_x^*}{\hat{\mu}_x^{\delta} + \hat{\mu}_x^*} \right) \right\}. \quad (14)$$

For $\alpha = 0$, the proposed estimator t becomes equal to estimator t_5 .

To obtain the expressions of mean squared error (MSE) of the proposed estimator, let us assume

$$\omega_y^* = \frac{1}{\sqrt{n}} \sum_{i=1}^n (y_i^* - \mu_Y), \quad \omega_U^* = \frac{1}{\sqrt{n}} \sum_{i=1}^n U_i^*, \quad \omega_x^* = \frac{1}{\sqrt{n}} \sum_{i=1}^n (x_i^* - \mu_X) \quad \text{and} \\ \omega_V^* = \frac{1}{\sqrt{n}} \sum_{i=1}^n V_i^*.$$

$$\text{Adding } \omega_y^* \text{ and } \omega_U^*, \text{ we have } \omega_y^* + \omega_U^* = \frac{1}{\sqrt{n}} \{ \sum_{i=1}^n (y_i^* - \mu_Y) + \sum_{i=1}^n U_i^* \}.$$

Multiplying both sides by $\frac{1}{\sqrt{n}}$, we have

$$\frac{1}{\sqrt{n}} (\omega_y^* + \omega_U^*) = \left\{ \frac{1}{n} \sum_{i=1}^n (y_i^* - \mu_Y) + \frac{1}{n} \sum_{i=1}^n (y_i^* - Y_i^*) \right\}$$

or

$$\frac{1}{\sqrt{n}} (\omega_y^* + \omega_U^*) = \hat{\mu}_y^* - \mu_Y,$$

or

$$\hat{\mu}_y^* = \mu_Y + \frac{1}{\sqrt{n}} (\omega_y^* + \omega_U^*).$$

Similarly, one can obtain

$$\hat{\mu}_x^* = \mu_X + \frac{1}{\sqrt{n}} (\omega_x^* + \omega_V^*).$$

Further

$$\left. \begin{aligned} E \left(\frac{\omega_y^* + \omega_U^*}{\sqrt{n}} \right)^2 &= \lambda_2 (\sigma_y^2 + \sigma_U^2) + \theta (\sigma_{y(2)}^2 + \sigma_{U(2)}^2) \\ E \left(\frac{\omega_x^* + \omega_V^*}{\sqrt{n}} \right)^2 &= \lambda_2 (\sigma_x^2 + \sigma_V^2) + \theta (\sigma_{x(2)}^2 + \sigma_{V(2)}^2) = B \text{ (say)} \\ E \left\{ \left(\frac{\omega_y^* + \omega_U^*}{\sqrt{n}} \right) \left(\frac{\omega_x^* + \omega_V^*}{\sqrt{n}} \right) \right\} &= \lambda_2 \rho_{yx} \sigma_y \sigma_x + \theta \rho_{yx(2)} \sigma_{y(2)} \sigma_{x(2)} = A \text{ (say)} \end{aligned} \right\},$$

Expressing t in terms of ω_i^* ; $i = x, y, U, V$; we have

$$t = (\mu_Y + W_y) \left(1 - \frac{nW_x}{(N-n)\mu_X} \right) \left[\alpha \exp \left\{ \frac{NW_x}{2(N-n)\mu_X} \left(1 + \frac{(N-2n)W_x}{2(N-n)\bar{X}} \right)^{-1} \right\} \right. \\ \left. + (1 - \alpha) \exp \left\{ \frac{-NW_x}{2(N-n)\mu_X} \left(1 + \frac{(N-2n)W_x}{2(N-n)\mu_X} \right)^{-1} \right\} \right],$$

where $W_y = \frac{1}{\sqrt{n}} (\omega_y^* + \omega_U^*)$ and $W_x = \frac{1}{\sqrt{n}} (\omega_x^* + \omega_V^*)$.

Simplifying and ignoring terms of order greater than two, one can obtain

$$\begin{aligned} t - \mu_Y &\cong W_y - \frac{1}{2\mu_X} \left(\frac{N+2n}{N-n} \right) W_x (\mu_Y + W_y) + \frac{3\mu_Y}{8\mu_X^2} \left(\frac{N}{N-n} \right)^2 W_x^2 \\ &\quad + \frac{\alpha}{\mu_X} \left(\frac{N}{N-n} \right) (\mu_Y + W_y) W_x. \end{aligned} \quad (15)$$

Taking expectation on both sides of (15), we get the bias of t as

$$\begin{aligned} B(t) &\cong \frac{1}{\mu_X (N-n)} \left\{ \alpha N - \frac{N+2n}{2} \right\} (\lambda_2 \rho_{yx} \sigma_y \sigma_x + \theta \rho_{yx(2)} \sigma_{y(2)} \sigma_{x(2)}) \\ &\quad + \frac{3}{8} \frac{\mu_Y^2}{\mu_X^2} \left(\frac{N}{N-n} \right)^2 \left\{ \lambda_2 (\sigma_X^2 + \sigma_V^2) + \theta (\sigma_{X(2)}^2 + \sigma_{V(2)}^2) \right\} \end{aligned}$$

Squaring both sides of equation (15) ignoring terms of order greater than two and taking expectations, the mean squared error (MSE) of t is

$$\begin{aligned} MSE(t) &= E(t - \mu_Y)^2 \cong E \left[W_y - \frac{\mu_Y}{2\mu_X} \left(\frac{N+2n}{N-n} \right) W_x + \alpha \frac{\mu_Y}{\mu_X} \left(\frac{N}{N-n} \right) W_x^2 \right]^2 \\ &= E \left[W_y^2 + \left(\frac{N+2n}{N-n} \right)^2 \frac{\mu_Y^2 W_X^2}{4\mu_X^2} - \frac{\mu_Y}{\mu_X} \left(\frac{N+2n}{N-n} \right) W_x W_y \right] \\ MSE(t) &= MSE_1(t) + MSE_2(t), \end{aligned} \quad (16)$$

where

$$\begin{aligned} MSE_1(t) &= \left\{ \lambda_2 \sigma_y^2 + \theta \sigma_{y(2)}^2 \right\} + \frac{\mu_Y^2}{\mu_X^2} \left\{ \frac{1}{4} \left(\frac{N+2n}{N-n} \right)^2 + \alpha^2 \left(\frac{N}{N-n} \right)^2 \right. \\ &\quad \left. - \alpha \left(\frac{N}{N-n} \right) \left(\frac{N+2n}{N-n} \right) \right\} \left\{ \lambda_2 \sigma_x^2 + \theta \sigma_{x(2)}^2 \right\} \\ &\quad - \frac{\mu_Y}{\mu_X} \left(\frac{N+2n-2\alpha N}{N-n} \right) \left\{ \lambda_2 \rho_{yx} \sigma_x \sigma_y + \theta \rho_{yx(2)} \sigma_{x(2)} \sigma_{y(2)} \right\} \end{aligned}$$

is the MSE of t without measurement error, and

$$\begin{aligned} MSE_2(t) &= \left\{ \lambda_2 \sigma_U^2 + \theta \sigma_{U(2)}^2 \right\} + \frac{\mu_Y^2}{\mu_X^2} \left\{ \frac{1}{4} \left(\frac{N+2n}{N-n} \right)^2 \right. \\ &\quad \left. + \alpha^2 \left(\frac{N}{N-n} \right)^2 - \alpha \left(\frac{N}{N-n} \right) \left(\frac{N+2n}{N-n} \right) \right\} \left\{ \lambda_2 \sigma_V^2 + \theta \sigma_{V(2)}^2 \right\} \end{aligned}$$

is the contribution of measurement error to the MSE of t .

Minimize equation (16) with respect to α yields its optimum value as

$$\alpha = \left(\frac{N+2n}{2N} \right) - \left(\frac{N-n}{N} \right) \left(\frac{\mu_X}{\mu_Y} \right) \left(\frac{A}{B} \right). \quad (17)$$

Substituting the value of α from equation (17) in equation (16), we get the optimum MSE of t as

$$\min .MSE(t) = \lambda_2 (\sigma_y^2 + \sigma_U^2) + \theta (\sigma_{y(2)}^2 + \sigma_{U(2)}^2) - \frac{A^2}{B}. \quad (18)$$

4. Efficiency Comparisons

To compare the efficiency of the proposed estimator t with the existing estimator $t_0, t_1, t_2, \hat{\mu}_y^*, t_3, t_4$ and t_5 , we have

$$Var(t_0) - \min .MSE(t) \geq 0; \text{ if } \frac{A^2}{B} \geq \theta (\sigma_{y(2)}^2 + \sigma_{U(2)}^2). \quad (19)$$

$$MSE(t_1) - \min .MSE(t) \geq 0;$$

$$\text{if } \frac{A^2}{B} + \lambda_2 \{R^2 (\sigma_x^2 + \sigma_V^2) - 2R\rho_{yx}\sigma_y\sigma_x\} \geq \theta (\sigma_{y(2)}^2 + \sigma_{U(2)}^2). \quad (20)$$

$$\min .MSE(t_2) - \min .MSE(t) \geq 0$$

$$\text{if } \frac{A^2}{B} + \lambda_2 \{\mu_0^2 R^2 (\sigma_x^2 + \sigma_V^2) - 2\mu_0 R\rho_{yx}\sigma_y\sigma_x\} \geq \theta (\sigma_{y(2)}^2 + \sigma_{U(2)}^2). \quad (21)$$

$$Var(\hat{\mu}_y^*) - \min .MSE(t) \geq 0; \text{ if } \frac{A^2}{B} \geq (\lambda_2 \sigma_U^2 + \theta \sigma_{U(2)}^2). \quad (22)$$

$$MSE(t_3) - \min .MSE(t) \geq 0$$

$$\begin{aligned} \text{if } \frac{A^2}{B} \geq \lambda_2 (\sigma_U^2 - R^2 \sigma_x^2 + 2R\rho_{yx}\sigma_y\sigma_x) \\ + \theta (\sigma_{U(2)}^2 - R^2 \sigma_{x(2)}^2 + 2R\rho_{yx(2)}\sigma_{y(2)}\sigma_{x(2)}). \end{aligned} \quad (23)$$

$$MSE(t_5) - \min .MSE(t) \geq 0;$$

$$\text{if } \frac{A^2}{B} \geq \lambda_2 (\sigma_U^2 + \rho_{yx}^2 \sigma_y^2) + \theta (\sigma_{U(2)}^2 + \rho_{yx(2)}^2 \sigma_{y(2)}^2). \quad (24)$$

From (19)-(24), we envisage that the proposed estimator t outperforms existing estimators viz. $t_0, t_1, t_2, \hat{\mu}_y^*, t_3, t_4$ and t_5 , respectively.

5. Empirical Study

In this section, we demonstrate the performance of different estimators over the usual unbiased estimators, generating four populations from normal distribution with different choices of parameters by using R language program. The auxiliary information on variable X has been generated from $N(5, 10)$ population. This type of population is very relevant in most socio-economic situations with one interest and one auxiliary variable.

Population I: $X = N(5, 10)$; $4Y = X + N(0, 1)$; $y = Y + N(1, 3)$; $x = X + N(1, 3)$; $N = 5000$; $\mu_Y = 4.927167$; $\mu_X = 4.924306$; $\sigma_y^2 = 102.0075$; $\sigma_x^2 = 101.4117$; $\sigma_U^2 = 8.862114$; $\sigma_V^2 = 9.001304$; $\rho_{yx} = 0.995059$

N_1	N_2	$\sigma_{y(2)}^2$	$\sigma_{x(2)}^2$	$\sigma_{U(2)}^2$	$\sigma_{V(2)}^2$	$\rho_{yx(2)}$
4500	500	99.99174	99.87471	9.150544	8.756592	0.994916
4250	750	100.9428	100.8224	9.053862	8.766538	0.995535
4000	1000	104.2711	103.2349	8.821278	8.339179	0.995472

Population II: $X = N(5, 10)$; $Y = X + N(0, 1)$; $y = Y + N(1, 5)$; $x = X + N(1, 5)$; $N = 5000$; $\mu_Y = 4.996681$; $\mu_X = 5.013507$; $\sigma_y^2 = 97.12064$; $\sigma_x^2 = 95.95803$; $\sigma_U^2 = 23.96055$; $\sigma_V^2 = 24.19283$; $\rho_{yx} = 0.994822$

N_1	N_2	$\sigma_{y(2)}^2$	$\sigma_{x(2)}^2$	$\sigma_{U(2)}^2$	$\sigma_{V(2)}^2$	$\rho_{yx(2)}$
4500	500	97.02783	94.54578	22.80557	25.43263	0.994546
4250	750	98.27616	97.42674	23.27837	24.13829	0.994992
4000	1000	96.09359	94.71923	24.42978	23.03076	0.99467

Population III: $X = N(5, 10)$; $Y = X + N(0, 1)$; $y = Y + N(2, 3)$; $x = X + N(2, 3)$; $N = 5000$; $\mu_Y = 4.730993$; $\mu_X = 4.741928$; $\sigma_y^2 = 101.2633$; $\sigma_x^2 = 100.2288$; $\sigma_U^2 = 9.1025$; $\sigma_V^2 = 9.052019$; $\rho_{yx} = 0.995187$

N_1	N_2	$\sigma_{y(2)}^2$	$\sigma_{x(2)}^2$	$\sigma_{U(2)}^2$	$\sigma_{V(2)}^2$	$\rho_{yx(2)}$
4500	500	102.7504	101.2097	9.095136	8.8123	0.995045
4250	750	99.55993	99.49764	9.233619	8.805872	0.995314
4000	1000	105.4334	103.8947	9.277715	9.072151	0.995105

Population IV: $X = N(5, 10)$; $Y = X + N(0, 1)$; $y = Y + N(2, 5)$; $x = X + N(2, 5)$; $N = 5000$; $\mu_Y = 4.961081$; $\mu_X = 4.96178$; $\sigma_y^2 = 102.2408$; $\sigma_x^2 = 100.868$; $\sigma_U^2 = 25.94111$; $\sigma_V^2 = 25.03951$; $\rho_{yx} = 0.394221$

N_1	N_2	$\sigma_{y(2)}^2$	$\sigma_{x(2)}^2$	$\sigma_{U(2)}^2$	$\sigma_{V(2)}^2$	$\rho_{yx(2)}$
4500	500	103.5361	102.1031	25.31099	22.84483	0.394622
4250	750	103.6790	102.7446	24.6859	26.12337	0.395036
4000	1000	100.1031	99.31665	25.80394	24.50468	0.394778

We have computed the percent relative efficiencies (PRE's) of the estimators. The results are shown in Tables (1-5).

TABLE 1: Percent relative efficiencies of the estimators with respect to the Hansen and Hurwitz (1946) estimator $\hat{\mu}_y^*$ for population I.

Percent relative efficiencies of the estimator with respect to Hansen and Hurwitz (1946) estimator.										
N_1	N_2	Esti.	PRE without measurement error				PRE with measurement error			
			(1/k)				(1/k)			
			(1/2)	(1/3)	(1/4)	(1/5)	(1/2)	(1/3)	(1/4)	(1/5)
4500	500	$\hat{\mu}_y^*$	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
		t_5	3589.91	3588.7	3587.87	3587.36	407.81	407.34	406.95	406.62
		$t_{(opt)}$	10115.23	10091.44	10071.7	10055.67	562.28	561.26	560.42	559.7
4250	750	$\hat{\mu}_y^*$	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
		t_5	3643.38	3683.63	3715.99	3742.73	408.25	408.16	408.08	408.02
		$t_{(opt)}$	10283.99	10391.55	10477.01	10546.55	563.33	563.18	563.07	562.98
4000	1000	$\hat{\mu}_y^*$	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
		t_5	3612.39	3627.65	3639.68	3649.71	410.12	411.32	412.19	412.87
		$t_{(opt)}$	10302.55	10414.41	10497.72	10562.18	569.27	573.31	576.29	578.59

TABLE 2: Percent relative efficiencies of the estimators with respect to the Hansen and Hurwitz (1946) estimator $\hat{\mu}_y^*$ for population II.

Percent relative efficiencies of the estimator with respect to (Hansen & Hurwitz 1946) estimator										
N_1	N_2	Esti.	PRE without measurement error				PRE with measurement error			
			(1/k)				(1/k)			
			(1/2)	(1/3)	(1/4)	(1/5)	(1/2)	(1/3)	(1/4)	(1/5)
4500	500	$\hat{\mu}_y^*$	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
		t_5	8810.15	8774.57	8746.13	8723.19	208.96	209.04	209.12	209.18
		$t_{(opt)}$	9626.52	9582.74	9546.95	9517.16	219.1813	219.21	219.23	219.25
4250	750	$\hat{\mu}_y^*$	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
		t_5	8865.94	8876.05	8886.02	8896.18	209.63	210.21	210.23	211.04
		$t_{(opt)}$	9727.01	9761.53	9788.52	9810.21	220.04	220.72	221.25	221.67
4000	1000	$\hat{\mu}_y^*$	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
		t_5	8822.61	8803.71	8793.64	8789.89	208.53	208.31	208.14	208.02
		$t_{(opt)}$	9630.34	9595.3	9569.72	9550.23	218.97	218.84	218.75	218.68

TABLE 3: Percent relative efficiencies of the estimators with respect to the Hansen and Hurwitz (1946) estimator $\hat{\mu}_y^*$ for population III.

Percent relative efficiencies of the estimator with respect to Hansen and Hurwitz (1946) estimator										
N_1	N_2	Esti	PRE without measurement error				PRE with Measurement Error			
			(1/k)				(1/k)			
			(1/2)	(1/3)	(1/4)	(1/5)	(1/2)	(1/3)	(1/4)	(1/5)
4500	500	$\hat{\mu}_y^*$	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
		t_5	3764.03156	3749.06	3736.85	3726.78	399.90	400.13	400.32	400.48
		$t_{(opt)}$	10382.13	10356.68	10335.67	10318.03	552.05	553.01	553.81	554.49
4250	750	$\hat{\mu}_y^*$	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
		t_5	3801.01	3815.27	3826.87	3836.65	399.59	399.56	399.54	399.53
		$t_{(opt)}$	10449.44	10477.23	10499.40	10517.49	550.39	550.04	549.79	549.54
4000	1000	$\hat{\mu}_y^*$	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
		t_5	3755.45	3737.65	3725.56	3717.28	400.31	400.78	401.12	401.38
		$t_{(opt)}$	10379.70	10356.86	10340.41	10328.01	553.30	554.95	556.19	557.12

From the above tables, the following points are noted:

1. The performance of the proposed estimator t at its optimum shows efficient PRE with respect to the Hansen and Hurwitz's estimator and t_5 estimator for all the four populations.

For population I, it is noted that

TABLE 4: Percent relative efficiencies of the estimators with respect to the Hansen and Hurwitz (1946) estimator $\hat{\mu}_y^*$ for population IV.

Percent relative efficiencies of the estimator with respect to Hansen and Hurwitz (1946) estimator										
N_1	N_2	Estimators	PRE without measurement error				PRE with measurement error			
			(1/k)				(1/k)			
			(1/2)	(1/3)	(1/4)	(1/5)	(1/2)	(1/3)	(1/4)	(1/5)
4500	500	$\hat{\mu}_y^*$	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
		t_5	418.06	303.07	276.20	252.057	287.68	231.78	199.54	178.57
		$t_{(opt)}$	788.35	460.99	347.07	289.31	321.79	242.38	202.99	179.52
4250	750	$\hat{\mu}_y^*$	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
		t_5	262.46	277.82	243.19	218.04	254.54	198.65	169.74	152.09
		$t_{(opt)}$	571.21	346.40	269.54	230.90	272.20	201.63	170.01	152.13
4000	1000	$\hat{\mu}_y^*$	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
		t_5	307.48	258.20	222.85	196.56	232.11	178.47	152.52	137.21
		$t_{(opt)}$	471.64	294.80	234.89	204.92	242.70	179.39	152.53	137.74

- When $N_1 = 4500$; $N_2 = 500$, with increase in the value of k , the PRE of the estimators t_5 and $t_{(opt)}$ with respected to $\hat{\mu}_y^*$ increases in both cases of with and without Measurement error, respectively.
- When $N_1 = 4250$; $N_2 = 750$, in the case with Measurement error, the PRE's decreases and in the case without Measurement error, the PRE's increases with the increase in the value of k .
- When $N_1 = 4000$; $N_2 = 1000$, with increase in the value of, k increases in both cases with and without Measurement error, respectively.

For population II, it is noted that

- When $N_1 = 4500$; $N_2 = 500$, with increase in the value of k , the PRE of the estimators t_5 and $t_{(opt)}$ with respected to $\hat{\mu}_y^*$ decreases in the case without Measurement error and increases in the case with Measurement error.
- When $N_1 = 4250$; $N_2 = 750$, in the case with Measurement error, the PRE's of the estimators increases in both cases with the increase in the value of k .
- When $N_1 = 4000$; $N_2 = 1000$, with the increase in the value of k decreases in both cases with and without Measurement error, respectively.

For population III, it is noted that

- When $N_1 = 4500$; $N_2 = 500$ and $N_1 = 4500$; $N_2 = 500$ in the case without Measurement error, the PRE's decreases and in the case with Measurement error, the PRE's increases with the increase in the value of k .
- When $N_1 = 4250$; $N_2 = 750$, in the case without Measurement error, the PRE's increases and in the case with Measurement error, the PRE's decreases with the increase in the value of k .
- For population IV, it is envisaged that with poor correlation between the study and the auxiliary variable, performance of the proposed estimator is better than the other estimators.

From Table 5; it is envisaged that the proposed estimator at its optimum performs more efficiently than the usual unbiased estimator, Shalabh's (t_1) and Shukla et al. (t_2) estimators for estimating the population mean in the presence of measurement error.

TABLE 5: Percent relative efficiencies of the estimators with respect to usual unbiased estimator in the presence of measurement error (no non-response).

Estimator	Population I	Population II	Population III	Population IV
$t_0 = \hat{\mu}_y$	100.00	100.00	100.00	100.00
t_1	587.229	247.12	578.26	587.22
t_2	612.476	273.193	600.33	612.47
$t_{(opt)}$	612.480	273.214	600.39	612.48

6. Conclusion

An important goal is understanding, managing, controlling, and reporting all known sources of error having impact on the quality of our statistics. In the present study, we have proposed an estimator for estimating the population mean of the study variable. The suggested estimator uses auxiliary information to improve efficiencies in situations where there are non-response and measurement errors concerning study variable and auxiliary variable. The relative performance of the proposed estimators is compared with conventional estimators. The proposed estimator performs better than the usual unbiased estimator t_0 , Shalabh's (1997) (t_1) and Shukla, Pathak & Thakue (2012) (t_2) estimators in the presence of measurement error; and its performance is better than the usual unbiased estimator $\hat{\mu}_y^*$ and estimator t_5 in the presence of measurement error and non-response. The study is supported by empirical study based on four populations.

We recommend our proposed estimator for future assess to study the characteristics of the variable in interest where measurement errors and non-response occur in the survey.

Acknowledgements

Authors wish to thank the Editor-in-Chief Professor Leonardo Trujillo and the learned referees for their critical and constructive comments regarding improvement of the paper.

[Received: January 2014 — Accepted: September 2014]

References

- Allen, J., Singh, H. & Smarandache, F. (2003), 'A family of estimators of population mean using multi-auxiliary information in presence of measurement errors', *International Journal of Social Economics* **30**, 837–849.
- Binder, D. A. (2008), Design-based estimation, in P. J. Lavrakas, ed., 'Encyclopedia of Survey Research Methods', SAGE Publications, Inc., pp. 192–194. doi: 10.4135/9781412963947.
- Cochran, W. (1963), *Sampling Techniques*, John Wiley, New York.
- Cochran, W. (1977), *Sampling Techniques*, 3 edn, John Wiley & Sons, Inc., New York.
- Dixon, J. (2010), Assessing nonresponse bias and measurement error using statistical matching, in R. Harter, ed., 'Section on Survey Research Methods - JSM', American Statistical Association, pp. 3388–3396.
*<http://www.bls.gov/osmr/pdf/st100190.pdf>
- Dorazio, R. M. (1999), 'Design based and model based inference in surveys of freshwater mollusks', *Journal of the North American Benthological Society* **18**, 118–131.
- Fuller, W. (1995), 'Estimation in the presence of measurement error', *International Statistical Review* **63**, 121–147.
- Gregoire, T. G. (1998), 'Design-based and model-based inference in survey sampling: Appreciating the difference', *Canadian Journal of Forestry Research* **28**, 1429–1447.
- Gregoire, T. G. & Salas, C. (2009), 'Ratio estimation with measurement error in the auxiliary variate', *Biometrics* **65**, 590–598.
- Groves, R. M. (1989), *Survey Errors and Survey Costs*, Wiley, New York.
- Hansen, M. & Hurwitz, W. (1946), 'The problem of non-response in sample surveys', *Journal of American Statistical Association* **41**, 517–529.
- Ilves, M. (2011), Estimation in the presence of non-response and measurement errors, Proceedings of the Third Baltic – Nortic Conference in Survey Statistics, Norrfällsviken, Sweden, pp. 13–17.
- Jackman, S. (1999), 'Correcting surveys for non-response and measurement error using auxiliary information', *Electoral Studies* **18**, 7–27.
- Khare, B. & Srivastava, S. (1997), 'Transformed ratio type estimators for the population mean in presence of non-response', *Communication in Statistics – Theory and Methods* **26**, 1779–1791.
- Kish, L. (1954), 'Differentiation in metropolitan areas', *American Sociological Review* **19**, 388–398.

- Kish, L. (1994), 'Classical and model based estimators for forest inventory', *Silva Fennica* **28**, 3–14.
- Koch, G. G. & Gillings, D. B. (2006), Inference, design based vs model based, in S. Kotz, C. B. Read, N. Balakrishnan & B. Vidakovic, eds, 'Encyclopedia of Statistical Sciences', John Wiley and Sons Inc., New York.
- Kumar, S., Singh, H., Bhogal, S. & Gupta, R. (2011), 'A class of ratio-cum-product type estimators under double sampling in the presence of non-response', *Hacetatepe Journal of Mathematics and Statistics* **40**, 589–599.
- Manisha & Singh, R. (2001), 'An estimation of population mean in the presence of measurement errors', *Journal of the Indian Society of Agricultural Statistics* **54**, 13–18.
- Manisha & Singh, R. (2002), 'Role of regression estimator involving measurement errors', *Brazilian Journal of Probability and Statistics* **16**, 39–46.
- Rao, P. (1986), 'Ratio estimation with sub sampling the non-respondents', *Survey Methodology* **12**, 217–230.
- Salas, C. & Gregoire, T. (2010), 'Statistical analysis of ratio estimators and their estimators of variances when the auxiliary variate is measured with error', *European Journal of Forest Research* **129**, 847–861.
- Sarndal, C. E., Swensson, B. & Wretman, J. (1992), *Model Assisted Survey Sampling*, Springer, New York.
- Shabbir, J., Haq, A. & Gupta, S. (2014), 'A new difference-cum-exponential type estimator of finite population mean in simple random sampling', *Revista Colombiana de Estadística* **37**, 199–211.
- Shalabh (1997), 'Ratio method of estimation in the presence of measurement errors', *Journal of the Indian Society of Agricultural Statistics* **50**, 150–155.
- Sharma, P. & Singh, R. (2013), 'A generalized class of estimators for finite population variance in presence of measurement errors', *Journal of Modern Applied Statistical Methods* **12**, 231–241.
- Shukla, D., Pathak, S. & Thakur, N. S. (2012), 'Class(es) of factor-type estimator(s) in presence of measurement error', *Journal of Modern Applied Statistical Methods* **11**, 336–347.
- Shukla, D., Pathak, S. & Thakur, N. (2012), 'An estimator for mean estimation in presence of measurement error', *Research and Reviews: A Journal of Statistics* **1**, 1–8.
- Singh, H. & Karpe, N. (2007), 'Effect of measurement errors on a class of estimators of population mean using auxiliary information in sample surveys', *Journal of Statistical Research of Iran* **4**, 175–189.

- Singh, H. & Karpe, N. (2008), 'Estimation of population variance using auxiliary information in the presence of measurement errors', *Statistics in Transition-New Series* **9**, 443–470.
- Singh, H. & Karpe, N. (2009), 'On the estimation of ratio and product of two population means using supplementary information in presence of measurement errors', *Statistica, Anno* **69**, 27–47.
- Singh, H. & Karpe, N. (2010), 'Estimation of mean, ratio and product using auxiliary information in the presence of measurement errors in sample surveys', *Journal of Statistical Theory and Practice* **4**, 111–136.
- Singh, H. & Kumar, S. (2008), 'A regression approach to the estimation of the finite population mean in the presence of non-response', *Australian and New Zealand Journal of Statistics* **50**, 395–408.
- Wang, L. (2002), 'A simple adjustment for measurement errors in some dependent variable models', *Statistics & Probability Letters* **58**, 427–433.