



Bragantia

ISSN: 0006-8705

editor@iac.sp.gov.br

Secretaria de Agricultura e  
Abastecimento do Estado de São Paulo  
Brasil

Jiang, Liangrong; Zhu, Zhenzhong; Huang, Rongyu; Huang, Yumin; Wang, Houcong;  
Zheng, Jingsheng; Fang, Xunjun

Mapping and allelic sequencing of a long sterile lemma trait in rice

Bragantia, vol. 76, núm. 2, abril-junio, 2017, pp. 229-237

Secretaria de Agricultura e Abastecimento do Estado de São Paulo  
Campinas, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=90850849007>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

# Mapping and allelic sequencing of a long sterile lemma trait in rice

Liangrong Jiang<sup>1</sup>, Zhenzhong Zhu<sup>1</sup>, Rongyu Huang<sup>1</sup>, Yumin Huang<sup>1</sup>, Houcong Wang<sup>1</sup>, Jingsheng Zheng<sup>1\*</sup>, Xunjun Fang<sup>2</sup>

1.Xiamen University - School of Life Sciences - Xiamen (Fujian), China.

2.Hainan Institute of Tropical Agricultural Resources - Sanya (Hainan), China.

**ABSTRACT:** Some outer floral organs are unique in gramineous plants, like the sterile lemma and rudimentary glume in rice. However, their development mechanisms are still poorly understood. In this study, we used 4 mutants with long sterile lemma (LSL), named JF11, JF12, JF13 and JNY-7, to be crossed with Aijiaonante (AJNT) and Nipponbare (NIP), respectively. Genetic analysis indicated that LSL trait exhibited recessive heredity and was controlled by a common allele named *sl-1(t)*. Using the method of bulk segregant analysis and linkage analysis between SSR markers and LSL trait based on  $F_2$  populations, the *sl-1(t)* gene was located at the interval between RM20903 and RM20948 on chromosome 7. The interval harbors a known *G1* gene, which regulates the sterile

lemma trait. The findings of allelic sequencing showed an 11-base deletion in gene *G1* happened in the mutants of JF11, JF12 and JF13, which led to a frame-shifting mutation, whereas the mutant of JNY-7 had a base substitution that caused the change of the amino acid residue. Eight substitutions in the ORF and 10 in the upstream region from -1 to -824 were found between *Indica* and *Japonica* rice by DNA sequence analysis, but those polymorphisms have no effect on the gene function. In conclusion, we fine mapped the LSL gene, *sl-1(t)*, and found 2 kinds of mutant alleles conferring the gene function and the DNA polymorphisms of *G1* between *Indica* and *Japonica* rice.

**Key words:** *Oryza sativa* L., gene mapping, ALOG domain.

\*Corresponding author: zheng21006@xmu.edu.cn

Received: Apr. 1, 2016 – Accepted: Aug. 17, 2016

## INTRODUCTION

A flower is the vital reproductive organ determining quality of fruits and seed yield in plants. Since the 1980s, the mechanism of floral organ development has become one of hot spots in developmental biology by means of the floral organ mutants of model plants. The ABC model of plant flower development was put forward to elaborate the molecular mechanism of floral organ identity in eudicots based on their previous findings (Carpenter and Coen 1990; Bowman et al. 1991; Coen and Meyerowitz 1991; Coen and Carpenter 1993). Then, *FBP7* and *FBP11* genes, which regulated the identity and development of an ovule in petunia, were cloned in 1995 (Angenent et al. 1995; Colombo et al. 1995) and designated to an additional class D MADS box gene. Furthermore, the *SEP1/2/3* genes were found as necessary for the development of petals, stamens, and carpels in *Arabidopsis* and were identified as a new class gene (E-class gene) of the floral quartet model (Pelaz et al. 2000; Honma and Goto 2001; Galimba et al. 2012). The findings of D-class and E-class enriched the ABC model and made the classical one extending to the ABCDE model.

Moreover, many researchers found that the genes among the Classes A, B, and C are, relatively, conserved between grasses and edicts based on comparison with the homology of the MADS-box genes and analysis of the transgenic plants (Schmidt et al. 1993; Chung et al. 1995; Kang et al. 1995; Mena et al. 1996; Ambrose et al. 2000). It is known that rice is a typical model plant of the grasses and the monocots in general and has 5 kinds of genes showing similar function like the ones in Classes A, B, C, D and E. So far, many genes related to floral organ development in rice were identified and cloned, with *OsMADS15*, *OsMADS14* and *OsMADS18* belonging to Class A (Kyoizuka et al. 2000; Lim et al. 2000; Fornara et al. 2004), *OsMADS2*, *OsMADS4*, *OsMADS16*, and *OsMADS32*, to Class B (Chung et al. 1995; Kang et al. 1998; Moon et al. 1999; Kyoizuka et al. 2000; Rensing et al. 2008; Wang et al. 2015), *OsMADS3*, *OsMADS58*, *RAG*, *DL*, *CPP1* and *DFO1*, to Class C (Coen and Carpenter 1993; Kang et al. 1995; Kang et al. 1998; Kyoizuka et al. 2000; Yamaguchi et al. 2004; Yamaguchi et al. 2006; Li et al. 2011a; Yan et al. 2015; Zheng et al. 2015), *OsMADS13* and *OsMADS21*, to Class D (Lopez-Dee et al. 1999; Dreni et al. 2007; Li et al. 2011a), *OsMADS1*, *OsMADS5*, *OsMADS6*, *OsMADS7*, *OsMADS8*, *OsMADS34*, *MJ706*,

and *EMF2B*, to Class E (Chung et al. 1994; Jeon et al. 2000; Agrawal et al. 2005; Kalika et al. 2005; Sun and Zhou 2008; Cui et al. 2010; Gao et al. 2010; Li et al. 2011b; Lin et al. 2013; Conrad et al. 2014). The function analysis of these genes is very helpful to elucidate the molecular regulation mechanism of rice inflorescence.

It was also known that the divergence of monocot and dicot happened about 200 million years ago (Wolfe et al. 1989). However, the morphological characteristics and development process of monocot flower are distinct from those of dicot flower. For example, the lemma, palea, and lodicule around the stamen and pistil in rice floret are obviously different from the calyx and petal in dicot flower; in addition, there is a pair of sterile lemmas and rudimentary glumes remained outside each rice floret, respectively (Yoshida and Nagato 2011), which seems an unique floral organ in grass species. In some studies, it was believed that rice lodicule is similar to the petal in dicot flower (Kang et al. 1998; Ambrose et al. 2000) and the lemma and the palea amount to the calyx (Prasad et al. 2001). Besides, the sterile lemmas might be derived from the lemma (Prasad et al. 2001; Yoshida et al. 2009). Nevertheless, many questions still remain to be further studied, such as “the lemma and the palea are the same organ?”, “where do the sterile lemma and the rudimentary glume derive from?”, and “how does every gene of floral organ work in perfect union?”. In this study, we attempted to analyze the genetic characteristics of sterile lemma trait and cloned the target gene using 4 mutants with long sterile lemma in order to reveal the gene function and structure.

## MATERIAL AND METHODS

The *Indica* long sterile lemma (LSL) mutants, JF11, JF12 and JF13, were selected from a mutant population that derived from rice mature pollens induced by <sup>60</sup>Co-γ ray irradiation with 30Gy in the Rice Genetics and Breeding Laboratory of Xiamen University, China (Figure 1a). The *Japonica* LSL mutant, JNY-7, was donated from China National Center for Crop Germplasm Preservation (Figure 1a). Aijiaonante (AJNT) and Nipponbare (NIP) are the germplasm materials preserved in our lab. F<sub>1</sub> and F<sub>2</sub> populations were constructed from the crossings between JF11 and AJNT (JF11/AJNT), NIP and JF11 (NIP/JF11), JNY-7 and AJNT

→

(JNY-7/AJNT), NIP and JNY-7 (NIP/JNY-7), JF11 and JNY-7 (JF11/JNY-7) and JF11 and JF13 (JF11/JF13), respectively.



**Figure 1.** Grain phenotypes of the parents and their  $F_2$ s. (a) The parent grains, from left to right, are: AJNT, JNY-7, JF13, JF12 and JF11. White arrows show sterile lemmas and the red ones show long sterile lemmas; (b) The grains, from left to right, are JNY-7, JF11, AJNT,  $F_1$  from JNY-7/JF11 and  $F_1$  from JNY-7/AJNT, respectively; (c) The grains, from left to right, are JF11, JF13, AJNT,  $F_1$  from JF11/AJNT and  $F_1$  from JF11/JF13, respectively; (d) The grains, from left to right, are NIP, JNY-7, and  $F_1$  from NIP/JNY-7, respectively; (e) The grains, from left to right, are NIP, JF11 and  $F_1$  from AJNT/JF11, respectively.

### DNA extraction and linked SSR marker screening

DNA extraction referred to Cetyltrimethyl Ammonium Bromide (CTAB) extracting method (Wang and Fang 2003). The marker tightly linked to the target trait was followed the approach of Bulk Segregant Analysis (BSA) (Michelmore et al. 1991). In this study, DNA was extracted from individual  $F_2$  plant. DNAs from 15 individuals with LSL trait were equally mixed and developed the mutant

gene pool. Similarly, the wild type (WT) gene pool was developed as well. These 2 gene pools were subjected to screen the simple sequence repeats (SSR) markers. The polymorphic markers between the 2 gene pools might link tightly to the locus of LSL trait.

### Genotyping and mapping

The SSR markers linked tightly to LSL trait were used to genotype 700 individuals with LSL in the  $F_2$  population derived from the JF13/AJNT crossing. Linkage analysis between genotype and phenotype was conducted by MAPMAKER/EXP 3.0 (Lander et al. 1987; Lincoln et al. 1992), and the linkage map was drawn by MapChart 2.2 (Voorrips 2002).

### SSR markers and specific primers

Sequences of 522 SSR markers used in this research were downloaded from the GRAMENE website (<http://www.gramene.org/microsat/ssr.html>). According to the NIP sequence at the 2 flanks of the target locus, 72 SSR markers were designed by Primer 5.0 and synthesized by Life Technologies Corporation.

A pair of primers, XMsl-7 (Forward primer: GAATGGAGGGTTGGGTCAC; XMsl-7, and Reverse primer: CGAAGCAACGGAACGAACAC), was designed and synthesized to amplify the Open Reading Frame (ORF), its upstream and downstream sequences of *G1*.

## RESULTS AND DISCUSSION

All  $F_1$  individuals generated from the crossings JF11/AJNT, NIP/JF11, JNY-7/AJNT and NIP/JNY-7 showed the phenotype of wild type (Figures 1b,c,d,e), whereas the  $F_1$  plants from the crosses JNY-7/JF11 and JF11/JF13 showed the phenotype of long sterile lemma (Figures 1b,c). In the  $F_2$  population from the crossing of JF11/AJNT, there are 4,741 wild type individuals and 1,565 mutant individuals, respectively, presenting 3:1 segregation ratio (WT/LSL) ( $\chi^2 = 0.102$ ;  $\chi^2_{0.05} = 3.84$ ). This indicated that the LSL trait of JF11, JF13, and JNY-7 should be controlled by a recessive locus named *sl-1(t)*.

A set of 137 SSR markers with polymorphism between JF11 and AJNT was used to test the genotype of the DNA pools

of wild type and mutant. Two markers with polymorphism on the short arm of chromosome 7, RM51 and RM295, were further applied to validate the genotype of 200  $F_2$  LSL individuals. Linkage analysis between molecular marker and phenotype trait showed that the locus of *sl-1(t)* was mapped on the inner side of RM51 and RM295, and the genetic distances from the target locus to RM51 and RM295 were 13.5 and 12.7 cM, respectively. In addition, another 6 SSR markers on chromosome 7, RM20828, RM20852, RM20903, RM20948, RM21004, and RM21035, were used to fine map the *sl-1(t)*, through genotyping 700 mutants of  $F_2$  plants. The *sl-1(t)* was narrowed in the interval between RM20903 and RM20948, and the genetic distances between *sl-1(t)* and the 2 markers were 3.8 and 4.9 cM, respectively (Figure 2).


The *G1* gene (LOC\_Os07g04670) with a 831-base length ORF was deemed to confer the sterile lemma trait (Yoshida et al. 2009). In the present research, *sl-1(t)* was located in the

interval between RM20903 and RM20948, which harbored *G1*. So we speculated that the *SL-1(t)* might be allelic to *G1*.

The PCR product with 1,873 bases included an 831-base ORF, an 874-base upstream sequence, and a 168-base downstream sequence of *G1*. Allelic sequencing showed that the homologous *G1* sequences of JF11, JF12 and JF13 mutants had a same deletion with 11 bases (GACGGCGCCGC) in the exon (Figure 3a), which directly led to the event of frame-shifting mutation, while JNY-7 had a base substitution which made an arginine convert into a histidine at the site of the 117th amino residue (Figures 3b,c). The results indicated that the mutation of *G1* should be able to generate the phenotype of long sterile lemma of the 4 mutants in this study.

Rice sterile lemma was a characteristic floral organ that was considered to be a vestigial floret in rice (Yoshida et al. 2009). Some genes related to the sterile lemma have been mapped or cloned in rice. *G1* was cloned by the map-based cloning method (Yoshida et al. 2009). *Osleg* was mapped in a 207-kb region on the short arm of chromosome 7, which was close to *G1* (Chen et al. 2010). In the present study, it was believed that *sl-1(t)* gene should be allelic to *G1* based on the locus location and allelic sequencing. It was very obvious that *G1* would be a key gene for further research on rice sterile lemma.

*G1* has only 1 exon with 831 bp, which codes a ALOG domain, a distinct N-terminal DNA-binding domain with an additional zinc ribbon (Figure 4) (Yoshida et al. 2009; Iyer and Aravind 2012). The *g1-4* mutant has a 68-base deletion in the downstream of ALOG domain; *g1-2* and *g1-3* have a same base transversion in the zinc ribbon insert region and a transition in the third helix of the ALOG domain, respectively (Yoshida et al. 2009). In this research, the mutant locus, an 11-base deletion near to the mutant site of *g1-4*, happened downstream the ALOG domain (Figure 4). In the *G1* gene of JNY-7, a base transition happened at the base No. 344 in the third helix of the ALOG domain (Figure 4). In the ORF and the upstream from -1 to -824 base of *G1*, there were 8 and 10 base substitutions between *Indica* and *Japonica* rice, respectively, and 2 of the 8 mutants located at the ORF caused the changes of the amino acid sequence. The amino acid No. 133, V, becomes G, and the No. 190, M, becomes R (Figure 4). These 2 amino acids are located at the ALOG domain, but do not impact on the gene function.



**Figure 2.** The location of *sl-1(t)* in the molecular linkage map on the short arm of chromosome 7. The bar is *sl-1(t)*, and the numbers on the left of the linkage map represent genetic distance (cM).



that the sterile lemma identity should also be related to the MADS gene family.

How is the rice sterile lemma regulated? Is there a real link between ALOG domain and MADS domain? How do the 2 domains work together? Now all of these questions are little known to us. The research on gene regulation and signal path should be further conducted in the future. The present study might provide some good mutant materials and theoretical basis for further studying the key domain, gene regulation, and signal path related to rice sterile lemma.

## CONCLUSION

Rice sterile lemma is a unique outer floral organ in gramineous plants. Long sterile lemma is a recessive trait and regulated by a key gene, *G1*. The mutant genes, *sl-1(t)*, in JF11, JF12, JF13 and JNY-7 are alleles of *G1*. JF11, JF12 and JF13 have an 11-base deletion, which gives rise to a frame-shifting mutation. JNY-7 has a base substitution which triggers a missense mutation.

*G1* codes an ALOG domain, which is a distinct N-terminal DNA-binding domain with an additional zinc ribbon. In this study, we found 2 mutations. One

locates the downstream of the ALOG domain, and the other, in the third helix of the ALOG domain. These 2 mutations both affect the gene function. Also, 8 and 10 base substitutions happen in the ORF and the upstream from -1 to -824 base of *G1* between *Indica* and *Japonica* rice, respectively, and only 2 of them in the ORF lead to the change of amino acid sequence. Although these 2 polymorphisms are located at the ALOG domain, there is no impact on the gene function, which shows these polymorphism loci are not the key conservative ones for supporting the structure and function of ALOG domain.

In a word, this study makes the researcher to further understand the structure and function of the *G1* gene and provides 2 available mutants for exploring the development mechanism of rice floral organ.

## ACKNOWLEDGEMENTS

This study was supported by grants from the National Natural Science Foundation of China (31100866), the Education Science Research Projects in Fujian Province for Young and Middle-aged Teachers (JAT160014), and the Fundamental Research Funds for the Central Universities (2013121040).

## REFERENCES

- Agrawal, G. K., Abe, K., Yamazaki, M., Miyao, A. and Hirochika, H. (2005). Conservation of the E-function for floral organ identity in rice revealed by the analysis of tissue culture-induced loss-of-function mutants of the *OsMADS1* gene. *Plant Molecular Biology*, 59, 125-135. <http://dx.doi.org/10.1007/s11103-005-2161-y>.
- Ambrose, B. A., Lerner, D. R., Ciceri, P., Padilla, C. M. P., Yanofsky, M. F. and Schmidt, R. J. (2000). Molecular and genetic analyses of the *Silky1* gene reveal conservation in floral organ specification between eudicots and monocots. *Molecular Cell*, 5, 569-579. [http://dx.doi.org/10.1016/S1097-2765\(00\)80450-5](http://dx.doi.org/10.1016/S1097-2765(00)80450-5).
- Angenent, G. C., Franken, J., Busscher, M., van Dijken, A., van Went, J. L., Dons, H. J. and van Tunen, A. J. (1995). A novel class of MADS box genes is involved in ovule development in petunia. *The Plant Cell*, 7, 1569-1582. <http://dx.doi.org/10.1105/tpc.7.10.1569>.
- Bowman, J. L., Smyth, D. R. and Meyerowitz, E. M. (1991). Genetic interactions among floral homeotic genes of *Arabidopsis*. *Development*, 112, 1-20.
- Carpenter, R. and Coen, E. S. (1990). Floral homeotic mutations produced by transposon-mutagenesis in *Antirrhinum majus*. *Genes and Development*, 4, 1483-1493. <http://dx.doi.org/10.1101/gad.4.9.1483>.
- Chen, D. B., Zhan, X. D., Chao, W. U., Shen, X. H., Wei-Ming, W. U., Gao, Z. Q. and Cheng, S. H. (2010). Genetic analysis and gene mapping of a long empty glumes mutant in rice (*Oryza sativa* L.). *Acta Agronomica Sinica*, 36, 1506-1511.
- Chung, Y. Y., Kim, S. R., Finkel, D., Yanofsky, M. F. and An, G. (1994). Early flowering and reduced apical dominance result from ectopic expression of a rice MADS box gene. *Plant Molecular Biology*, 26, 657-665. <http://dx.doi.org/10.1007/BF00013751>.

- Chung, Y. Y., Kim, S. R., Kang, H. G., Noh, Y. S., Min, C. P., Finkel, D. and An, G. (1995). Characterization of two rice MADS box genes homologous to *GLOBOSA*. *Plant Science*, 109, 45-56. [http://dx.doi.org/10.1016/0168-9452\(95\)04153-L](http://dx.doi.org/10.1016/0168-9452(95)04153-L).
- Coen, E. S. and Carpenter, R. (1993). The metamorphosis of flowers. *The Plant Cell*, 5, 1175-1181. <http://dx.doi.org/10.1105/tpc.5.10.1175>.
- Coen, E. S. and Meyerowitz, E. M. (1991). The war of the whorls: genetic interactions controlling flower development. *Nature*, 353, 31-37. <http://dx.doi.org/10.1038/353031a0>.
- Colombo, L., Franken, J., Koetje, E., van Went, J., Dons, H. J., Angenent, G. C. and van Tunen, A. J. (1995). The petunia MADS box gene *FBP11* determines ovule identity. *The Plant Cell*, 7, 1859-1868. <http://dx.doi.org/10.1105/tpc.7.11.1859>.
- Conrad, L. J., Khanday, I., Johnson, C., Guiderdoni, E., An, G., Vijayraghavan, U. and Sundaresan, V. (2014). The polycomb group gene *EMF2B* is essential for maintenance of floral meristem determinacy in rice. *The Plant Journal*, 80, 883-894. <http://dx.doi.org/10.1111/tpj.12688>.
- Cui, R., Han, J., Zhao, S., Su, K., Wu, F., Du, X., Xu, Q., Chong, K., Theissen, G. and Meng, Z. (2010). Functional conservation and diversification of class E floral homeotic genes in rice (*Oryza sativa*). *The Plant Journal*, 61, 767-781. <http://dx.doi.org/10.1111/j.1365-313X.2009.04101.x>.
- Dreni, L., Jacchia, S., Fornara, F., Fornari, M., Ouwerkerk, P. B. F., An, G., Colombo, L. and Kater, M. M. (2007). The D-lineage MADS-box gene *OsMADS13* controls ovule identity in rice. *The Plant Journal*, 52, 690-699. <http://dx.doi.org/10.1111/j.1365-313X.2007.03272.x>.
- Fornara, F., Pařenicová, L., Falasca, G., Pelucchi, N., Masiero, S., Ciannamea, S., Lopez-Dee, Z., Altamura, M. M., Colombo, L. and Kater, M. M. (2004). Functional characterization of *OsMADS18*, a member of the *AP1/SQUA* subfamily of MADS box genes. *Plant Physiology*, 135, 2207-2219. <http://dx.doi.org/10.1104/pp.104.045039>.
- Galimba, K. D., Tolkin, T. R., Sullivan, A. M., Melzer, R., Theißen, G. and Di Stilio, V. S. (2012). Loss of deeply conserved C-class floral homeotic gene function and C- and E-class protein interaction in a double-flowered ranunculid mutant. *Proceedings of the National Academy of Sciences of the United States of America*, 109, E2267-E2275. <http://dx.doi.org/10.1073/pnas.1203686109>.
- Gao, X., Liang, W., Yin, C., Ji, S., Wang, H., Su, X., Guo, C., Kong, H., Xue, H. and Zhang, D. (2010). The *SEPALLATA*-like gene *OsMADS34* is required for rice inflorescence and spikelet development. *Plant Physiology*, 153, 728-740. <http://dx.doi.org/10.1104/pp.110.156711>.
- Honma, T. and Goto, K. (2001). Complexes of MADS-box proteins are sufficient to convert leaves into floral organs. *Nature*, 409, 525-529. <http://dx.doi.org/10.1038/35054083>.
- Iyer, L. M. and Aravind, L. (2012). ALOG domains: provenance of plant homeotic and developmental regulators from the DNA-binding domain of a novel class of DIRS1-type retrotransposons. *Biology Direct*, 7, 39. <http://dx.doi.org/10.1186/1745-6150-7-39>.
- Jeon, J. S., Jang, S., Lee, S., Nam, J., Kim, C., Chung, Y. Y., Kim, S. R., Lee, Y. H., Cho, Y. G. and An, G. (2000). *Leafy hull sterile1* is a homeotic mutation in a rice MADS box gene affecting rice flower development. *The Plant Cell*, 12, 871-884. <http://dx.doi.org/10.1105/tpc.12.6.871>.
- Kalika, P., Sriram, P. and Usha, V. (2005). *OsMADS1*, a rice MADS-box factor, controls differentiation of specific cell type in lemma and plea and is an early-acting regulator of inner floral organs. *The Plant Journal*, 43, 915-928. <http://dx.doi.org/10.1111/j.1365-313X.2005.02504.x>.
- Kang, H. G., Jeon, J. S., Lee, S., Lee, S. H. and An, G. (1998). Identification of class B and class C floral organ identity genes from rice plants. *Plant Molecular Biology*, 38, 1021-1029. <http://dx.doi.org/10.1023/A:1006051911291>.
- Kang, H. G., Noh, Y. Y., Chung, Y. Y., Costa, M. A., An, K. and An, G. (1995). Phenotypic alterations of petal and sepal by ectopic expression of a rice MADS box gene in tobacco. *Plant Molecular Biology*, 29, 1-10. <http://dx.doi.org/10.1007/BF00019114>.
- Kyozuka, J., Kobayashi, T., Morita, M. and Shimamoto, K. (2000). Spatially and temporally regulated expression of rice MADS box genes with similarity to *Arabidopsis* class A, B and C genes. *Plant and Cell Physiology*, 41, 710-718. <http://dx.doi.org/10.1093/pcp/41.6.710>.
- Lander, E. S., Green, P., Abrahamson, J., Barlow, A., Daly, M. J., Lincoln, S. E. and Newburg, L. (1987). MAPMAKER: an interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. *Genomics*, 1, 174-181. [http://dx.doi.org/10.1016/0888-7543\(87\)90010-3](http://dx.doi.org/10.1016/0888-7543(87)90010-3).
- Li, H., Liang, W., Hu, Y., Zhu, L., Yin, C., Xu, J., Dreni, L., Kater, M. M. and Zhang, D. (2011b). Rice *MADS6* interacts with the floral homeotic genes *SUPERWOMAN1*, *MADS3*, *MADS58*, *MADS13*, and *DROOPING LEAF* in specifying floral organ identities and meristem fate. *The Plant Cell*, 23, 2536-2552. <http://dx.doi.org/10.1105/tpc.111.087262>.

- Li, H., Liang, W., Yin, C., Zhu, L. and Zhang, D. (2011a). Genetic interaction of *OsMADS3*, *DROOPING LEAF*, and *OsMADS13* in specifying rice floral organ identities and meristem determinacy. *Plant Physiology*, 156, 263-274. <http://dx.doi.org/10.1104/pp.111.172080>.
- Lim, J., Moon, Y. H., An, G. and Jang, S. K. (2000). Two rice MADS domain proteins interact with *OsMADS1*. *Plant Molecular Biology*, 44, 513-527. <http://dx.doi.org/10.1023/A:1026517111843>.
- Lin, X., Wu, F., Du, X., Shi, X., Liu, Y., Liu, S., Hu, Y., Theissen, G. and Meng, Z. (2013). The pleiotropic *SEPALLATA*-like gene *OsMADS34* reveals that the 'empty glumes' of rice (*Oryza sativa*) spikelets are in fact rudimentary lemmas. *The New Phytologist*, 202, 689-702. <http://dx.doi.org/10.1111/nph.12657>.
- Lincoln, S., Daly, M. and Lander, E. (1992). Constructing genetic maps with MAPMAKER/EXP 3.0. 3. ed. Cambridge: Whitehead Institute for Biomedical Research.
- Lopez-Dee, Z. P., Wittich, P., Enrico, P. M., Rigola, D., Del Buono, I., Gorla, M. S., Kater, M. M. and Colombo, L. (1999). *OsMADS13*, a novel rice MADS-box gene expressed during ovule development. *Developmental Genetics*, 25, 237-244. [http://dx.doi.org/10.1002/\(SICI\)1520-6408\(1999\)25:3<237::AID-DVG6>3.0.CO;2-L](http://dx.doi.org/10.1002/(SICI)1520-6408(1999)25:3<237::AID-DVG6>3.0.CO;2-L).
- Mena, M., Ambrose, B. A., Meeley, R. B., Briggs, S. P., Yanofsky, M. F. and Schmidt, R. J. (1996). Diversification of C-function activity in maize flower development. *Science*, 274, 1537-1540. <http://dx.doi.org/10.1126/science.274.5292.1537>.
- Michelmore, R. W., Paran, I. and Kesseli, R. V. (1991). Identification of markers linked to disease-resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions by using segregating populations. *Proceedings of the National Academy of Sciences of the United States of America*, 88, 9828-9832. <http://dx.doi.org/10.1073/pnas.88.21.9828>.
- Moon, Y. H., Jung, J. Y., Kang, H. G. and An, G. (1999). Identification of a rice *APETALA3* homologue by yeast two-hybrid screening. *Plant Molecular Biology*, 40, 167-177. <http://dx.doi.org/10.1023/A:1026429922616>.
- Pelaz, S., Ditta, G. S., Baumann, E., Wisman, E. and Yanofsky, M. F. (2000). B and C floral organ identity functions require *SEPALLATA* MADS-box genes. *Nature*, 405, 200-203. <http://dx.doi.org/10.1038/35012103>.
- Prasad, K., Sriram, P., Kumar, C. S., Kushalappa, K. and Vijayraghavan, U. (2001). Ectopic expression of rice *OsMADS1* reveals a role in specifying the lemma and palea, grass floral organs analogous to sepals. *Development Genes and Evolution*, 211, 281-290. <http://dx.doi.org/10.1007/s004270100153>.
- Rensing, S. A., Lang, D., Zimmer, A. D., Terry, A., Salamov, A., Shapiro, H., Nishiyama, T., Perroud, P. F., Lindquist, E. A., Kamisugi, Y., Tanahashi, T., Sakakibara, K., Fujita, T., Oishi, K., Shin-I, T., Kuroki, Y., Toyoda, A., Suzuki, Y., Hashimoto, S. I., Yamaguchi, K., Sugano, S., Kohara, Y., Fujiyama, A., Anterola, A., Aoki, S., Ashton, N., Barbazuk, W. B., Barker, E., Bennetzen, J. L., Blankenship, R., Cho, S. H., Dutcher, S. K., Estelle, M., Fawcett, J. A., Gundlach, H., Hanada, K., Heyl, A., Hicks, K. A., Hughes, J., Lohr, M., Mayer, K., Melkozernov, A., Murata, T., Nelson, D. R., Pils, B., Prigge, M., Reiss, B., Renner, T., Rombauts, S., Rushton, P. J., Sanderfoot, A., Schween, G., Shiu, S. H., Stueber, K., Theodoulou, F. L., Tu, H., van de Peer, Y., Verrier, P. J., Waters, E., Wood, A., Yang, L., Covem, D., Cuming, A. C., Haseb, M., Lucasm, S., Mishler, B. D., Reskim, R., Grigoriev, I. V., Quattronom, R. S. and Boorem, J. L. (2008). The *Physcomitrella* genome reveals evolutionary insights into the conquest of land by plants. *Science*, 319, 64-69. <http://dx.doi.org/10.1126/science.1150646>.
- Schmidt, R. J., Veit, B., Mandel, M. A., Mena, M., Hake, S. and Yanofsky, M. F. (1993). Identification and molecular characterization of *ZAG1*, the maize homolog of the Arabidopsis floral homeotic gene *AGAMOUS*. *The Plant Cell*, 5, 729-737. <http://dx.doi.org/10.1105/tpc.5.7.729>.
- Sun, Q. and Zhou, D. X. (2008). Rice *jmjC* domain-containing gene *JMJ706* encodes H3K9 demethylase required for floral organ development. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 13679-13684. <http://dx.doi.org/10.1073/pnas.0805901105>.
- Voorrips, R. E. (2002). MapChart: software for the graphical presentation of linkage maps and QTLs. *Journal of Heredity*, 93, 77-78. <http://dx.doi.org/10.1093/jhered/93.1.77>.
- Wang, H., Zhang, L., Cai, Q., Hu, Y., Jin, Z., Zhao, X., Fan, W., Huang, Q., Luo, Z., Chen, M., Zhang, D. and Yuan, Z. (2015). *OsMADS32* interacts with PI-like proteins and regulates rice flower development. *Journal of Integrative Plant Biology*, 57, 504-513. <http://dx.doi.org/10.1111/jipb.12248>.
- Wang, Z. and Fang, X. (2003). Plant DNA isolation. *Molecular Plant Breeding*, 2, 281-288.
- Wolfe, K. H., Gouy, M., Yang, Y. W., Sharp, P. M. and Li, W. H. (1989). Date of the monocot-dicot divergence estimated from chloroplast DNA sequence data. *Proceedings of the National Academy of Sciences of the United States of America*, 86, 6201-6205. <http://dx.doi.org/10.1073/pnas.86.16.6201>.

- Yamaguchi, T., Lee, D. Y., Miyao, A., Hirochika, H., An, G. and Hirano, H. Y. (2006). Functional diversification of the two C-Class MADS Box Genes *OsMADS3* and *OsMADS58* in *Oryza sativa*. *The Plant Cell*, 18, 15-28. <http://dx.doi.org/10.1105/tpc.105.037200>.
- Yamaguchi, T., Nagasawa, N., Kawasaki, S., Matsuoka, M., Nagato, Y. and Hirano, H. Y. (2004). The *YABBY* gene *DROOPING LEAF* regulates carpel specification and midrib development in *Oryza sativa*. *The Plant Cell*, 16, 500-509. <http://dx.doi.org/10.1105/tpc.018044>.
- Yan, D., Zhang, X., Zhang, L., Ye, S., Zeng, L., Liu, J., Li, Q. and He, Z. (2015). *CURVED CHIMERIC PALEA 1* encoding an EMF1-like protein maintains epigenetic repression of *OsMADS58* in rice palea development. *The Plant Journal*, 82, 12-24. <http://dx.doi.org/10.1111/tpj.12784>.
- Yoshida, A., Suzuki, T., Tanaka, W. and Hirano, H. Y. (2009). The homeotic gene long sterile lemma (G1) specifies sterile lemma identity in the rice spikelet. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 20103-20108. <http://dx.doi.org/10.1073/pnas.0907896106>.
- Yoshida, H. and Nagato, Y. (2011). Flower development in rice. *Journal of Experimental Botany*, 62, 4719-4730. <http://dx.doi.org/10.1093/jxb/err272>.
- Zheng, M., Wang, Y., Wang, C., Ren, Y., Lv, J., Peng, C., Wu, T., Liu, K., Zhao, S., Liu, X., Guo, X., Jiang, L., Terzaghi, W. and Wan, J. (2015). *DEFORMED FLORAL ORGAN1* (DFO1) regulates floral organ identity by epigenetically repressing the expression of *OsMADS58* in rice (*Oryza sativa*). *The New Phytologist*, 206, 1476-1490. <http://dx.doi.org/10.1111/nph.13318>.