

**Tropical and Subtropical
Agroecosystems**
An international multidisciplinary journal

www.redalyc.org/articulo.oa?id=93912996002

Tropical and Subtropical Agroecosystems

E-ISSN: 1870-0462

ccastro@uady.mx

Universidad Autónoma de Yucatán

México

Kondamudi, Rajesh; Sri Rama Murthy, K.; Pullaiah, T.

EUPHORBIACEAE - A CRITICAL REVIEW ON PLANT TISSUE CULTURE

Tropical and Subtropical Agroecosystems, vol. 10, núm. 3, septiembre-diciembre, 2009, pp. 313-335

Universidad Autónoma de Yucatán

Mérida, Yucatán, México

Available in: <http://www.redalyc.org/articulo.oa?id=93912996002>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

***Tropical and
Subtropical
Agroecosystems***

REVIEW [REVISIÓN]

EUPHORBIACEAE - A CRITICAL REVIEW ON PLANT TISSUE CULTURE

**[EUPHORBIACEAE – UNA REVISIÓN CRÍTICA SOBRE CULTIVO DE
TEJIDOS]**

Rajesh Kondamudi¹, K. Sri Rama Murthy^{1*} and T. Pullaiah²

¹*School of Conservation Biology and Plant Biotechnology, Department of
Biotechnology, Montessori Mahila Kalasala, Vijayawada - 520 010,
Andhra Pradesh, India,*

²*Department of Botany, Sri Krishnadevaraya University, Anantapur – 515 203,
Andhra Pradesh, India. E-mail: drksrmurthy@yahoo.com*

**Corresponding author*

SUMMARY

The members of Euphorbiaceae are valuable source of different kinds of useful products like dyes, edible tubers, oil crops, furniture, agricultural implements, ornamental plants, pharmacological products, rubber, timber and aesthetic items. Micropropagation is an alternative mean of propagation that can be employed in conservation of the flora in relatively shorter time. Tissue culture is useful for multiplying and conserving the species, which are difficult to regenerate by conservation methods and save them from extinction. Cryopreservation of germplasm would help in maintaining the genetic diversity of the endangered population. Improved cell and tissue culture technologies would help in producing the active compounds *in vitro* with better productivities without cutting down the natural resources. There is sufficient progress at research level to suggest that the tissue culture of Euphorbiaceae can and should be further developed. This review emphasizes the *in vitro* manipulation and remarkable achievements with biotechnology in this family made during the last six decades.

Key words: Euphorbiaceae; plant tissue culture; micropropagation.

RESUMEN

Los miembros de la familia Euphorbiaceae son una fuente valiosa de diversos productos valiosos como tintes, tubérculos comestibles, aceites, implementos agrícolas, plantas ornamentales, productos farmacológicos, lates, madera y productos estéticos. La micropropagación es una herramienta que puede ser empleada como una herramienta para la conservación de la flora en un lapso de tiempo menor. El cultivo de tejidos es útil para multiplicar y conservar especies, las cuales son difíciles de regenerar con otros métodos de conservación y salvarlas así de la extinción. La criopreservación del germoplasma ayudaría a mantener la diversidad genética de población en peligro. Tecnologías mejoradas para el cultivo de células y tejidos ayudaría a producir compuestos activos *in vitro* con una mejor productividad y sin afectar los recursos naturales. Existe suficiente progreso en materia de investigación que sugiere que el cultivo de tejidos de Euphorbiaceae puede y debe ser desarrollada. Esta revisión enfatiza la manipulación *in vitro* y los logros que la biotecnología ha logrado en esta familia en las últimas seis décadas.

Palabras clave: Euphorbiaceae; cultivo de tejidos vegetales; micropropagación.

INTRODUCTION

Researchers are constantly exploring new evidence for natural resources. A lot of economical factors are based on our resources which makes a transition from one element to another very difficult. The family Euphorbiaceae comprises nearly 322 genera and 8910 species (Bingtao Li *et al.*, 2008) many of which have their own economic value and hence contribute to the floristic wealth of tropical and subtropical countries of the world. The family comprises a number of endemic

and endangered taxa. However the *in vitro* studies are confined only to a few genera of aesthetic, medicinal, timber yielding, rubber yielding, dye yielding, cottage industries, ornamental and food crops like *Acalypha*, *Baliospermum*, *Codiaeum*, *Cleistanthus*, *Croton*, *Euphorbia*, *Embllica*, *Eryngium*, *Excoecaria*, *Givotia*, *Glochidion*, *Hevea*, *Jatropha*, *Mallotus*, *Manihot*, *Phyllanthus*, *Putranjiva*, *Ricinus*, *Sapium* and *Uapaca*.

Major components of *Euphorbia* latex are sterols, terpenoids vitamins and insecticides and anti cancer drugs (Stohs and Rosenberg, 1975; Heftmann, 1975; Deshmukh and Borle 1975; Biesboer and Mahlberg, 1979; Yamamoto *et al.*, 1981; Saigo and Saigo, 1983; Itokawa *et al.*, 1989; Wu *et al.*, 1991; Rani *et al.*, 2003). Abel-Fattah and Rozk (1987) published on chemical constituents and economic important plants of Euphorbiaceae. The family is also well reputed for the production of valuable secondary metabolites like alkaloids and flavonoids in nature (Puebla *et al.*, 2003; Maciel *et al.*, 1998; Martin *et al.*, 2002; Rani *et al.*, 2002). Several of *Phyllanthus* species produce useful secondary metabolites, which have been extracted from whole plants (Unander, 1996). The species of *Phyllanthus* contain alkaloids, tannins, flavonoids, lignans, phenols, terpenes and show antinociceptive action in mice and other therapeutic activities (Filho *et al.*, 1996). The members of the family are rich in reduced hydrocarbon materials that can be extracted and converted to petroleum like compounds (Tidemann and Hawker, 1982). Cracked or fermented latex can be used as fuel (Nielsen *et al.*, 1977; Calvin, 1980; Depeyre *et al.*, 1994). Depletion of petroleum resources is creating an opportunity for exploitation of vegetable oils as bio-diesel. All over the world major sources of diesel include rape seed (USA), sun flower (Italy and South France), soybean (USA and Brazil), oil palm (Malaysia), lin seed (Spain), cotton seed (Greece), beef tallow (Ireland) and *Jatropha* (Nicaragua and South Africa) (Jayasingh, 2004).

Jatropha oil is a real bio product, CO₂ neutral, non toxic, bio degradable, free of sulphur and chlorine, caloric equivalently to mineral oil; *Jatropha* oil is non consumable, 100% pure biomass and as a result, CO₂ neutral (green) energy production. Global castor seed production is around 1 million tons per year. Leading producing areas are India (with over 60% of the global yield), China and Brazil. (Arifin *et al.*, 2008)

The latex of *Euphorbia pulcherrima* has been reported to be poisonous to livestock (Anonymous, 1978) however, in veterinary medicine it is used to kill Maggots in the wounds of livestock. *Euphorbia lagascae* is a spurge which is present wildy at South Eastern Spain, it produces 50% of seed oil with 60% of cis 12, 13-epoxy oleic or Vernolic acid (Kleiman *et al.*, 1964). Present objective for breeding in *Euphorbia lagascae* would be the reduction of skin irritant compounds in the latex (Turley *et al.*, 2000). The leaf extracts of this family are reported to have many medicinal properties including purgative, sedative, antifungal, antiamoebic and anti cancerous activities (Desmukh and Borle, 1975; Kupchan *et al.*, 1976). *Jatropha curcas*, a multipurpose plant, is valued not only for its medicinal properties and resistance to

various stresses but also for its use as an oil (curcas) seed crop (Heller, 1996; Openshaw, 2000). The seeds press cake and oil of *Jatropha curcas* cannot be used for human or animal consumption but can be used as organic fertilizer due to the presence of several toxic substances including a lectin (curcin) phorbol esters, saponins, protease inhibitors and phytates (Makkar *et al.*, 1998). Antiviral effects against hepatitis-B virus and possibly against the reverse transcriptase of retroviruses have also been reported (Thyagarajan *et al.*, 1988; Shead *et al.*, 1992). Pharmacological studies carried out with callus extracts of *Phyllanthus niruri*, *P. tenellus* and *P. urinaria* have shown antinociceptive properties and the main compounds identified in the extracts were flavonoids, tannins and phenols (Santos *et al.*, 1994). In Brazil, infusion of leaves, stems and roots of *Phyllanthus* species are used in folk medicine for treating intestinal infections, diabetes, Hepatitis-B and disturbances of kidney and urinary bladder (Calixto *et al.*, 1998). Additional studies on callus and root extracts of these different species have shown the presence of phyllembilin, tannin that has antimicrobial activity. Hydrolysable tannins which inhibited DNA polymerase and reverse transcriptase of geraniin and its derivatives which showed high activity in the inhibition of Human Immunodeficiency Virus (HIV) reverse transcriptase and angiotensin converting enzyme involved in diabetic complications (Ueno *et al.*, 1988; Ogata *et al.*, 1992; Unander, 1996). One of the main impediments in tissue culture studies in this family is the presence of latex.

IN VITRO MANIPULATIONS

Murashige and Skoog (1962) medium was mostly used to initiate and improve the response in *in vitro* cultures, White's basal medium, Woody plant medium and B₅ media (Gomborg's medium) have also been employed in some of the cases. Now a day, the transgenic plants produced through tissue culture methods showed superior abiotic and biotic stress tolerance (Ganesan and Jayabalan, 2006). Roy and Jinnah (2001) studied the *in vitro* propagation of poinsettias (*Euphorbia pulcherrima*). The hormonal control of triterpinols synthesis in *Euphorbia characias* calli was studied by Ferriera *et al.*, (1992). Successful *in vitro* vegetative propagation has been reported for *Euphorbia* species (Lang he *et al.*, 1974; Lee *et al.*, 1982; Jakobek *et al.*, 1986; Zhang *et al.*, 1987; Nielsen *et al.*, 2003). Few studies available on the tissue culture of *Phyllanthus* species are on callus culture of *P. emblica*, *P. urinaria*, *P. amarus*, *P. abnormis*, *P. carolinensis*, *P. tenellus*, *P. niruri* and on transformed root cultures of *P. niruri* (Khanna and Nag, 1973; Unander, 1991; Ishimaru *et al.*, 1992; Santos *et al.*, 1994). *Croton* was chosen for micropropagation due to its rare success in conventional breeding and very little data is available

for its *in vitro* production (Shibata *et al.*, 1996; Orlikowska *et al.*, 1995; Orlikowska *et al.*, 2000). Regarding castor bean regeneration, few reports were available for whole plant regeneration through shoot tip and embryonal axis culture (Sujatha and Reddy, 1998) and genetic transformation through direct regeneration methods (Sujatha and Sailaja, 2005). Carron and Enjalric (1983), Sompong and Muangkaewangam (1992) and Asokan *et al.*, (1988) gave protocols for micropropagation of *Hevea brasiliensis* and Ferriere *et al.*, (1992) focused his

research on origin and ontogenesis of somatic embryos. Qin *et al.* (2006); Compos *et al.* (2007) studied the tissue culture and plant regeneration of *Jatropha curcus* and *Jatropha elliptica* respectively. Chitra and Madhusoodanan (2005) studied the influence of auxins in direct *in vitro* morphogenesis of *Euphorbia nivulia*. Rout *et al.*, (2006) reviewed critically on present scenario and future prospects of tissue culture of some Euphorbiaceae members. In Table 1 a summary of tissue culture research is presented.

Table 1. Tissue culture studies in Euphorbiaceae.

Taxa	Type of culture	Source of Explant	Result	Reference
<i>Baliospermum montanum</i> (Willd.) Mull-Arg	Node	Node, <i>in vitro</i> shoots	Shoot proliferation, root initiation.	Johnson and Manickam, 2003
<i>Cleistanthus collinus</i> Roxb.	Shoot bud	Nodal explants	Apical and axillary shoot proliferation, rooted <i>in vitro</i> ,	Quraishi and Mishra, 1998
<i>Codiaeum variegatum</i> L.	Shoot tip.	Shoots	Shooting, roots induced.	Asma Nasib <i>et al.</i> , 2008
<i>Codiaeum variegatum</i> L.	Leaf	Leaf explants	Somatic embryos, adventitious buds, Shoot multiplication, rooted <i>in vitro</i> .	Marconi and Radice, 1997
<i>Codiaeum variegatum</i> L.	Shoot	Shoots	Axillary shoot proliferation (increased when shoot tips were removed and inverted).	Orlikowska <i>et al.</i> , 2000.
<i>Croton sublyratus</i> Kurz.	Shoot tip	Shoots	Axillary shoot proliferation, shoot regeneration.	Shibata <i>et al.</i> , 1996
<i>Croton urucurana</i> Baill.	Callus	Leaf segments	Callus induced	Lima <i>et al.</i> , 2008
<i>Emblica officinalis</i> Gaertn.	Callus	Mature endosperm	Shoots, embryo like bodies, plantlets.	Sehgal and Khurana., 1985
<i>Emblica officinalis</i> Gaertn.	Callus	Mature embryo	Embryos, multiple shoots, rooted <i>in vitro</i> .	Tyagi and Govil, 1999
<i>Euphorbia antisyphilitica</i> Zucc.	Shoot	Shoot explant	Shoot proliferation	Jakobek <i>et al.</i> , 1986
<i>Euphorbia characias</i> L.	Callus	Callus	Deferent levels of triterpinols	Ferriera <i>et al.</i> , 1992
<i>Euphorbia esula</i> L.	Callus and rhizobacteria	Callus	Cell alterations noticed.	Souissi <i>et al.</i> , 1996
<i>Euphorbia esula</i> L.	Hypocotyl segment	Hypocotyls / all the parts.	Studied the effect of auxins, cytokinins,	Davis and Olson, 1993

Taxa	Type of culture	Source of Explant	Result	Reference
<i>Euphorbia esula</i> L.	Cell suspension	Stem callus, hypocotyls, Seedling root segments,	Organogenesis, plant lets, shoots, rooted <i>in vitro</i>	Davis <i>et al.</i> , 1988
<i>Euphorbia lagascae</i> Spreng	Bud, shoot	Apical shoot, axillary shoot	Rooting	Torres, 2004
<i>Euphorbia lathyris</i> L.	Shoot tips	Shoot tip	Adventitious shoot and organogenesis etiolated micro shoot rooting.	Ripley and Preece, 1986
<i>Euphorbia nivulia</i> Buch.-Ham.	Mesophyll	Mesophyll cells	Somatic embryos, multiple shoots, rooted <i>in vitro</i> .	Martin <i>et al.</i> , 2005
<i>Euphorbia pulcherrima</i> Willd.	Node	Node	Somatic embryogenesis	Jasrai <i>et al.</i> , 2003
<i>Euphorbia pugniformis</i>	Shoot	Tubercle	Shoot proliferation Crestate shoot proliferation	Balotis and Papafotiou, 2003
<i>Euphorbia tirucalli</i> L.	Axillary bud	Inter node	Adventitious bud proliferation, shoot regeneration	Uchida <i>et al.</i> , 2004
<i>Excoecaria agallocha</i> L.	Node	Node	Axillary sprouting	Rao <i>et al.</i> , 1998
<i>Givotia rottleriformis</i> Griff.	Embryo	Zygotic embryo	Callus, seedlings, shoots, rooted <i>in vitro</i>	Rambabu <i>et al.</i> , 2006
<i>Givotia rottleriformis</i> Griff.	Seed	Seeds	Seed germination	Rambabu <i>et al.</i> , 2005
<i>Glochidion multiloculare</i> Mull-Arg.	Callus, shoot tip	Shoot apical meristem, nodes	Multiple shoots, rooted <i>in vitro</i> .	Yamuna <i>et al.</i> , 1995
<i>Hevea brasiliensis</i> Müll.Arg.	Callus	Callus	Embryogenesis	Ferriere <i>et al.</i> , 1992
<i>Hevea brasiliensis</i> Müll.Arg.	Anther	Immature anther	Somatic embryogenesis, callus induction, plant regeneration.	Jayasree <i>et al.</i> , 1999
<i>Hevea brasiliensis</i> Müll.Arg.	Immersion culture	Embryogenic callus	Somatic embryogenesis	Etienne <i>et al.</i> , 1997
<i>Hevea brasiliensis</i> Müll.Arg.	Callus	Tissue	CaCl ₂ effect, embryogenic calli, friable calli.	Montoro <i>et al.</i> , 1995
<i>Hevea brasiliensis</i> Müll.Arg.	Somatic embryos	Somatic embryos	Observed the effects of desiccation, osmolarity of the medium and ABA on maturation.	Etienne <i>et al.</i> , 1993
<i>Jatropha curcas</i> L.	Node	Node	Axillary shoot proliferation	Datta <i>et al.</i> , 2007
<i>Jatropha curcas</i> L.	Axillary bud	Leaf segments, axillary buds, nodes	Adventitious shoots, multiple shoots.	Sujatha <i>et al.</i> , 2005
<i>Jatropha curcas</i> L.	Hypocoty, petiole and leaf bits.	Hypocotyls, petiole and leaf bits	Regeneration of adventitious shoots, rooted <i>in vitro</i> .	Sujatha and Mukta, 1996

Taxa	Type of culture	Source of Explant	Result	Reference
<i>Jatropha curcas</i> L.	Shoot tip	Shoot tip	Shooting rooted <i>in vitro</i>	Rajore and Batra, 2005
<i>Jatropha elliptica</i> (Pohl) Müll.Arg.	Nodal multiplication	Nodes	Shooting, rooting, acclimatization	Campos <i>et al.</i> , 2007
<i>Mallotus repandus</i> (Willd.) Müll.Arg	Node, inter node	Node, inter node	Shoot induction, Shoot elongation, organogenesis.	Prathanaturug <i>et al.</i> , 2007
<i>Mallotus repandus</i> (Willd.) Müll. Arg	Node	Nodal, shoots	Shoot formation, rooted <i>in vitro</i> ,	Kaewsuwan <i>et al.</i> , 2005
<i>Manihot esculenta</i> Crantz	Somatic embryo	Somatic embryo	Shoot organogenesis, somatic embryogenesis	Ma, 1998.
<i>Manihot esculenta</i> Crantz	Leaf	Young leaf lobes	Induction of somatic embryos, germination of somatic embryos shoot development	Raemakers <i>et al.</i> , 1993 b
<i>Manihot esculenta</i> Crantz	Apical meristem	Apical meristem	Virus free plants	Kartha <i>et al.</i> , 1974
<i>Manihot esculenta</i> Crantz	Leaf	Leaf explant	Direct and indirect somatic embryogenesis	Raemakers <i>et al.</i> , 2000
<i>Manihot esculenta</i> Crantz	Somatic embryo culture	Somatic embryo	Somatic embryo culture Differentiated, developed and germinated.	Groll <i>et al.</i> , 2002
<i>Manihot esculenta</i> Crantz	Root	Primary root	Observed the ability of primary root storage.	Ricardo <i>et al.</i> , 2007
<i>Manihot esculenta</i> Crantz	Axillary bud	Mononodal micro cutting	Plants were recovered.	Rommanee <i>et al.</i> , 2003
<i>Manihot esculenta</i> Crantz	Protoplast	Leaf	Protoplast division was observed.	Anthony <i>et al.</i> , 1995
<i>Manihot esculenta</i> Crantz	Cryopreserved shoot tips	<i>In vitro</i> grown shoot tips	Regeneration of shoots	Escobar <i>et al.</i> , 1997
<i>Manihot esculenta</i> Crantz	Primary embryo culture	Leaf lobes	Production of embryos	Raemakers <i>et al.</i> , 1993b
<i>Manihot esculenta</i> Crantz	Axillary bud culture	Node, axillary buds, meristems	Formation of shoots and multiple shoots.	Konan <i>et al.</i> , 1997
<i>Manihot esculenta</i> Crantz	Node	Node	Multiple shoots	Bhagwat <i>et al.</i> , 1996
<i>Manihot esculenta</i> Crantz	Cotyledon	Cotyledons	Somatic embryogenesis	Konan <i>et al.</i> , 1994
<i>Manihot esculenta</i> Crantz	Callus	Floral tissue	Somatic embryogenesis	Woodward and Kaerlas., 2001
<i>Manihot esculenta</i> Crantz	Leaf lobes	Young leaf lobes	Somatic embryos, shoot and root axis.	Mathews <i>et al.</i> , 1993
<i>Manihot esculenta</i> Crantz	Node, Shoot apices	Shoot apices, nodes	Callus formation, shoot formation.	Villaluz, 2006

Taxa	Type of culture	Source of Explant	Result	Reference
<i>Manihot esculenta</i> Crantz	Meristems	Meristem	<i>In vitro</i> flowering	Tang <i>et al.</i> , 1983
<i>Manihot esculenta</i> Crantz	Leaf	Immature leaves	Adventitious Shoot, somatic embryogenesis	Ma and Xu, 2002
<i>Phyllanthus amarus</i> Schum. & Thonn.	Shoot, node and inter node culture.	Shoot tip, node, inter node.	Multiple shoots, rooted <i>in vitro</i>	Ghanti <i>et al.</i> , 2004
<i>Phyllanthus amarus</i> Schum. & Thonn.	Synthetic seed	Shoot tip (alginic encapsulate)	Growth reduced plantlet formed	Singh <i>et al.</i> , 2006
<i>Phyllanthus amarus</i> Schum. & Thonn.	Shoot tip	Shoot tip	Multiple shoots, rooted <i>in vitro</i>	Bhattacharyya and Bhattacharya, 2001
<i>Phyllanthus beddomii</i> (Gamble) Mohanan	Seed germination	Seeds	Plantlet	Maridass and Thangavel, 2008.
<i>Phyllanthus coroliniensis</i> Walter	Axillary shoot	Nodal segments	Multiple shoots rooting initiated.	Catapan <i>et al.</i> , 2000
<i>Phyllanthus fraternus</i> Webster	Shoot tip, nodal	Shoot tips, nodal parts	Multiple shoots, axillary bud sprouting	Rajasubramaniam and Saradhi, 1997
<i>Phyllanthus niruri</i> L.	Node	Nodal segments	Shoots, multiple shoots, flowers and fruiting	Liang and Keng, 2006
<i>Phyllanthus stipulatus</i> Raf.	Axillary bud	Nodal segment	Callus induction, micro shoots, multiple shoots, rooted <i>in vitro</i> , <i>ex-vitro</i> flowering, regeneration.	Catapan <i>et al.</i> , 2001
<i>Phyllanthus urinaria</i> L.	Node and Callus	Node	Callus induction, multiple shoots, rooted <i>in vitro</i>	Catapan <i>et al.</i> , 2002
<i>Ricinus communis</i> L.	Cotyledon culture	Cotyledon	Shoot proliferation, multiple shoots, callus, <i>in vitro</i> rooted,	Kumari <i>et al.</i> , 2008
<i>Ricinus communis</i> L.	Bud culture	Apical and axillary buds	Shoot regeneration, Rooted <i>in vitro</i> , elongated.	Molina and Schobert, 1995
<i>Ricinus communis</i> L.	Shoot tip	Meristematic explant	Deferential cytokinin effects were studied.	Sujatha and Reddy, 1998

Seed germination

Sterilization is must for the seeds prior to inoculation in order to avoid the contamination. Jamal *et al.* (1993) studied the negative effect of 2,4-Dichlorophenoxy acetic acid (2,4-D) on seed germination of the *Euphorbia esula*. In *Givotia rottleriformis* Rambabu *et al.*, (2005) studied the effect of Gibberellic acid (GA₃) on enhancement of *in vivo* seed germination. Thapliyal (2004) reported twin seedlings in *Putranjiva roxburghii*. The transmission of juvenile rooting

ability from seedlings to adults of *Hevea brasiliensis* was studied by Muzik and Cruzado (1958).

Cotyledonary nodes and hypocotyl tissue

Ricinus communis cotyledons, cultured on the medium with Murashige and Skoog's medium salts (MS) + B₅ Vitamins, resulted in the formation of multiple shoots, callusing, roots and the plantlets were later acclimatized (Kumari *et al.*, 2008). Hypocotyl tissue from zygotic embryo axis of *Ricinus communis* produced adventitious shoots when treated with either

Thidiazuron (TDZ) or BA (N6-Benzyladenine). While Indole butyric acid (IBA) induced the rooting very effectively and plants established in the soil (Ahn *et al.*, 2007). Molina and Schobert.(1995) studied the micropropagation of *Ricinus communis*. Cotyledinory explants of *Manihot esculenta* were first treated with 2,4-D (4mg/l^{-1}) then placed on MS + NAA (1-naphthaleneacetic acid) + BA + GA_3 , MS medium or 4 × MS micro salts were efficient for the formation of somatic embryos. 6% Sucrose was optimal concentration for the development of somatic embryos after an induction treatment using 2% sucrose, ABA (0.52 mg/l^{-1}) induced good number of somatic embryos production (Konan *et al.*, 1994). Seneviratne (1991) studied the micropropagation of juvenile and mature *Hevea brasiliensis*.

Shoot and root regeneration

Viruses makes plants to grow less by their effects like necrosis, curling of leaves, decreased yield and plant death (Quak, 1997). Shoot apical meristems and first set of primordial leaves in an elongating shoot are generally not connected to the vascular system of the plant and therefore are not contaminated by viruses as that travel through the vascular system. The confusion in the true shoot meristem consists only of the isolated apical dome without visible primordial leaves attached. The nodal segments of *Baliospermum montanum* were cultured on MS + BAP ($3.10\mu\text{M}$). *In vitro* raised shoots were rooted on half MS medium fortified with various concentrations and combinations of auxins (Johnson and Manickam, 2003). *In vitro* micropropagation of *Baliospermum axillare* was recorded by Kamalendra and Sudarshanam (2003). For the micropropagation of *Hevea brasiliensis* the shoot tips were used (Gunatilleke and Samaranyake 1988). Another important medicinal plant *Phyllanthus amarus* apical meristems were cultured on MS + Kn(6-furfurylamino purine (Kinetin))/BAP singly or in combination with Indole acetic acid (IAA). Growth regulators at lower range ($0.1\text{-}1.0\text{ mg L}^{-1}$) stimulated direct regeneration of shoots. Kn is superior to BAP and Kn + IAA combination was more suitable than Kn alone. The micropropagated plants were successfully established in the soil with high survival rate of 80% (Bhattacharyya and Bhattacharya, 2001). Ghanti *et al.*, (2004) reported high frequency shoot regeneration from shoot tip, nodal and internodal segments of *Phyllanthus amarus* and studied the effect of different concentrations of BAP, Kn and coconut milk on shoot tips, axillary and internodal segments. Hardening was done in the liquid MS medium and the survival rate of the acclimatized plants was 85%. Singh *et al.* (2006), encapsulated the shoot tips of *Phyllanthus amarus* with 3% sodium alginate and 75 mM CaCl_2 , H_2O . They were grown *in vitro* and 90 % of shoots were regenerated. There are so many factors effecting the

growth of the shoot tips like, MS salts in the sodium alginate beads, concentration of sodium alginate and storage duration etc.

In the genus *Phyllanthus* the nodal culture, callus culture and rooting were observed in many species. viz., *Phyllanthus carolinensis* (Catapan *et al.*, 2000); *Phyllanthus fraternus* (Rajasubramaniam and Saradhi, 1997) *Phyllanthus stipulatus* (Catapan *et al.*, 2001) *Phyllanthus urinaria* (Catapan *et al.*, 2002).

The nodal explants of *Manihot esculenta* were cultured on two different media, liquid MS + TDZ ($0.11\text{-}0.22\mu\text{M}$) where internodal elongation was observed, followed by solidified MS+BA ($2.2\text{ }\mu\text{M}$)+ GA_3 ($1.6\mu\text{M}$) generated clusters of buds and fasciated stems which developed into shoots (Bhagwat *et al.*, 1996). The nodes of *Manihot esculenta* were cultured on medium containing BAP (10mg / l) to get enlarged axillary buds, they gave multiple shoots on shifting to the fresh medium (Konan *et al.*, 1997). Kartha *et al.*, (1974) produced viral free plants from the Apical meristems of *Manihot utilissima*. Cryopreserved and thawed apices of *Manihot esculenta* were cultured on a osmoprotected medium. Successfully vitrified apices showed growth within a week and developed into shoots. The average recovery rate was about 70% (Rommanee *et al.*, 2003). Escobar *et al.*, (1997) presented a protocol for recovering *Manihot esculenta* plants from shoot tips maintained in liquid nitrogen. The shoot tips responded as per the pre and post freezing steps. Finally, they have succeeded to produce 50-70% of plants. Villaluz (2006) developed a protocol for the rapid meristem development and mass propagation of *Manihot esculenta* and the explants were cultured on either liquid or solid medium containing GA_3 , BAP, NAA at (0.25 , 0.1 and 0.2 mg/l) respectively. MS medium with NAA induced both shoot and root development (Ricardo *et al.*, 2007).

High frequency regeneration from various explants of *Jatropha integerrima* has been reported (Sujatha and Dhingra, 1993). Sujatha and Mukta.(1996) worked on *Jatropha curcas* to study the morphogenesis and plant regeneration from tissues. The shoot tips of the *Jatropha curcas* were cultured on the MS + BAP (2.0mg/l) and IAA (0.5mg/l) along with adenine sulphate, glutamine and activated charcoal, these shoots were rooted on the half MS + IBA ($0.5\text{-}5.0\text{mg/l}$) the high frequency rooting was on MS + IBA 3.0 mg/l . The survival rate of acclimatized plants was 60-70% (Rajore and Batra, 2005). Shoot buds from the axillary nodes and leaf segments of non-toxic *Jatropha curcas* were cultured on MS + Kn ($2.3\text{-}46.5\mu\text{M}$), BA ($2.2\text{-}44.4\mu\text{M}$) and TDZ ($2.3\text{-}45.4\mu\text{M}$) individually, the leaf segments were cultured on MS+ BA ($8.9\mu\text{M}$) + IBA ($2.5\mu\text{M}$) and got adventitious shoots (Sujatha *et*

al., 2005). Datta *et al.*, (2007) reported *in vitro* clonal propagation of biodiesel plant *Jatropha curcas*.

The shoot tips of *Glochidion multiloculare* produced multiple shoots when cultured on MS + BA (1.0 mg/l) and IAA (1.0 mg/l) callus derived from the leaf and stem explants on a medium containing 2,4-D (0.5-2.0 mg/l) produced shoot buds when transferred to MS + BA (1.0-2.0 mg/l) + CM (Coconut milk, 10% v/v) the rooting combination was MS + IBA (1.0 mg/l) and were acclimatized (Yamuna *et al.*, 1995). The shoots of *Euphorbia antisyphilitica* were cultured on modified MS medium + NAA (0.13µM) and BAP (4.44µM), the rooting combination was half MS+IBA 0.49µM and acclimatization was very simple (Jakobek *et al.*, 1986). The axillary shoots of *Euphorbia lagascae* were dipped in IBA (50 mg/l) for 2 min. IBA increased the survival rate up to 80-100% (Torres, 2004). The tips of cristate lateral shoots of *Euphorbia pugniformis* were cultured on the MS + sucrose 2% and NAA 0.1 mg/l⁻¹ and BA, gave cristate shoots of both forms. While 20% gave normal shoots. The rooting combination for cristate shoots was MS+ IBA (0.5mg/l⁻¹), 95-100% of rooted shoots were acclimatized *ex vitro*. Quite a lot of cristates reverted into the normal form. Effect of MS nitrogen salts had a considerable effect on the cristate form shoot reversion (Balotis and Papafotiou, 2003). Effects of several factors on rooting of Poinsettias was recorded by Sochacki and Chimid (1994). Acclimatization systems for micro cuttings of *Euphorbia fulgens* was developed by Zhang and Stoltz (1989). *In vitro* production of shoot buds in *Euphorbia pulcherrima* was carried out by Nataraja *et al.*, 1973.

Shibata *et al.* (1996) developed a micropropagation protocol for *Croton sublyratus* the results demonstrate that the feasibility of applying the shoot tip culture technique for enhancing production of plaunotol by cultivating uniform populations of *Croton sublyratus* with higher plaunotol level. The morphogenetic studies on cultured endosperm of *Croton banplandianum* was carried out by Bhojwani and Johri (1971).

Shoots of *Codiaeum variegatum* cultured on MS + BAP (4.4µM) and GA₃ (5.8µM) doubled the number of axillary shoots in comparison to non defoliated controls. An additional increase in shoot number was achieved when the explant shoots were placed on the medium horizontally or vertically in an inverted position with shoot tip down. Exogenous IAA decreased shoots number and 2,3,5-Triiodo benzoic acid did neither replace defoliation nor diminish the size of callus formed at the base of the shoots. Almost all the shoots obtained were able to root on the medium with IBA (1mg/l) and acclimatized (Orlikowska *et al.*, 2000). Asma Nasib *et al.*, (2008)

grew the shoot tips of *Codiaeum variegatum* on MS + BAP (0.5mg/l) + peptone (25mg/l) which enhanced the shoot formation and rooting was induced on MS + IBA (2.0mg/l) medium and they were acclimatized with 95% survival rate. Marconi and Radice (1997) studied organogenesis and somatic embryogenesis in *Codiaeum variegatum*. To study the laticiferous systems of the *Hevea*, Bouychou (1953) of the Institute Francia Caoutchouc started *in vitro* cultures first time and followed by Wilson and Street, (1975). Gunitilleke and Samaranyake (1988) used shoot tips for the micropropagation of *Hevea brasiliensis*.

Leaf cultures

The leaves of *Uapaca kirkiana*, *Uapaca nitida* and *Jatropha curcas* were co-cultured for the callus induction. The objective of this investigation was to evaluate the early signs of grafting compatibility of *Uapaca kirkiana* (Mng'omba *et al.*, 2007). The propagation of *Uapaca kirkiana* using tissue culture techniques was carried out by Maliro, (1997) and the effect of cytokinins on *in vitro* propagation of *Uapaca kirkiana* was studied by Chishimba *et al.*, (2000). A clonal propagation protocol was developed for *Uapaca kirkiana* by Mng'omba, (2007). Anthony *et al.*, (1995) developed a protocol for the leaf protoplast culture of *Manihot esculenta*. The leaf lobes of *in vitro* grown *Manihot esculenta* were cultured on MS + 2,4-D (4 mg/l) + CuSO₄ (2µM) to improve the frequency of somatic embryogenesis. They were transferred to basal medium supplemented with activated charcoal. The desiccated bipolar somatic embryos showed 92% germination and 83% plants regeneration (Mathews *et al.*, 1993). Immature leaves of *Manihot esculenta* were cultured on MS + 2,4-D or NAA to induce somatic embryogenesis and adventitious shoots. Results showed that auxin was a key factor for inducing embryogenic cells, BAP stimulated adventitious shoots. Histological examinations supported the conclusion (Ma and Xu, 2002). The leaf bits of *Croton urucurana* were cultured on woody plant medium + different concentrations of 2,4-D in combination with TDZ, BAP induced the callus, the highest callus fresh mass was observed when treated with 2,4-D (Lima *et al.*, 2008).

Nodes and internodes

Nodal meristems are an important tissue source for micropropagation and plants raised from these are comparatively more resistant to genetic variation (Pierik, 1991). Konan *et al.*, (1994) reported an efficient mass propagation system for Cassava (*Manihot esculenta*) based on nodal explants and axillary bud-derived meristems. The nodes and basal sprouts of the *Cleistanthus collinus* were cultured on + Citric acid (CA, 104.1µM) and Polyvinylpyrrolidone

(PVP) 40 (12.5 or 25 μ M) + BA (0.44 μ M) resulted in shooting. In general, the explants from basal sprouts were more suitable than terminal twig explants for the micropropagation (Qurashi and Mishra, 1998). The nodes of *Excoecaria agallocha* were cultured on MS + BA (13.3 μ M) + Zeatin (4.65 μ M) + IBA (1.23 μ M) resulted in shoot proliferation and for rooting, IBA (0.23 μ M) was suitable. The regenerated plants were successfully acclimatized with 85% survival rate (Rao *et al.*, 1998). Nodes and shoots of *Mallotus repandus* cultured on MS + BA, 2ip (6- γ , γ -dimethylallylamine) purine), Kn and BA (4.44 μ M) induced shoots, where as the roots were induced with NAA (32.23 μ M), the regenerated plants were acclimatized properly and then analyzed chemically to compare with mother plants (Kaewsuwan *et al.*, 2005).

The nodal explants of *Euphorbia pulcherrima*, cultured on MS + NAA, 2,4-D, Kn and 2ip generated the embryogenic callus which on subculture to MS medium supplemented with 2ip (9.8 μ M) and NAA (0.54 μ M) gave somatic embryoids. The embryoids germinated successfully and were acclimatized with high survival rate (Jasrai *et al.*, 2003). *Euphorbia tirucalli* inter node explants, cultured on Linsmaier and Skoog's medium (LS) + TDZ (0.02 mg/l), resulted in the induction of the adventitious buds which on further culture on LS basal medium grew into shoots. The roots were induced by growing the shoots on the LS + NAA (0.02 mg/l) followed by half LS basal medium and were successful to obtain whole plantlets (Uchida *et al.*, 2004). The nodes and internodes of *Mallotus repandus* were precultured on MS basal and then shifted to shoot induction medium comprising of BAP (4.44 μ M), the shoots were immediately shifted to shoot elongation medium. The results suggest that the explants need to acquire competence before shoot organogenesis. Rooting was obtained by incubating the regenerated shoots on half MS + NAA (10.74 μ M) and the plants were successfully transferred to soil (Prathanurug *et al.*, 2007). Sujatha *et al.* (2005) reported the proliferation of shoot bud from nodes and leaf sections of non toxic *Jatropha curcas*. Nodal explants of *Jatropha curcas* were raised on MS + BA (22.2 μ M) and adenine sulphate (55.6 μ M) and thus shoots were multiplied on MS + Kn (2.3 μ M), IBA (0.5 μ M) and adenine sulphate (27.8 μ M). For rooting, MS + IBA (1.0 μ M) and subsequent transfer to MS basal and later acclimatized. (Datta *et al.*, 2007). The effect of TDZ on nodal proliferation was recorded by Seneviratne and Flemingmann (1996).

Callus initiation

As first step in many tissue culture experiments, it is necessary to induce callus from the primary explant, Callus is produced in response to injury. Callus was

defined as tissue constituted by differentiated cells, which develop in response to a chemical or physical lesion, under determinate hormonal conditions. It can be obtained from a tissue fragment and only some of the callus cells exhibit the totipotency, *i. e.* the ability to differentiate into tissues, organs and even embryos, being able to regenerate whole plants (Pierik, 1990). The callus cultures will be used to study protoplast isolation, cell type, cellular selection, somatic embryogenesis, organogenesis and secondary metabolite production. Oil body formation in *Euphorbia tirucalli* cell suspension culture was noted by Ohyama *et al.* (1984b) and protoplast isolation from *Euphorbia tirucalli* cell suspension cultures and sustained cell division was observed by Ohyama *et al.*, (1984a). It was Chua, (1966), who studied the tissue culture of *H. brasiliensis* role of osmotic concentration, carbohydrate and pH value in induction of callus growth in plumule tissue from *Hevea* seedlings. Asokan *et al.* (2001), studied the isoenzyme markers for distinguishing embryogenic calli from non embryogenic during somatic embryogenesis in *H. brasiliensis*.

In *Mallotus philippensis* (Abbas, 1993) obtained a continuously growing callus on MS + 2,4-D (5.78 μ M) + Kn (2.5 μ M). These calli when sub cultured on MS + BA (13.3 μ M) + CH (Casein hydrolysate, 100mg/l) gave rise to four types of morphologically distinct cell lines. Among these four lines, only the green compact cell line was responsive for organogenic differentiation. Shoot regeneration occurred in this callus when sub cultured on MS + BA (13.3 μ M) + NAA (1.1 μ M). *Euphorbia esula* is a vigorous, highly competitive perennial weed of Eurasian origin that has spread rapidly over the Northern US and Canada and has become an economic pest in the past several years (Watson, 1985). Lee and Starratt (1972) were the first to establish callus cultures from roots of leafy spurge seedlings, using NAA. Nevertheless, they did not attempt to regenerate plants from their cultures.

Suspension cultures and co-cultures

The cell suspension obtained from the stem callus of *Euphorbia esula* on basal medium containing a reduced oxidized nitrogen ratio of 33:67 under the fluorescent lights. Roots and shoots formed and then acclimatized (Davis *et al.*, 1988). Soussi *et al.*, (1996) studied the interaction of *Pseudomonas fluorescens* isolate LS 102 and *Flavobacterium balustinum* isolate 105 with *Euphorbia esula* callus. They concluded that the callus tissue might provide an excellent working model to investigate the mode and /or mechanism of potential bio-control agents on their host plants. The growth anatomy and morphogenetic potential of callus and cell suspension of *Hevea brasiliensis* was studied by Wilson and Street (1975).

Somatic embryogenesis and embryo culture

Studies utilizing immature zygote embryos help researcher gain greater insight into embryo development and seed maturation. Embryo rescue is a valuable tool for plant breeders to obtain hybrids from crosses that would otherwise abort on the plant. Multiplication of *Croton urucurana* can be achieved by micropropagation, either organogenesis or somatic embryogenesis (Grattapaglia and Machado., 1998). Tyagi and Govil (1999) worked on *Emblica officinalis* to observe somatic embryogenesis and micropropagation.

There have been several reports of cassava plant regeneration via somatic embryogenesis that are of reproducible (Stamp and Henshaw, 1982, 1987a ; Stamp 1987; Szabados *et al.*, 1987; Cabral *et al.*, 1992) and secondary somatic embryogenesis (Stamp and Henshaw, 1987b). In *Manihot esculenta* the cyclic system of somatic embryogenesis was improved by using both liquid and solid media of which liquid medium was suitable and helped in the induction of multiple embryos (30) and they were converted into shoots (Raemakers *et al.*, 1993a,b). Somatic embryo fragments of *Manihot esculenta* cultivar "Nanzhi 188" cultured on the media containing cytokinins like (BA and TDZ) induced the shoot organogenesis. While auxins stimulated somatic embryogenesis. The efficiency depends on the different cytokinins and their combinations with auxins (Ma, 1998). Raemakers *et al.*, (1999) developed a protocol for the direct cyclic somatic embryogenesis of *Manihot esculenta* for mass production. The leaf callus derived cotyledons of primary somatic embryo of Cassava (*Manihot esculenta*), cultured on auxin supplemented MS medium (liquid / solid), resulted in the formation of the secondary somatic embryos which were cultured on the Gresshoff and Doy's basal medium supplemented with auxins results in indirect somatic embryogenesis. Depending on BA concentration, plants can be transferred either directly to green house or after using standard multiplication protocols (Raemakers *et al.*, 2000). Woodward and Kaerlas (2001) studied the floral material embryogenic potential of *Manihot esculenta*.

Groll *et al.*, (2002) studied the effects of quarter, half, full and double strength medium salt concentration on differentiation and maturation of somatic embryos of Cassava (*Manihot esculenta*) finally they suggested that half to full strength MS medium was quite suitable for the induction of proliferative nodular embryogenic callus after the desiccation in the saturated K₂SO₄ for 10 days. The zygotic embryos of *Givotia rottleriformis* were cultured on MS + 3% sucrose. On shifting, the seedling to medium containing tyrosine 100 mg l⁻¹ the

survival rate of acclimatized seedlings was elevated to 60-70% (Rambabu *et al.*, 2006).

Somatic embryos of *Hevea brasiliensis* were grown in the specialized conditions like desiccation, medium osmolarity, ABA concentration in the medium for the conversion into plantlet. Slow desiccation or maturation on 351 mol m⁻³ sucrose supplemented with 1m mol m⁻³ ABA strongly improved germination ability and conversion of embryos into plantlets indicated the increased level of vigor (Etienne *et al.*, 1993). Etienne *et al.*, (1997) developed a protocol for improvement of somatic embryogenesis in *Hevea brasiliensis* using the temporary immersion technique. For the last two decades, considerable progress has been made on *in vitro* techniques for multiplication and improvement of *Hevea*. The effect of thidiazuron on axillary proliferation of *H. brasiliensis* was reported by Seneviratne and Flemingmann, (1996). Jha *et al.*, (2007) reported somatic embryogenesis in *Jatropha curcas* Somatic embryogenesis has been reported from anthers as well as inner integumental tissue (Wang *et al.*, 1980; Nataraja, 1975) noted the morphogenesis in embryonal calli of *Euphorbia pulcherrima*.

Effect of ABA and cytokinins on the somatic embryogenesis of *Hevea brasiliensis* was reported by Veisseire *et al.* (1994b). Effect of conditioned media on the somatic embryogenesis of *H. brasiliensis* was reported by Veisseire *et al.*, (1994a). The influence of growth regulators and sucrose on somatic embryogenesis from immature inflorescences of *H. brasiliensis* was reported by Sushamakumari *et al.*, (2000). In *H. brasiliensis* multiple shoots were induced from the somatic embryos (Sushamakumari *et al.*, 1999). Effect of ABA and high concentration of polyethylene glycol on *H. brasiliensis* somatic embryo development was studied by Linossier *et al.*, (1997) GA₃ regulated embryo induction and germination in *H. brasiliensis* was recorded by Kumarijayasree and Thulaseedharan, (2001). Kumarijayasree *et al.*, (2001a), optimized the parameters effecting somatic embryogenesis and long term somatic embryos were initiated and maintained by Kumarijayasree and Thulaseedharan (2004) and Kumarijayasree *et al.*, (2001b). Jayatilake (2007) studied the micrografting, female floret culture and somatic embryogenesis of *H. brasiliensis*. *In vitro* micro grafting of *H. brasiliensis* was studied by Kala *et al.*, (2002). The water status of callus from *H. brasiliensis* during induction of somatic embryoids was recorded by Etienne *et al.*, (1991). Hadrami *et al.*, (1989) and Carron *et al.*, (1995) studied the effect of polyamines on the somatic embryogenesis. Carron and Enjarlic (1982) and Carron *et al.*, (1985) developed protocols for vegetative propagation of *H. brasiliensis* by somatic embryogenesis and *in vitro* microcutting.

Organogenesis and morphogenesis

The organogenesis in *Euphorbia esula* was studied by Davis and Olson (1993) IAA induced the roots very effectively, high concentrations of IAA, NAA, 2,4-D promoted callusing. High or low concentrations of picloram reduced the efficiency of rooting. Martin *et al.*, (2005) reported the influence of auxins in direct *in vitro* morphogenesis of mesophyll cells of *Euphorbia nivulia*, Kn reduced the rate of morphogenesis, where as BAP induced somatic embryogenesis. The combination of BA with NAA and IAA had positive effect on morphogenesis. IBA (13.3 μ M) induced shooting and half MS + IBA (2.46 μ M) suitable for rooting; BA (4.44 μ M) +2,4-D (2.26 μ M) were optimum for somatic embryogenesis of proximal explant. GA₃ supplementation to half MS medium resulted in the conversion of embryos into plantlets with survival rate of 90%. Paranjothy and Gandimathi, (1975, 1976) reported on callus cultures and morphogenesis and tissue and organ cultures of *Hevea brasiliensis*

Anther culture

The aim here is the production of haploid plants through the induction of androgenesis in the haploid cells of the immature pollen grain. Haploid plants are important for number of reasons. Because they possess single set of chromosomes, plant breeders interested in haploid plants because either spontaneous doubling of chromosome number (to 2n) or an application of the colchicine to double the chromosome number gives rise to homozygous plants (Razdan, 2003). Immature anthers of *Hevea brasiliensis* were cultured on the modified MS medium supplemented with 2,4 -D (2.0 mg/l) and Kn (0.5mg/l) resulted in the induction of the callus. By culturing, the callus on MS + Kn (0.7mg/l) and NAA (0.2mg/l) produced maximum number of somatic embryoids. They were further developed from embryos to plantlets on the hormone free medium and then acclimatized (Jayasree *et al.*, 1999). Effect of altered temperature on plant regeneration frequencies in stamen culture of rubber trees was studied by Wang *et al.*, (1998). Wang and Chen (1995) studied the effects of temperature on stamen culture and somatic plant regeneration frequencies in stamen culture of rubber tree. Shiji *et al.*, (1990) prepared a report on anther cultures of *H. brasiliensis*. Propagation of calli from anthers of *H. brasiliensis* was done by Satchuthananthavale and Irugalbandara, (1972). Chen *et al.*, (1979) prepared a protocol for obtaining pollen plants of *H. brasiliensis*. Chen *et al.*, (1982) discussed the recent advances in anther culture of *H. brasiliensis*.

Endosperm culture

Morphogenesis from endosperm tissues has been reported in *Jatropha panduraefolia* (Srivastava and Johri, 1974). Endosperm callus of *Emblica officinalis* sub cultured on MS + BAP (0.2mg/l) and IAA (0.1mg/l) resulted in the formation of shoots and embryo like structures in 50 and 8% cultures respectively. Regeneration of shoots was more frequent when both BAP (0.2mg/l) and IAA (0.1mg/l) were present than on BAP (0.2mg/l) alone. The embryo like structures produced plantlets (Sehgal and Khurana, 1985). Thomas and Chaturvedi (2008) focused on Endosperm cultures of Euphorbiaceae. The growth and metabolism of *Ricinus communis* endosperm in tissue culture was noted by Brown *et al.*, (1970), Johri and Srivastava (1972). The organogenesis in endosperm tissue culture of *Codiaeum variegatum* was reported by Chikkannaiah and Gayatri (1974) and Gayatri (1978). Sehgal *et al.*, (1981) reported the growth responses of mature endosperm of *Euphorbia geniculata*. Sehgal and Abbas (1996), induced triploid plants from the endosperm cultures of *Mallotus philippensis*. Srivastava (1973) reported the formation of triploid plantlets in endosperm cultures of *Putranjiva roxburghii*.

In vitro Flowering and fruiting

Flower heads had more number of meristematic initials, which facilitates the active proliferation of calli and respective organs. The nodes from the flower heads will give an excellent multiple shoots too. When we know the complete flowering mechanism, we can alter the growth and development patterns, or else we can revert the growth of the plant material. In case of *Euphorbia milli* the reversion of inflorescence development and its applications to large-scale micropropagation in airlift bioreactor was done by Dewir *et al.*, (2005a). Tang *et al.*, (1983) reported *in vitro* flowering in five genotypes of *Manihot esculenta* out of his 13 genotypes. It has given the recent trend towards selecting late flowering and non-branching cassava genotypes, *in vitro* flowering system may be the first step towards the possibility of recombining genetic material via *in vitro* fertilization in otherwise non-hybridizable lines. Liang and Keng (2006) reported the role of affecting factors (light, GA₃, sucrose, number of subcultures) on *in vitro* flowering and fruiting while they are working on the micropropagation of valuable medicinal plant *Phyllanthus niruri*.

Factors affecting organogenesis and plant regeneration

This includes effect of temperature, light, pH, plant growth regulators and orientation of the explant on the medium. They have great effect on the explant response but very few workers concentrated on these aspects. The physical status of the plant and the genotype are significant. Continuous light period was found optimum for endosperm proliferation of *Ricinus communis* (Srivastava 1971a). The optimum temperature for the growth of endosperm reported as 25°C and the pH was 5.0 for *Ricinus communis* (Johri and Srivastava, 1973). For *Jatropha* and *Putranjiva* the optimum pH was 5.6 (Srivastava, 1971b). Ripley and Preece (1986) working on *Euphorbia lathyris* found that minimal wounding and vertical orientation of the explants during inoculation are must for the shoot tip rise for which BA is not required where as for callusing and adventitious shooting BA is must. The etiolated shoot tips were rooted well on MS + NAA (1.0 mg/l⁻¹) and the plantlets were acclimatized. Hadrami and Auzac (1992) studied the effect of growth regulators polyamine content and peroxidase activity in *Hevea brasiliensis* callus. CaCl₂ at a concentration between 0-9 mM was used in the callogenesis medium of *Hevea brasiliensis*, low concentrations induced the embryogeny, at high concentrations (12 mM CaCl₂) induced the friable callus. They observed that the CaCl₂ has effective interactions with growth regulators and can decrease the rate of nitrogen metabolism, reduction in the water content (Montoro *et al.*, 1995). Effect of strictly plant related factors on the response of *H. brasiliensis* was studied by Lardet *et al.*, (1998). As per Matysiak and Nowak (1995), increased concentration of CO₂ also lessens water stress of microcuttings by closing the stomata. Unander *et al.*, (1995) studied the factors effecting the germination and stand establishment of *Phyllanthus amarus*. Sochacki and Chimid, (1994) studied the effect of several factors on rooting of Poinsettias (*Euphorbia pulcherrima*). The effect of light emitting diode on growth and shoot proliferation and effect of paclobutrazole, light emitting diodes and sucrose on flowering of *Euphorbia milli* was studied by Dewir *et al.* (2005b, 2006).

A major focus of research during the past two decades has been the manipulation of growth media and growth conditions. Another is the testing of a variety of explant sources to obtain somatic embryogenesis, organogenesis and regeneration of plants. If we want to reduce the micropropagated seedling production cost and automation of the work we can use robots (Kozai *et al.*, 1988; Kozai, 1991a, b). Rapid multiplication, germplasm conservation, pathogen elimination, genetic manipulations and for secondary metabolite production bioreactor technology may

reduce the production cost provided proper precautions are taken to prevent contamination. Teixeira da Silva, (2003) reviewed on the usage of thin cell layer technology in ornamental plant micropropagation and biotechnology, which highlights organogenesis and somatic embryogenesis for plant regeneration and genetic transformation. Somatic embryogenesis facilitated cryopreservation, synthetic seed development, mutations and genetic transformation. Plant transformation methods and gene silencing technology can effectively used to evaluate and authenticate newly discovered endogenous genes to characterize their function in plants as well as genetically manipulate trait quality and productivity (Dandekar, 2003).

CONCLUSION

The review given here can be used for multiplication of the above said medicinal and economical plants commercially. Farmers, tribal people, pharmaceutical companies can utilize this information and can benefit economically. Active principles isolated from the cultures and from plants can be utilized as leads for further drug development by the pharmaceutical industry. Grafting compatibility studies are very helpful for the pathological studies.

ACKNOWLEDGEMENTS

The authors are grateful to Council of Scientific and Industrial Research (CSIR), New Delhi for financial support and Dr. V. Koteswaramma, Architect of Montessori Educational Complex for her encouragement and moral support.

REFERENCES

- Abbas, N. 1993. Studies on somatic embryogenesis and organogenesis in three economic plants, *Mallotus philippensis*, *Aleurites fordii* - mature endosperm, *Trachyspermum ammi*, seedling explants. Thesis submitted to University of Delhi, India.
- Abel-Fattah, M. and Rozk. 1987. The chemical constituents and economic plants of the Euphorbiaceae. Botanical Journal of the Linnean Society, 94: 293-326.
- Ahn, Y. J., Vang, L., Thomas, A., Mc Keon and Chen, G.Q. 2007. High-frequency plant regeneration through adventitious shoot formation in castor (*Ricinus communis* L.). In *Vitro Cell. Developmental Biology-Plant*, 43: 9-15.

- Anonymous. 1978. The Wealth of India, Council of Scientific and Industrial Research, New Delhi. 3: 224 - 230.
- Anthony, P., Davey, M.R., Power, J.B. and Lowe, K.C. 1995. An improved protocol for the culture of Cassava leaf protoplasts. *Plant Cell Tissue and Organ Culture*, 42: 299-302.
- Arifin, B., Poapongsakorn, N., Fatimah, M., Arshad and Amna A.A. Hameed 2008. Bio-fuel Trends in Asia -1: Indonesia, Malaysia and Thailand- IPC-ICONE Seminar on "Climate Change, Agriculture and Trade" in Bogor, Indonesia. ALARN (Asia-Latin America Agricultural Research Network) - May 12, 2008.
- Asma Nasib., Kashif Ali and Saifullah Khan. 2008. *In vitro* propagation of *Croton* (*Codiaeum variegatum*), *Pakistan Journal of Botany*, 40: 99-104.
- Asokan, M.P., Narasimhan, K., Kumarijayasree, P., Thulasidharan, A., Sushmakumari, S., Kala, R.G. and Jayasree, R. 2001. Isoenzyme markers for distinguishing embryogenic calluses from non embryogenic during somatic embryogenesis in *Hevea brasiliensis* Mull.- Arg. *Journal of Tree Science*, 20: 78-85.
- Asokan, M.P., Sobhana, P., Sushmakumari, S. and Sethuraj, M.R. 1988. Tissue culture propagation of rubber (*Hevea brasiliensis* Mull.- Arg.) Clone GT1. *Indian Journal of Natural Rubber Research*, 1:10-12.
- Balotis, G. and Papafotiou, M. 2003. Micropropagation and stability of *Euphorbia pugniformis* cristate form. *Acta Horticulturae*, (ISHS) 616: 471-474.
- Bhagwat, B., Vieiral, L.G. F. and Erickson, L.R., 1996. Stimulation of *in vitro* shoot proliferation from nodal explants of Cassava by thidiazuron, benzyladenine and gibberellic acid, *Plant Cell Tissue and Organ Culture*, 46:1-7.
- Bhattacharyya, R. and Bhattacharya, S. 2001. High frequency *in vitro* propagation of *Phyllanthus amarus* Schum. & Thonn. by shoot tip culture. *Indian Journal Experimental Biology*, 39:1184-1187.
- Bhojwani, S.S. and Johri, B.M. 1971. Morphogenetic studies on cultured mature endosperm of *Croton bonplandianum*, *New Phytologist*, 70:761-766.
- Biesboer, D.D. and Mahlberg, P.G. 1979. The effect of the medium modification and selected precursors on sterol production by short-term callus cultures of *Euphorbia tirucalli*. *Journal Natural Products*, 42: 648-657.
- Bingtao, Li., Huaxing, Qiu., Jin-shuang Ma., Hua Zhu., Michael, G., Gilbert, Hans-Joachim (Hajo) Esser, Stefan Dressler, Petra Hoffmann, Lynn J. Gillespie, Maria Vorontsova and Gordon D. McPherson 2008. *Flora of China*. Vol. 11 pp. 163, <http://www.efloras.org>, dated 22nd September. 2008.
- Bouychou, J. G. 1953. La culture *in vitro* des tissue d'*Hevea*. *Proceedings Rubber Conference Bogor*, *Archives of Rubber Cultivation*, 30: 50-53.
- Brown, D.J., Canvin, D.T. and Zilkey, B.F. 1970. Growth and metabolism of *Ricinus communis* endosperm in tissue culture. *Canadian Journal Botany*, 48: 2323-2331.
- Cabral, G.B., Aragao, F.J.L., Matsumoto, K., Monte Neshich, D.C., Rech, E.L., 1992. First International Scientific Meeting of the Cassava. *Biotechnology Network (CBN)*, Cartagena Colombia, Abstracts: 48.
- Calixto, J.B., Santos, A.R.S., Filho, C.V. and Yunes, R.A. 1998. A review of the plants of the genus *Phyllanthus*: Their chemistry, pharmacology and therapeutic potential. *Medical Research Review* 4: 225-258.
- Calvin, M. 1980. Hydrocarbons from plants: analytical methods and observations. *Naturewissen*, 67: 525-533.
- Carron, M.P. and Enjalric, F. 1982. Studies on vegetative propagation of *Hevea brasiliensis* by somatic embryogenesis and *in vitro* Microcutting. In : Fujiwara, A (ed.). *Plant tissue culture*, Maruzen, Tokyo. pp. 751-752.
- Carron, M.P. and Enjalric, F. 1983. Prospects for the Micropropagation of *Hevea brasiliensis* *Review Generale des. Caoutch Plant*, 628: 65-68.
- Carron, M.P., Deschamps, A., Enjalric, F. and Lardet, L. 1985. Vegetative *in vitro* propagation by microcuttings of selected rubber tree: a wide

- spread technique before 2000. Proceedings of 1st International Rubber Tissue Culture Workshop, Kuala Lumpur, Malaysia. pp. 22-26.
- Carron, M.P., Etienne, H., Lardet, L., Campagna, S., Persin, Y., Leconte, A. and Chaîne, C., 1995. Somatic embryogenesis in Rubber (*Hevea brasiliensis* Mull.- Arg.) In: Jain. S.M., Gupta. P and Newton. R (ed) Somatic embryogenesis in woody plants, Kluwer Academic Publisher, The Netherlands, 2: 117-136.
- Catapan, E., Luis, M.d.Silva, B., Moreno, F.N. and Viana, A.M. 2002. Micropropagation, callus and root culture of *Phyllanthus urinaria* (Euphorbiaceae). Plant Cell Tissue and Organ Culture, 70: 301-309.
- Catapan, E., Otuki, M.F. and Vianna, A.M. 2001. *In vitro* culture of *Phyllanthus stipulatus* (Euphorbiaceae). Revista Brasileira Botanica, Sao Paulo, 24: 25-34.
- Catapan, E., Fleith, O.M. and Maria, V.A. 2000. *In vitro* culture of *Phyllanthus carolinensis* (Euphorbiaceae). Plant Cell Tissue and Organ Culture, 62: 195-202.
- Chen, C., Chen, F., Chien, C., Wang, C., Chang, S., Hus, H., Ou, S., Ho, Y. and Lu, T.1979. A Process of obtaining pollen plants of *Hevea brasiliensis* Mull.- Arg. Scientia Sinica, 22: 81-90.
- Chen, F., Qian, C., Qin, M., Xu, X. and Xiao, Y.1982. Recent advances in anther culture of *Hevea brasiliensis* (Muell.- Arg.), Theoretical Applied Genetics, 62:103-108.
- Chikkannaiah, P.S. and Gayatri, M.C. 1974. Organogenesis in endosperm tissue culture of *Codiaeum variegatum* Blume. Current Science, 43: 23-24.
- Chishimba, W.K., Lingumbwanga, E., Tembo, L.M. and Hang'andu, A.K. 2000. Effect of cytokinins on *in vitro* propagation of *Uapaca kirkiana*, Journal Tropical Forest Science, 12: 28-36.
- Chitra, P. and Madhusoodanan, V. 2005. Influence of auxins in direct *in vitro* morphogenesis of *Euphorbia nivulia*, a lectinaceous medicinal plant. *In Vitro Cellular Developmental Biology -Plant*, 41: 314-319.
- Chua, S.E.1966. Studies on tissue culture of *Hevea brasiliensis* Mull.- Arg Role of osmotic concentration, carbohydrate and pH value in induction of calluses growth in plumule tissue from *Hevea brasiliensis* seedling. Journal of Rubber Research Institute of Malaya, 19: 272-276.
- Compos, R.A.S., Anez, L.M.M., Dombroski, J.L.D. and Dignart, S.L. 2007. Micropropagation of *Jatropha elliptica* (Pohl) Mull.- Arg. Revista Brasileira Plantas Medicinai, 9: 30-36.
- Dandekar, A.M., 2003. Techniques for manipulating quality and productivity traits in horticultural crops, Acta Horticulturae, 625: 293-305.
- Datta, M.M., Mukherjee, P., Ghosh, B. and Jha, T.B., 2007. *In vitro* clonal propagation of biodiesel plant (*Jatropha curcas* L.), Current Science, 93:1438-1442.
- Davis D. G., Olson P. A. and Stolzenberg, L.R. 1988. Organogenesis in cell culture of leafy spurge (*Euphorbia esula* L.) accessions from Europe and North America, Plant Cell Reports, 7:253-256.
- Davis D. G. and Olson P. A. 1993. Organogenesis in Leafy spurge (*Euphorbia esula* L.) *In Vitro Cellular Developmental Biology-Plant*, 29: 97-101.
- Depeyre, D., Isambert, A. and Sow, D. 1994. Des plantes a latex, source de methane. Biofuture juillet-Aout : 25-28 - 30.
- Deshmukh, S.D and Borle, M.N. 1975. Studies on the insecticidal properties of indigenous plant products. Indian Journal Entomology, 37:11-18.
- Dewir, Y.H., D. Chakrabarty, E.J. Hahn and Paek. K.Y. 2005a. Reversion of inflorescence development in *Euphorbia millii* and its application to large-scale micropropagation in an air-lift bioreactor. Journal of Horticultural Science & Biotechnology, 80: 581-587.
- Dewir, Y. H., D. Chakrabarty, S. J. Kim, E. J. Hahn and K.Y. Paek. 2005b Effect of light emitting diode on growth and shoot proliferation of *Euphorbia millii* and *Spathiphyllum cannifolium*. Horticulture, Environment, and Biotechnology, 46: 375-379.

- Dewir Y. H., Chakrabarty, D., Hahn, E.J. and Paek, J.Y. 2006. The effects of paclobutrazole, light emitting diodes and sucrose on flowering of *Euphorbia millii* plantlets *in vitro*. European Journal of Horticultural Science, 71: 240-244.
- Escobar, R.H., Mafia, G., Roca, W.M. 1997. A methodology for recovering cassava plants from shoot tips maintained in liquid nitrogen. Plant Cell Reports, 16: 474-478.
- Etienne, H., Berger, A. and Carron, M.P. 1991. Water status of callus from *Hevea brasiliensis* during induction of somatic embryogenesis, Physiological Plantarum, 82:213-218.
- Etienne, H., Lartaud, M., Ferriere, N.M., Carron, M.P., Berthouly, M. and Teisson, C. 1997. Improvement of somatic embryogenesis in *Hevea brasiliensis* Müll.-Arg. using the temporary immersion technique. *In Vitro Cellular Developmental Biology-Plant*, 33: 81-87.
- Etienne, H., Montaoro, P., Ferriere, M. and Carron, M.P. 1993. Effect of desiccation, Medium osmolarity and abscisic acid on the maturation of *Hevea brasiliensis* Somatic embryos, Journal of Experimental Botany, 44: 1613-1619.
- Ferriera, M.F., Novais, J.M. and Pais, M.S.S. 1992. Hormonal Control of triterpinols synthesis in *Euphorbia characias* calli. Elsevier sciences, Oxford, Royaume-UNI, 39:31-37.
- Ferriere, N.M., Grout, H. and Carron, M.P. 1992. Origin and ontogenesis of somatic embryos in *Hevea brasiliensis* (Euphorbiaceae). American Journal of Botany, 79:174-180.
- Filho, V.C., Santos, A.R.S., Campos, R.O.P., Miguel, O.G., Yunes, R.A., Ferrari, F., Messana, J. and Calixto, J.B. 1996. Chemical and pharmacological studies of *Phyllanthus carolinensis* in mice. Journal of Pharmaceutical Pharmacology, 48:1231-1236.
- Ganesan, M. and Jayabalan, N. 2006. Role and Importance of chitinase gene expression in plants for fungal disease tolerance-In: Jayabalan, N. (ed). Plant Biotechnology. pp 1-28. APH Publishing corporation, New Delhi.
- Gayatri, M.C. 1978. *In vitro* studies on *Codiaeum variegatum*: Growth and organogenesis in Endosperm tissue. Phytomorphology, 28:395-400.
- Ghanti, K.S., Govindaraju, B., Venugopal, R.B., Rao, S.R., Kaviraj, C.P., Jabeen, F.T.Z., Barad, A. and Rao, S., 2004. High frequency shoot regeneration from *Phyllanthus amarus* Schum. & Thonn. Indian Journal Biotechnology, 3:103-107.
- Grattapaglia, D. and Machado, M. A. 1998. Micropropagação. In: Torres, A. C, Caldas L S, Buso J A. Cultura de tecidos e transformação genética de plantas. Brasília, DF: Embrapa SPI/ Embrapa CNPH, 1:183-260.
- Groll, J., Mycock, D.J. and Gray, V.M. 2002. Effect of medium salt concentration on differentiation and maturation of somatic embryos of Cassava (*Manihot esculenta* Crantz). Annals Botany, 89: 645-648.
- Gunatilleke, I.D. and Samaranyake, C. 1988. Shoot tip as a methods of micropropagation of *Hevea brasiliensis*. Journal of Rubber Research Institute of Sri Lanka, 68:33-44.
- Hadrami, I.E. and Auzac, J.D. 1992. Effect of growth regulators on polyamine content and peroxidase activity in *Hevea brasiliensis* callus. Annals Botany, 69:323-325.
- Hadrami, I.E., Ferriere, M.N., Carron, M.P. and Auzac, J.D., 1989. Polyamines, a possible limiting factor in somatic embryogenesis of *Hevea brasiliensis*. Comptes Rendus de Academic des Science de Paris 308. SeriesIII:205-211.
- Heftmann, E., 1975. Steroid hormones in plants. Lloydia, 38:195-209.
- Heller, J., 1996. Physic nut- *Jatropha curcas* L. Promoting the conservation and use of underutilized and neglected crops.No1. International Plant Genetic Resource Institute, Rome.
- Ishimaru, K., Yoshimatsu, K., Yamakawa, T., Kamada, H. and Shimomura, K. 1992. Phenolic constituents in tissue cultures of *Phyllanthus niruri*. Phytochemistry, 34:2015-2016.
- Itokawa, H., Ichihara, Y., Watanabe, K. and Takeya, K.1989. An antitumor principles from

- Euphorbia lathyris*. *Planta Medica*, 55:271-272.
- Jakobek, J.L., Backhaus, R.A. and Herman, K. 1986. Micropropagation of Candelilla, *Euphorbia antisiphilitica* Zucc. *Plant Cell Tissue Organ Culture*, 7:145-148.
- Jamal, A.H.S., Mark, F.A. and Stephen, M.D. 1993. Effect of 2,4-D on Leafy spurge (*Euphorbia esula*) viable seed production. *Weed Technology*, 7:76-78.
- Jasrai, T., Thaker, K.N. and D'Souza, M.C. 2003. *In vitro* propagation of *Euphorbia pulcherrima* Willd. through somatic embryogenesis. *Plant Tissue Culture*, 13: 31-36.
- Jayasingh, M. 2004. The use of Bio diesel by Indian railways. In: Hedge, N.G., J.N. Daniel and S. Dhar (eds), *Jatropha* and other perennial oil seed crops. BAIF Development Research Foundation, Pune, India, 31-33.
- Jayasree, P.K., Asokan, M.P., Sobha, S., Ammal, L. S., Rekha, K., Kala, R.G., Jayasree, R. and Thulaseedharan, A. 1999. Somatic embryogenesis and plant regeneration from immature anthers of *Hevea brasiliensis* Muell.- Arg. *Current Science*, 76:1242-1245.
- Jayatillake, K.M.D.P. 2007. Studies on micrografting of *Hevea brasiliensis* and female florets culture of *Hevea brasiliensis* for somatic embryogenesis. Research Report of Bachelor of Science in Agriculture, University of Ruhuna, Kamburupitiya, Sri Lanka. pp. 22-36.
- Jha, M.M., Mukherjee, P., Datta, M.M. 2007. Somatic embryogenesis in *Jatropha curcas* L. an important biofuel plant. *Plant Biotechnology Reports*, 1: 135-140.
- Johnson, M. and Manickam, V.S. 2003. *In vitro* micropropagation of *Baliospermum montanum* (Willd.)- Muell. Arg. A medicinal plant. *Indian Journal Experimental Biology*, 41:1349-1351.
- Johri, B.M. and Srivastava, P.S. 1972. *In vitro* growth response of mature endosperm of *Ricinus communis* L. In: Murty, Y. S., B.M. Johri, H.Y. Mohan Ram, T.M. Verghes (eds) *Advances in plant morphology* (Prof. V. Puri comm. volume.). Sarita prakashan, Meerut (India), pp 339-358.
- Johri, B.M. and Srivastava, P.S. 1973. Morphogenesis in endosperm cultures. *Z Pflanzenphysiologie*, 70:285-304.
- Kaewsuwan, S., Soonthornchareonnon, N. and Prathanturug, S. 2005. *In vitro* propagation of *Mallotus repandus* (Willd.) Muell. Arg. *Acta Horticulturae*. (ISHS) 676:95-101.
- Kala, R.G., Asokan, M.P., Kumarijayasree, P., Sobha, S., Jayasree, R., Rekha, K. and Tulaseedharan, A. 2002. *In vitro* micrografting in *Hevea brasiliensis*: Optimization of conditions. *Indian Journal of Natural Rubber Research*, 15:165-171.
- Kamalendra, S. and Sudarshanam, S. 2003. *In vitro* micropropagation of *Baliospermum axillare* Blume. *Indian Journal of Plant Physiology*, 8:125 -128.
- Kartha, K.K., Gamborg, O.L., Constabel, F. and Shyluk, J.P. 1974. Regeneration of Cassava plants from apical meristems. *Plant Science Letters*, 2: 107-113.
- Khanna, P. and Nag, T.N. 1973. Isolation, identification and screening of phyllembelin from *Embllica officinalis* Gaertn. tissue culture. *Indian Journal of Pharmacology*, 35:23-25.
- Kleiman, R., Smith, C.R. and Yates, S.G. 1964. Search for new industrial oils. XII. Fifty eight Euphorbiaceae oils, including one rich in vernolic acid. *Journal of the American Oil Chemists Society*, 42: 169-171.
- Konan, N.K., Schöpke, C., Carcamo, R., Beachy, R.N. and Fauquet, C. 1997. An efficient mass propagation system for Cassava (*Manihot esculenta* Crantz) based on nodal explants and axillary bud derived meristems. *Plant Cell Reports*, 16: 444 - 449.
- Konan, N.K., Sangwan, R.S. and Sangwan, B.S. 1994. Somatic embryogenesis from cultured mature cotyledons of Cassava (*Manihot esculenta* Crantz). *Plant Cell Tissue Organ Culture*, 37:91-102.
- Kozai, T. 1991a. Autotropic micropropagation. In: Bajaj Y.P.S. (ed). *Biotechnology in agriculture and forestry. High tech and micropropagation 1*. Springer-Verlag, New York. Pp: 313-343.

- Kozai, T. 1991b. Controlled environments in conventional and automated micropropagation. In: Levin R, Vasil IK, editors. Cell culture and somatic cell genetics of plants. Scale-up and automation in plant tissue culture. London: Acad. Press Inc, pp. 213-230.
- Kozai, T., Kubota, C. and Watanabe, I. 1988. Effect of basal medium composition on the growth of carnation plantlets in auto and mixotropic tissue culture. *Acta Horticulturae*, 230: 159-166.
- Kumarijayasree. and Thulaseedharan, A. 2001. Gibberellic acid regulated embryo induction and germination in *Hevea brasiliensis*. *Indian Journal of Natural Rubber Research*, 14: 106 - 111.
- Kumarijayasree. and Thulaseedharan, A. 2004. Initiation and Maintenance of long-term somatic embryos in *Hevea brasiliensis*. IRRDB Biotechnology work shop, 9-11 Feb 2004, Kuala Lumpur, Malaysia. pp.56.
- Kumarijayasree, P., Sunitha, S.S. and Thulaseedharan, A. 2001a. Effect of cytokinin on *in vitro* germination of *Hevea* somatic embryos. In Proceedings of the IRRDB Meeting, Montpellier, France, pp. 106-109.
- Kumarijayasree, P., Thomas, V., Saraswathyamma, C.K. and Thulaseedharan, A. 2001b. Optimization of parameters affecting somatic embryogenesis in *Hevea brasiliensis*. *Indian Journal of Natural Rubber Research*, 14:20-29.
- Kumari, K.G., Ganesan, M. and Jayabalan, N. 2008. Somatic organogenesis and plant regeneration in *Ricinus communis*. *Biologia Plantarum*, 52: 17-25.
- Kupchan, S.M., Uchida, I., Branfman, A.R., Dailey, R.C. and Fei, B.Y. 1976. Antileukemic principles isolated from Euphorbiaceae. *Plant Science*, 191: 571-572.
- Lang he, E.D., Debergh, P. and Rijk, V.R. 1974. *In vitro* culture as a method for vegetative propagation of *Euphorbia pulcherrima*. *Zeitschrift fur Pflanzenphysiologie* 71:271-274.
- Lardet, L., Aguilar, M.E., Michaux-Ferrere, N. and Berthouly, M. 1998. Effect of strictly plant related factors on the response of *Hevea brasiliensis* and *Theobroma cacao* nodal explants cultured *in vitro*. *In Vitro Cellular Developmental Biology-Plant*, 34: 34 - 40.
- Lee, C.W., Jeckes, J. and Thomas, J.C. 1982. Culture Propagation of *Euphorbia lathyris*, *Horticultural Science*, 17:533 -553.
- Lee, T.T. and Starratt, A.N. 1972. Growth substance requirements and major lipid constituents of tissue cultures of *Euphorbia esula* and *E. cyparissias*, *Canadian Journal Botany*, 50:723-726.
- Liang, O.P. and Keng, C.L. 2006. *In vitro* plant regeneration, flowering and fruiting of *Phyllanthus niruri* L (Euphorbiaceae). *International Journal of Botany*, 2:409-414.
- Lima, E.C., Paiva, R., Nogueira, R.C., Soares, F.P., Emrich, E.B. and Silva, A.A.N. 2008. Callus induction in leaf segments of *Croton urucurana* Baill. *Ciencia Agrotecnologia*, 32:17-22.
- Linossier, L., Vessiere, P., Cailloux, F. and Coudret. 1997. Effect of Absciscic acid and high concentrations of PEG on *Hevea brasiliensis* somatic embryos development. *Plant Science*, 125:183-191.
- Ma, G., 1998. Effects of cytokinins and auxins on Cassava shoot organogenesis and Somatic embryogenesis from somatic embryo explants. *Plant Cell Tissue Organ Culture*, 54: 1-7.
- Ma, G. and Xu, Q. 2002. Induction of somatic embryogenesis and adventitious shoots from immature leaves of Cassava. *Plant Cell Tissue Organ Culture*, 70: 281-288.
- Maciel, A.M., Pinto, A.C., Brabo, S.N. and Silva, M.N. 1998. Terpenoids from *Croton cajura*. *Phytochemistry*, 49: 823-826.
- Makkar, H.P.S., Aderibigbe, A.O. and Bekker, K. 1998. Comparative evaluation of non toxic and toxic varieties of *Jatropha curcas* for chemical composition, digestibility, protein degradability and toxic factors, *Food Chemistry*, 62: 207-215.
- Maliro, M. 1997. Propagation of *Uapaca kirkiana* using tissue culture techniques. M.Sc., Thesis, Bunda College of Agriculture, Lilongwe, Malawi, pp 98.

- Marconi, P.L. and Radice, S.1997. Organogenesis and somatic embryogenesis in *Codiaeum variegatum* (L.) Blume cv. Corazon de Oro. *In Vitro Cellular Developmental Biology-Plant*, 33: 258-262.
- Maridass, M. and Thangavel, K. 2008. Conservation efforts through *in vitro* seed germination of a hepatoprotective plant *Phyllanthus beddomei* (Gamble) Mohanan. *Ethnobotanical Leaflets*, 12: 419-424.
- Martin, A.P., Salgueiro, L.R., Conclaves, M.J., Vila, R., Tomi, F., Adzet, T., Cunha, A.P., Canigüeral, S. and Casanova, J. 2002. Antimicrobial activity and chemical composition of bark oil of *Croton stellulifer*. *Planta Medica*, 66:647-652.
- Martin, K.P., Sundarakumari, C., Chitra, M. and Madhusoodhanan, P.V. 2005. Influence of auxins in direct *in vitro* morphogenesis of *Euphorbia nivulia*, a lectinaceous medicinal plant, *In Vitro Cellular Developmental Biology-Plant*, 41: 314-319.
- Mathews, H., Scopke, C., Carcamo, R., Chavarriaga, P., Fauquet, C. and Beachy, R.N. 1993. Improvement of somatic embryogenesis and plant recovery in Cassava. *Plant Cell Reports*, 12: 328-333.
- Matysiak, B. and Nowak, J. 1995. Acclimatization of *ex vitro* Homolemena Emerald Gem' as effected by nutrient solution concentration and CO₂ enrichment. *Acta Horticulturae*. 390:157-160.
- Mng'omba, S.A. 2007. Development of Clonal propagation protocols for *Uapaca kirkiana* and *Pappea capensis*, two South African trees with economical potential. Ph.D., Thesis, University of Pretoria, Pretoria, South Africa, pp.195.
- Mng'omba, S.A., DuToit, E.S., Akinnifesi, F.K. and Venter, H.M. 2007. Histological evaluation of early graft compatibility of *Uapaca kirkiana* Muell.-Arg. scion / stock combinations. *Hortic Sci.*, 42: 1-5.
- Molina, S.M. and Schobert, C.1995. Micropropagation of *Ricinus communis*. *Journal of Plant Pathology*, 147: 270-272.
- Montoro, P., Etienne, H. and Carron, M.P. 1995. Effect of calcium on callus friability and Somatic embryogenesis in *Hevea brasiliensis* Mull.-Arg.: relations with callus mineral nutrition, nitrogen metabolism and water parameters. *Journal of Experimental Botany*, 46: 255-261.
- Murashige, T. and Skoog, F.1962. A revised medium for rapid growth and bioassays with tobacco tissue culture. *Physiol. Plant*, 15: 473-497.
- Muzik, T.J. and Cruzado, H.J. 1958. Transmission of juvenile rooting ability from seedlings to adults of *Hevea brasiliensis*. *Nature* 181:1288 - 1290
- Nataraja, K. 1975. Morphogenesis in embryonal callus of *Euphorbia pulcherrima in vitro*. *Current Science*, 44: 136-137.
- Nataraja, K., Chennaveeraiah, M.S. and Girigowda, P. 1973. *In vitro* production of shoot buds in *Euphorbia pulcherrima*. *Current Science*, 42:577-578.
- Nielsen, M D., Farestveit, B. and Andersen, A.S. 2003. Adventitious shoot development from decapitated plants of periclinal chimeric poisetia plants (*Euphorbia pulcherrima* Willd. ex Klotsch). *European Journal of Horticultural Science* 68:161-168.
- Nielsen, P.E., Nishimura, H., Otvos, J.W. and Calvin, M. 1977. Plant Crops as a source of fuel and hydrocarbon like materials. *Science*, 198: 942 - 944.
- Ogata, T., Higuchi, H., Mochida, S., Matsumoto, H., Kato, A., Endo, T., Kaji, A. and Kaji, H.1992. HIV-1 reverse transcriptase inhibitor from *Phyllanthus niruri*. *AIDS Research Human Retro viruses*, 8:1937-1944.
- Ohyama, K., Misawa, N., Yamano, Y. and Komano, T.1984a. Protoplast isolation from *Euphorbia tirucalli* L. cell suspension cultures and sustained cell division ZPflanzenphysiology, 113:367-370.
- Ohyama, K., Uchida, Y., Misawa, N., Komano, T., Fujita, M. and Ueno, T. 1984b. Oil body formation in *Euphorbia tirucalli* L. cell suspension cultures. *Plant Cell Rep.*, 3:21-22.
- Openshaw, K., 2000. A review of *Jatropha curcas*: an oil plant of unfulfilled promise. *Biomass and Bioenergy*.19:1-15.
- Orlikowska, T., Sabata, I. and Kucharska, D. 2000. The effect of leaf and shoot tip removal and explant orientation on axillary shoot

- proliferation of *Codiaeum variegatum* Blume var. *pictum* Muell.- Arg. cv. excellent. *Scientia Horticulturae*, 85:103-111.
- Orlikowska, T., Sabata, I. and Nowak, E. 1995. Adventitious shoot regeneration on explants of *Anthurium*, *Codiaeum*, *Dieffenbachia*, *Gerbera* and *Spathiphyllum* for breeding purposes. *Acta Horticulturae*. 420:115-117.
- Paranjothy, K. and Gandimathi, H. 1975. Morphogenesis in callus cultures of *Hevea brasiliensis* Muell.- Arg. Proceedings of Plant tissue culture. Culture symposium, Kaula Lumpur, Malaysia. pp19-25.
- Paranjothy, K. and Gandimathi, H. 1976. Tissue and Organ culture of *Hevea brasiliensis*. Proceedings of International Rubber conference, Kaula Lumpur, Malaysia. pp 59-84.
- Pierik, R.L.M. 1990. Cultivo *in vitro* de las plantas superiores. Sao Paulo: M. Nijoff. Brazil, 329 p.
- Pierik, R.L.M. 1991. Commercial aspects of micropropagation. In *Horticulture-New Technologies and Applications* (eds) Prakash J.K. and R.L.M Pierik), Dordrecht, The Netherlands.
- Prathantharug, S., Soonthornchareonnon, N., Chuakul, W., Phaidee, Y. and Saralamp, P. 2007. An Improved protocol for Micropropagation of *Mallotus repandus* (Willd.) Mull.-Arg. *In Vitro Cellular Developmental Biology-Plant*, 43:275-279.
- Puebla, P., Lopez, J.L., Guerrero, M., Carron, R., Martin, M.L., Roman, L.S. and Feliciano, A.S. 2003. Neo-clerodane diterpenoids from *Croton schiedeana*. *Phytochemistry*, 62:551-554.
- Qin, H., Song, S.Q. and Long, C.L. 2006. Tissue culture and plant regeneration of *Jatropha curcas* (Euphorbiaceae) *Acta Botanica Yunnanica* 28: 649-652.
- Quak, F. 1997. Meristem culture and virus free plants In: J. Reinert and Y.P.S. Bajaj (eds.), *Applied and fundamental aspects of plant cell, tissue and organ culture*. Springer Verlag, pp. 598-646.
- Quraishi, A. and Mishra, S. K. 1998. Micropropagation of nodal explants from adult trees of *Cleistanthus collinus*. *Plant Cell Reports*, 17: 430-433.
- Raemakers, C.J.J.M., Bessembinder, J.J.E., Staritsky, G., Jacobsen, E. and Visser, R.G.F. 1993a. Induction, germination and shoot development of somatic embryos in Cassava. *Plant Cell Tissue Organ Culture*, 33:151-156.
- Raemakers, C.J.J.M., Schavemaker, C.M., Jacobsen, E. and Visser, R.G.E. 1993b. Improvements of cyclic somatic embryogenesis of cassava (*Manihot esculenta* Crantz) *Plant Cell Reports*, 12: 226-229.
- Raemakers, K., Jacobsen, E. and Visser, R. 2000. The use of somatic embryogenesis for plant propagation in Cassava. *Molecular Biotechnology*. Humana press, Totowa, NJ. 14: 215-221.
- Raemakers, K.J.J.M., Jacobsen, E. and Visser, R.G.F. 1999. Direct cyclic somatic embryogenesis of Cassava for production purposes. *Plant cell culture protocols*. 111: 61-70.
- Rajasubramaniam, S. and Saradhi, P.P. 1997. Rapid multiplication of *Phyllanthus fraternus* a plant with anti hepatitis viral activity. *Industrial Crops and Products* 6: 35-40.
- Rajore, S. and Batra, A. 2005. Efficient plant regeneration via shoot tip explant in *Jatropha curcas* L. *Journal Plant Biochemistry*, 14: 73-75.
- Rambabu, M., Ujjwala, D., Ugandhar, T., Praveen, M., Upender, M. and Swamy, N.R. 2005. Effect of GA₃ on enhancement of *in vivo* seed germination in *Givotia rottleriformis* (Euphorbiaceae) an endangered forest tree. *Indian Forestry*, 131: 25-30.
- Rambabu, M., Upender, D., Ujjwala, T., Ugandhar, M., Praveen, and Swamy, N.R. 2006. *In vitro* zygotic embryo culture of an endangered forest tree *Givotia rottleriformis* and factors affecting its germination and seedling growth. *In Vitro Cellular Developmental Biology-Plant*, 42: 418- 421.
- Rani, S.S., Murthy, K.S.R. and Pullaiah, T. 2002. Dye yielding plants of Andhra Pradesh, India. *Journal of Economic and Taxonomic Botany* 26: 739 - 749.

- Rani, S.S., Murthy, K.S.R., Goud, P.S.P. and Pullaiah, T. 2003. Tree wealth in the life and economy of the tribes' people of Andhra Pradesh, India. *Journal of Tropical Forest Science*. 15: 259 - 278.
- Rao, C.S., Eganathan, P., Anand, A., Balakrishna, P. and Reddy, T.P. 1998. Protocol for *in vitro* propagation of *Excoecaria agallocha* L. a medicinally important mangrove species, *Plant Cell Reports*, 17: 861- 865.
- Razdan, M.K. 2003. Introduction to Plant Tissue Culture (2nd ed.) Oxford & IBH Publishing Co. Pvt. Ltd., New Delhi p. 87- 106.
- Ricardo, M.D., Faloci., Mitra, M., Gonzalez., Ana, M., Mroginski. and Luis, A. 2007. *In vitro* cultured primary roots derived from stem segments of Cassava (*Manihot esculenta*) can behave like a storage organs. *Annals Botany*, 99: 409 - 423.
- Ripley, K.P. and Preece, J.E. 1986. Micropropagation of *Euphorbia lathyris* L. *Plant Cell Tissue Organ Culture*, 5: 213-218.
- Rommanee, C., Salak, P., Wichien, Y. and Akira, S. 2003. Routine cryopreservation of *In vitro* grown axillary apices of Cassava (*Manihot esculenta* Crantz) by vitrification: importance of simple mononodal culture. *Scientia Horticulturae*, 98: 485- 492.
- Rout, G.R., Mohapatra A. and Jain, S.M. 2006. Tissue culture of ornamental pot plant: A critical review on present scenario and future prospects, *Biotechnology Advances* 24: 531 - 560.
- Roy, S.K. and Jinnah, M.A. 2001. *In vitro* micropropagation of Poinsettia (*Euphorbia pulchirrima* Willd.). *Plant Tissue Culture*, 11: 133-140.
- Saigo, R.H. and Saigo, B.W. 1983. *Botany Principles and Applications*. Prentice-Hall. Englewood Cliffs, 121-136.
- Santos, A.R.S., Cechinel Filho, V., Viana, A.M., Moreno, F.N., Compos, M.M., Yunes, R.A. and Calixto, J.B. 1994. Analgesic effects of callus culture extracts from selected species of *Phyllanthus* in mice. *Journal of Pharmaceutical Pharmacology*, 46: 755-759.
- Satchuthananthavale, R. and Irugalbandara, Z.E. 1972. Propagation of Callus from *Hevea brasiliensis* anthers. *Quarterly Journal of Rubber Research Institute of Ceylon*, 49: 65-68.
- Sehgal, C.B. and Abbas, N.S. 1996. Induction of Triploid plantlets from the endosperm culture of *Mallotus philippensis* Mull.-Arg. *Phytomorphology*, 46: 283-289.
- Sehgal, C.B., Narang, K.H. and Sunila, A. 1981. Growth responses of mature endosperm of *Euphorbia geniculata* Ortega in cultures. *Beitr. Biol Pflanz.*, 55: 385-392.
- Sehgal, C.B. and Khurana, S. 1985. Morphogenesis and plant regeneration from cultured endosperm of *Embllica officinalis* Gaertn. *Plant Cell Reports*, 4: 263-266.
- Seneviratne, P. and Flemingmann, A.W. 1996. The effect of thidiazuron on axillary proliferation of *Hevea brasiliensis in vitro*. *Journal of Rubber research Institute of Sri Lanka*. 77: 1-14.
- Seneviratne, P. 1991. Micropropagation of Juvenile and Mature *Hevea brasiliensis*. Ph.D., Thesis, University of Bath, UK. pp 48-167.
- Shed, A., Vickery, K., Pajkos, A., Medhurst, R., Freiman, J., Dixon, R. and Cossart, Y. 1992. Effects of *Phyllanthus* plant extracts on duck hepatitis B virus *in vitro* and *in vivo*. *Antiviral Research*, 18: 127-138.
- Shibata, W., Murai, F., Akiyama, T., Siriphol, M., Matsunaga, E. and Morimoto, H. 1996. Micropropagation of *Croton sublyratus* Kuge. a tropical tree of medicinal importance. *Plant Cell Reports*, 16: 147-152.
- Shiji, Z., Zherghua, C. and Xueng, X. 1990. A summary report on anther culture for haploid plants of *Hevea brasiliensis*. In: *Proceedings of IRRDB Symposium on Breeding of Hevea brasiliensis*, Kuming, China, 5-6 October, 67-78.
- Singh, A.K., Sharma, M., Varshney, R., Agarwal, S.S. and Bansal, K.C. 2006. Plant regeneration from alginate encapsulated shoot tips of *Phyllanthus amarus* Schum & Thonn, a medicinally important plant species. *In Vitro Cellular Developmental Biology- Plant*, 42: 109-113.

- Sochacki, D. and Chimid, H.1994. Effect of several factors on rooting of Poinsettias. Kwiaistwaw Skirnewach, 1:74-78.
- Sompong, T.C. and Muangkaewangam, A.1992. Tissue culture of rubber-I. *In vitro* micropropagation of rubber. Journal of Science and Technology, 14:123-132.
- Soussi, T., Kremer, R.J. and White, J.A. 1996. Interaction of rhizobacteria with leafy spurge (*Euphorbia esula* L.) callus tissue cells. Plant Cell Tissue Organ Culture, 47:279-287.
- Srivastava, P.S. and Johri, B. M. 1974. Morphogenesis in mature endosperm cultures of *Jatropha panduraefolia*. Beitr. Biol. Pflanz. 50: 255-268.
- Srivastava, P.S. 1971a. *In vitro* induction of triploid roots and shoots from mature endosperm of *Jatropha panduraefolia*. Zpflanzenphysiology, 66: 93- 96.
- Srivastava, P.S. 1971b. *In vitro* growth requirements of mature endosperm of *Ricinus communis*. Current Science, 40: 337 - 339.
- Srivastava, P.S. 1973. Formation of Triploid plantlets in Endosperm culture of *Putranjiva roxburghii*. Zpflanzenphysiology, 69: 270-273.
- Stamp, J.A. 1987. Somatic embryogenesis in Cassava: the anatomy and morphology of the regeneration process, Annals Botany, 59: 451-459.
- Stamp, J.A. and Henshaw, G.G. 1982. Somatic embryogenesis in Cassava Zpflanzenphysiology 105:183-187.
- Stamp, J.A. and Henshaw, G.G. 1987a. Somatic embryogenesis from clonal leaf tissue of Cassava. Annals Botany, 59:445-450.
- Stamp, J.A. and Henshaw, G.G. 1987b. Secondary somatic embryogenesis and Plant regeneration in cassava. Plant cell Tissue Organ Culture, 10:227-233.
- Stohs, S.J. and Rosenberg, H.1975. Steroid and steroid metabolism in plant tissue culture. Lloydia, 38:181-194.
- Sujatha, M. and Dhingra, M.1993. Rapid plant regeneration from various explants of *Jatropha integerrima*. Plant cell Tissue Organ Culture, 35:293-296.
- Sujatha, M. and Mukta, N.1996. Morphogenesis and plant regeneration from tissue culture of *Jatropha curcas* L. Plant cell Tissue Organ Culture, 44:135-141.
- Sujatha, M., Makkar, H.P.S. and Becker, K. 2005. Shoot bud Proliferation from axillary nodes and leaf sections of non-toxic *Jatropha curcas* L. Plant Growth Regulation, 47: 83-90.
- Sujatha, M. and Reddy, T.P. 1998. Differential cytokinin effects on the stimulation of *in vitro* shoot proliferation from meristematic explants of Castor (*Ricinus communis*) Plant Cell Reports, 17:561-566.
- Sujatha, M. and Sailaja. M. 2005. Stable genetic transformation of Castor (*Ricinus communis* L.) via *Agrobacterium tumefaciens* mediated gene transfer using embryo axes from mature seeds. Plant Cell Reports, 23: 803-810.
- Sushamakumari, S., Rekha, K., Thomas, V., Sobha, S. and Jayasree, R. 1999. Multiple shoot formation from somatic embryos of *Hevea brasiliensis* Muell.- Arg. Indian Journal of Natural Rubber Research, 12:23-28.
- Sushamakumari, S., Sobha, S., Rekha, K., Jayasree, R. and Asokan, M.P. 2000. Influence of growth regulators and sucrose on somatic embryogenesis from immature inflorescence of *Hevea brasiliensis* Muell.- Arg. Indian Journal of Natural Rubber Research, 13:19-29.
- Szabados, L., Hoyos, R. and Roca, W.1987. *In vitro* somatic embryogenesis and plant regeneration of Cassava. Plant Cell Reports, 6:248-251.
- Tang, A.F., Cappadocia, M. and Byrne, D.1983. *In vitro* flowering in cassava (*Manihot esculenta* Crantz). Plant cell Tissue Organ Culture, 2:199-206.
- Teixeira de Silva, J.A. 2003. Thin cell layer technology in ornamental plants micropropagation and biotechnology. African Journal Biotechnology, 2: 683 - 691.
- Thapliyal, M., 2004. Twin seedlings in *Putranjiva roxburghii* Wall. Indian Journal of Forestry, 27: 43 - 44.

- Thomas, T.D. and Chaturvedi, R. 2008. Endosperm culture: a novel method for triploid plant production. *Plant cell Tissue Organ Culture*, 93: 1-14.
- Thyagarajan, S.P., Subrahmanian, S., Thirunalasudary, S., Venkateswaran, P.S. and Blumberg, B.S. 1988. Preliminary study: the effect of *Phyllanthus amarus* on chronic carriers of hepatitis B virus. *Lancet*, 2: 764 - 766.
- Tidemann, J. and Hawker, J.S. 1982. *In vitro* propagation of latex producing plants. *Annals Botany*, 49: 273-279.
- Torres, A.I. 2004. Rooting experiments with *Euphorbia lagascae* cuttings. *Anales de Biologia*, 26:101-104.
- Turley, D., Froment, M. and Cook, S. 2000. Development of *Euphorbia lagascae* as a new industrial oil crop ADAS, Woodthorne, Wolverhampton, UK pp-38.
- Tyagi, T. and Govil, C.M. 1999. Somatic embryogenesis and micropropagation in *Embllica officinalis* Gaertn. *Journal of the Indian Botanical Society*, 78:363-365.
- Uchida, H., Nakayachi, O., Otani, M., Kijikawa, M., Kohzu, Y., Yamato, K. T., Fukuzawa, H., Shimada, T. and Ohyama, K. 2004. Plant regeneration from inter node explants of *Euphorbia tirucalli*. *Plant Biotechnology*, 21: 397-399.
- Ueno, H., Horie, S., Nishi, Y., Shogawa, H., Kawasakim, M., Suzuki, S., Hayashi, T., Arisawa, M., Shimizu, M., Yoshizakim, M., Morita, N., Berganza, L.H., Ferro, E. and Basualdo, I. 1988. Chemical and pharmaceutical studies on medicinal plants in Paraguay. Geraniin, an angiotensin covering enzyme inhibitor from "papaparaime", *Phyllanthus niruri*. *Journal Natural Products*, 51:357-359.
- Unander, D.W. 1991. Callus induction in *Phyllanthus* species and inhibition of viral DNA polymerase and reverse transcriptase by callus extracts. *Plant Cell Reports*, 10: 461-466.
- Unander, D.W., Bryan, H.H., Lance, C.J. and Mc Millan, R.T. 1995. Factors effecting germination and stand establishment of *Phyllanthus amarus* (Euphorbiaceae). *Economic Botany*, 49:49-55.
- Unander, D.W. 1996. *Phyllanthus* species: *in vitro* culture and reverse transcriptase by callus extracts. *Plant Cell Reports*, 10:461-466.
- Veisseire, P., Cailloux, F. and Coudret, A. 1994a. Effect of conditioned media on the somatic embryogenesis of *Hevea brasiliensis*. *Plant Physiological Biochemistry*, 32: 571-576.
- Veisseire, P., Linssier, L. and Coudret, A. 1994b. Effect of abscisic acid and cytokinins on the development of somatic embryos in *Hevea brasiliensis*. *Plant cell Tissue Organ Culture*, 39: 219 - 223.
- Villaluz, Z.A. 2006. Improvement of *in vitro* techniques for rapid meristem development and mass propagation of Philippine Cassava (*Manihot esculenta* Crantz); *Journal of Food Agriculture and Environment*, 4:220-224.
- Wang, Z., Zeng, X., Chen, C., Wu, H., Li, G., Fan, Q. and Lu, W. 1980. Induction of Rubber plantlets from anther of *Hevea brasiliensis* Muell.Arg. *in vitro*. *Chinese Journal of Tropical Crops*. 1: 25 - 26.
- Wang, Z.Y. and Chen, X.T. 1995. Effect of temperature on stamen culture and somatic plant regeneration frequencies in stamen cultures of rubber tree. *Journal of Tropical and Subtropical Botany*, 6:166-168.
- Wang, Z.Y., Wu, H.D. and Chen, X.T. 1998. Effect of altered temperature on plant regeneration frequencies in stamen culture of rubber trees. *Journal of Tropical and Subtropical Botany*, 6: 166-168.
- Watson, A.K., 1985. Leafy spurge monograph No.3, Weed Science Society America. Champaign, IL. USA.
- Wilson, H.M. and Street, H.E. 1975. The growth, anatomy and Morphogenetic potential of calluses and cell suspension cultures of *Hevea brasiliensis*, *Annals Botany*, 39: 671- 682.
- Woodward, B. and Kaerlas, J.P. 2001. Somatic embryogenesis from floral tissue of cassava (*Manihot esculenta* Crantz). *Euphytica* 120: 1- 6.
- Wu, T., Lin, Y., Haruna, M., Pan, D., Shingu, T., Chen, Y., Hsu, H., Nakano, T. and Lee, K. 1991. Anti tumor agents, 119. Kansuiphorins A and B. two antileukemic diterpene esters

- from *Euphorbia kansui*. Journal Natural Products, 54: 823-829.
- Yamamoto, Y., Mizuguchi, R. and Yamada, Y. 1981. Chemical constituents of cultured cells of *Euphorbia tirucalli* and *Euphorbia milli*. Plant Cell Reports, 1: 29-30.
- Yamuna, P., Das, S., Jha, T.B. and Jha, S. 1995. Regeneration and multiplication of shoots in *Glochidion multiloculare* Muell-Arg., Journal of Herbs, Spices and Medicinal Plants, 3: 67-74.
- Zhang, B. and Stoltz, L.P. 1989. Acclimatization Systems for *Euphorbia fulgens* micro cuttings. Horticultural Science, 24: 1025-1026.
- Zhang, B, Stoltz, L.P. and Snyder, J.C. 1987. *In vitro* propagation of *Euphorbia fulgens*, Horticultural Science, 22: 486-488.

Submitted January 04, 2009 – Accepted April 03, 2009
Revised received April 19, 2009