



Boletín de la Sociedad Geológica
Mexicana

ISSN: 1405-3322

sgm_editorial@geociencias.unam.mx

Sociedad Geológica Mexicana, A.C.
México

R. Tévez, Hector; dos Santos Afonso, Maria
pH dependence of Glyphosate adsorption on soil horizons
Boletín de la Sociedad Geológica Mexicana, vol. 67, núm. 3, 2015, pp. 509-516
Sociedad Geológica Mexicana, A.C.
Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=94348267016>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative



pH dependence of Glyphosate adsorption on soil horizons

Hector R. Tévez¹, Maria dos Santos Afonso^{2,*}

¹ Facultad de Ciencias Forestales, Universidad Nacional de Santiago del Estero. Avenida Belgrano Sur 1912. 4200 Santiago del Estero, Argentina.

² INQUIMAE, Facultad de Ciencias Exactas y Naturales- Universidad de Buenos Aires. Ciudad Universitaria, Pabellón II, 3er Piso, Ciudad Autónoma de Buenos Aires, C1428EHA, Argentina.

* dosantos@qi.fcen.uba.ar

Abstract

Pesticides bring many problems to the environment and to human health. The first rationale for their use is increased food production. Glyphosate N-(phosphonomethyl)glycine (PMG) is a non-selective, post emergent, and broad spectrum herbicide, very well known for its extensive application in agriculture worldwide. PMG adsorption experiments were carried out in three horizons of a Typic Haplustoll soil from the Province of Santiago del Estero, Argentina.

Adsorption isotherms were fitted using Freundlich and Langmuir models. The affinity constants (K_F and K_L), the adsorption intensity ($1/n$) and the maximum surface coverage (Γ_{max}) were obtained. The results show the dependence of the parameters K_L and Γ_{max} with pH and also with the different horizons and particle size.

Keywords: Glyphosate, Adsorption isotherm, horizon, profile.

Resumen

Los pesticidas producen muchos problemas al ambiente y a la salud humana. La primera racionalización que considera el aumento en su aplicación es incrementar la producción de alimentos. El glifosato, N-fosfonometilglicina (PMG) es un herbicida no selectivo, post-emergente y de amplio espectro que es de uso extensivo en agricultura a nivel mundial. La adsorción de PMG fue llevada a cabo en tres horizontes de un suelo de la Provincia de Santiago del Estero, Argentina, clasificado como Haplustol Típico.

Las isotermas de adsorción fueron ajustadas utilizando los modelos de Freundlich y de Langmuir. Se determinaron las constantes de afinidad (K_F y K_L), la intensidad de adsorción ($1/n$) y el recubrimiento superficial máximo (Γ_{max}). Los resultados muestran la dependencia con el pH de los parámetros de K_L y Γ_{max} así como con la identidad de los horizontes y el tamaño de partícula.

Palabras clave: Glifosato, isoterma de adsorción, horizonte, perfil.

1. Introduction

In the last century the Province of Santiago del Estero, Argentina, lost more than 80 % their natural forest as a result of irrational logging. In the arid soil of the province, agriculture replaced forests. Soybean cultivation was developed at the expense of others crops, native forests and

livestock (Pérez-Carrera *et al.*, 2008).

Pesticides are chemicals widely used in agriculture and their use increase with the increasing of crop areas. In particular, glyphosate (N-phosphonomethylglycine, PMG) is used to remove annual grasses and perennial broadleaf weeds and woody species in agricultural, forestry and landscape. Thus, PMG is a non selective, post-emergent

and broad-spectrum commercial herbicide used worldwide in soybean agriculture. In the soil, the main way of PMG degradation is microbial mediated, considering that abiotic mineralization of PMG for the horizon A exceeds 1 % reaching a maximum of 12 % degradation for 60 days exposure (Jacobsen *et al.*, 2008).

PMG exhibits fast vertical mobility in soil, reaching high concentrations in deeper horizons where degradation is slower (Veiga *et al.*, 2001).

PMG is a good chelating agent and can coordinate metal ion in aqueous solution, especially at near-neutral pH levels where carboxylate and phosphonate chemicals groups are deprotonated forming strong complexes (Barja *et al.*, 2001). It can also be retained in the soil through adsorption onto aluminum and iron oxides (Nowack and Stone, 1999; Barja and dos-Santos-Afonso, 2005), clays (Damonte *et al.*, 2007; Khoury *et al.*, 2010 and references cited therein) and organic matter (Sposito, 1984; Piccolo *et al.*, 1996).

The PMG adsorption on soils or clays was studied by several authors (Nomura and Hilton, 1977; McConnell and Hossner, 1985; Morillo *et al.*, 2000; Sheals *et al.*, 2002; Pessagno *et al.*, 2008). In all the cases, adsorption decreased with a pH increase following an anionic adsorption behavior.

The study of competitive adsorption between PMG and phosphate on iron oxides had shown that PMG could be exchanged by phosphate (McBride and Kung, 1989; Gimsing and Borggaard, 2001), but the exchange on clays or soils is not so easy to characterize (Dion *et al.*, 2001). When the initial phosphate level is high the glyphosate sorption decreases (Dion *et al.*, 2001).

Thus, the surface coverage and the affinity constants for adsorption on iron oxides are higher than on clays minerals or soils, but the experimental results suggest that phosphonate is the liable chemical group for the surface coordination through inner-sphere surface complexes formation (dos-Santos-Afonso *et al.*, 2004; Pessagno *et al.*, 2005, 2008; Tévez *et al.*, 2008). Previous studies on the surface coverage and the adsorption isotherms of this herbicide on soil fractions from different provinces of Argentina (Santa Cruz, Misiones, and Corrientes) followed similar patterns to those of the pure minerals that form these soils (Pessagno *et al.*, 2005).

The aim of this work is to study the adsorption of PMG onto three different horizons of soils from Santiago del Estero Province, Argentina, with different mineral composition to understand how the mineral composition and pH are controlling the environmental fate of glyphosate.

2. Materials and methods

2.1. Chemicals

All chemicals utilized were of analytical reagent grade and were used without further purification. All solutions and soil dispersions were prepared using Milli-Q water.

PMG solutions were fresh and prepared daily by dissolving the herbicide in Milli-Q water. All PMG solution concentrations ranged from 0.05 to 10 mM.

2.2. Study area

The study area is located near Quimili in the center-east of Santiago del Estero Province between 62° 06' W and 61° 52' W and, 27° 24' S and 28° 00' S (Figure 1). Climate is semiarid mesothermal, with an average annual temperature of 19.6 °C and rainfall of between 600 and 750 mm per year concentrated in the spring-summer period (Torres-Bruchman, 1981).

The sampling area is a soil catena corresponding to that found in recent agriculture (9 years) and low for ancient agriculture (25 – 27 years).

The soil is derived from loessic sediments and it is located at a depression relief, and classified as Typic Pachic Haplustoll with grasslands, *Elionurus muticus* (Lorenz *et al.*, 2000).

Samples were taken up to 130 cm of depth (Figure 2.A) from three very well differentiated horizons classified as Ap (0 – 18 cm), AB (18 – 50 cm) and BC (105 – 130 cm). The Ap is the uppermost mineral horizon, disturbed by plowing or other agricultural practices. AB and BC are transitional horizons, wherein the horizon properties are dominated by horizon A properties but also have characteristics of B horizon (AB) or by horizon B properties but also have characteristics of C horizon (BC). The soil at depths between 50 and 105 cm has intermediate characteristics varying in a continuous from AB to BC and was not considered in this study. An schematic top view of pit trial is shown in Figure 2.B, each horizon was sampled at three different places indicated as 1, 2 and 3, and afterward they were mixed to get one sample per horizon.

2.3. Characterizations

The fresh soil samples were air-dried and ground to pass a sieve of 2 mm. pH was measured in 0.01 M CaCl₂ solution at 1:2.5 ratio of soil suspension (Hendershot *et al.*, 1993; Schlichting *et al.*, 1995) using a combined glass electrode. Organic matter (OM) content and soils chemical analysis were determined by the dichromate oxidation method (Schlichting *et al.*, 1995). The available phosphorus (P) is the inorganic P, that is extractable at pH 8.5 and was determined following the experimental procedure described in Olsen *et al.*, 1954 and Page *et al.*, 1982. The total surface area (Sw) was measured by H₂O adsorption (Torres-Sánchez and Falasca, 1997). The total iron oxides (Fe_{tot}) and amorphous iron oxides (Fe_{amorph}) were established by dithionite (Holmgren, 1967) and oxalate method (McKeague, 1967), respectively.

Soils samples were mixed with Lithium Metaborate/Lithium Tetraborate (LiBO₂/Li₂B₄O₇) and fused in a furnace. The molten melt was completely dissolved in acidic




Figure 1. Map of South America, Argentina and Santiago del Estero Province. The sampling area near Quimilí City is represented by ●.




Figure 2. A: Pit profile from where samples were taken and B: schematic top view of trial pits and sampling site.

media of 5 % nitric acid. This solution was analyzed for major and selected trace elements by Inductively Coupled Plasma - Atomic Emission Spectroscopy (ICP-AES). The sample composition are reported as oxide percentage.

The mineralogical composition and quantitative analysis of the soils were determined by X-ray Diffraction (XRD) and using the Rietveld method (Rietveld, 1969).

Point of zero net proton charge (PZNPC) or point zero salt effect (PZSE) is the pH where the net adsorption of protons and hydroxyl ions on the surfaces is independent of electrolyte concentration. Titration curves, when surface charge is plotted against pH, frequently showed a common intersection point that match with PZNPC.

PZNPC was determined by potentiometric titration of soils dispersions starting from natural pH values near neutrality and by adding increasing amounts of standardized NaOH or HCl 0.1 M at different initial ionic strength (0.1, 0.01 and 0.0012 M KCl) under N_2 atmosphere. Soils dispersions were prepared using Milli-Q water free of CO_2 , and treated by continuous bubbling of N_2 , for at least one hour before starting the titrations. The extent of adsorbed H^+ and OH^- by soil material was determined by subtracting the quantity of HCl or NaOH required to bring the dispersion and the electrolyte solution without soil to the same pH. The surface charge was calculated as follows:

$$Q = \{ \equiv SOH_2^+ \} - \{ \equiv SO^- \} = \frac{(V_{tit} - V_b) \cdot [tit]}{m} \quad (1)$$

where Q , V_{tit} , V_b , $[tit]$ and m are the surface charge, the volume of titrant used for dispersion titration, the volume of titrant used for electrolyte titration, titrant concentration and the mass of the solid, respectively.

2.4. Adsorption experiment

The adsorption of herbicide by the soils was studied using

batch experiments. Solutions of different concentration of glyphosate in a final volume of 11 mL were added to 0.100 g of soil samples dispersions. Dispersions were kept in constant agitation overnight at constant pH, ionic strength and room temperature to reach equilibrium. pH was adjusted during the experiment using small aliquots of HCl or NaOH solutions (0.10 M) and ionic strength was kept constant (0.1 M) using a KNO_3 solution. The sample was filtered through a 0.45 μm membrane and adsorbed glyphosate was calculated from the difference between the total added ligand and the supernatant concentration (C_e). PMG was evaluated by ion chromatography (Zhu *et al.*, 1999) using a DIONEX DX-100 instrument with a conductivity detector, a sample injection valve, and a 25 μL sample loop. Two plastic anion columns were coupled in series to serve both as pre-column (DIONEX AG-4) and analytical chromatographic column (DIONEX AS-4). The suppressor was regenerated with 50 mM H_2SO_4 with a flow rate of 12.5 $mL \cdot min^{-1}$. A mixture of NaOH/ CO_3^{2-} 4 mM/9 mM was chosen as eluent with a flow rate of 1 $mL \cdot min^{-1}$. The typical experimental error is lower than 5 % for all results.

2.5. pH effect

The pH dependence of the glyphosate uptake by soil horizons was investigated using batch isotherm experiments in a pH range from 2 to 8 with a soil concentration of 9.1 $g \cdot L^{-1}$ and different initial concentrations of PMG at a constant ionic strength of 0.1 M of KNO_3 . The pH was measured using a Metrohm 644 pH-meter with a combined glass microelectrode, and the pH was adjusted throughout the experiments using 0.1 M HCl or 0.1 M NaOH. Adsorption experiments were conducted in triplicate following the procedure described above. There were no significant differences within each replicate ($p < 0.01$). The expressed values represent the average of the obtained results.

2.6. Isotherms Modeling

The relationship between the ligand uptake and the sorbate equilibrium concentration at constant temperature is known as the adsorption isotherm. The adsorbent capacity of a certain material is related to the material balance adsorption: the sorbate that disappears from solution must be in the adsorbent. There are a considerable number of expressions that describe adsorption isotherms and between them, Freundlich and Langmuir models were chosen and applied for describing the equilibrium data.

2.6.1. Freundlich Model

The Freundlich isotherm fits many soil adsorption systems and is represented by equation (2):

$$\Gamma_e = K_F \cdot C_{eq}^{1/n} \quad (2)$$

where, Γ_e is adsorption per unit area of adsorbent; K_F is the Freundlich constant indicating the relative adsorption capacity while $1/n$ is the index of the heterogeneity of the surface or the adsorption intensity and C_{eq} is the equilibrium concentration of adsorbate in solution.

2.6.2. Langmuir Model

The Langmuir isotherm is a well known model that indicates a decrease of the available surface sites as the adsorbent concentration increases. The Langmuir isotherm assumes monolayer adsorption:

$$\Gamma = \Gamma_{max} \frac{K_L C_e}{1 + K_L C_e} \quad (3)$$

where Γ is the amount of glyphosate adsorbed ($\mu\text{mol.m}^{-2}$), Γ_{max} is the maximum amount of glyphosate adsorbed on the surface ($\mu\text{mol.m}^{-2}$) at a fixed pH and temperature, K_L is the Langmuir adsorption constant (mM^{-1}) which is related to the free energy of the reaction and C_e is the equilibrium concentration of herbicide in the solution (mM). In other

words, Γ_{max} is the concentration of PMG surface saturation.

3. Results and Discussion

Soil characteristics, chemical analysis, mineralogical composition and quantitative analysis are presented in Table 1, 2 and 3 respectively. XRD of the three soil horizons are shown in Figure 3.

The experimental curves of PZNPC recorded for the BC horizon are illustrated in Figure 4. Similar behavior was found for all the horizons that showed PZNPC values in the range of 7.1 – 8.1 (Table 1) following the sequence: $\text{Ap} < \text{AB} < \text{BC}$. PZNPC value can be explained by the absence of clay minerals with a negative permanent charge, while the presence of 2:1 clays shift the PZNPC to lower pH values (Table 3).

The higher PZNPC value for the horizons corresponds to horizon BC that contains similar amount of quartz, lower amount of feldspars (andesine) and high amount of illite. PZNPC increase with andesine feldspar content and OM decrease. The determination coefficients of a linear fit were $R^2_{\text{andesine}} = 0.9971$ and $R^2_{\text{OM}} = 0.9189$. The analysis of the three parameters variations in a 3D plot presented a determination coefficient of $R^2 = 1.0000$ and a constant variance test of $p < 0.0001$.

To describe the adsorption behavior Freundlich and Langmuir models were applied to the equilibrium data.

The PMG adsorption isotherms of soils dispersions equilibrated at different pH values are shown in Figure 5. The Freundlich model parameters values (K_F and $1/n$) were calculated using equation 2 and are given in Table 4. The $1/n$ values vary between 0.1 and 1, which indicates that this model could be used for interpreting the data. The correlation between experimental and calculated curves had a p-level between 0.137 and 0.0035; the determination

Table 1. Characteristics of agriculture soils profile from Santiago del Estero/Argentina.

Horizon	pH (CaCl_2 1:2.5)	OM (g C.Kg ⁻¹)	P ($\mu\text{g.g}^{-1}$)	Sw (m ² .g ⁻¹)	Fe _{amorph} (mg.g ⁻¹)	Fe _{tot} (mg.g ⁻¹)	PZNPC (pH)
Ap	5.90	23.30	43.34	188	0.239	1.66	7.1
AB	5.75	17.10	6.67	259	0.158	1.91	7.4
BC	6.02	12.10	1.19	242	0.095	0.99	8.1

Table 2. Chemical Analysis of agriculture soils profile from Santiago del Estero, Argentina.

Horizon	Chemical Analysis (%)						
	SiO ₂	Al ₂ O ₃	Fe ₂ O ₃	CaO	MgO	Na ₂ O	K ₂ O
Ap	64.6	12.25	3.51	1.47	1.14	1.70	2.43
AB	63.0	13.65	4.10	1.34	1.40	1.50	2.53
BC	63.0	14.05	4.42	1.44	1.62	1.56	2.66

coefficients were between 0.7578 and 0.9953 for different pHs and horizons.

The Langmuir model was also applied to make an interpretation of PMG adsorption isotherms on soils dispersions equilibrated at different pH values. This is shown in Figure 5, where solid lines are calculated using this model (equation 3) and Γ_{\max} and K_L are given. The isotherm model parameters were obtained by a non-linear optimization using the Solver-Excel tool. The parameters values were obtained from the plot of the inverse of the surface coverage as a function of the inverse of the equilibrium concentration. Results of the adsorption and surface coverage calculations were normalized with S_w data and the various horizons were contrasted. The correlation between experimental and calculated curves had a p-level between 0.050 and 0.001; the determination coefficients (R^2) obtained were between 0.9300 and 0.9999; and were higher than those obtained using the Freundlich model. Thus, the

Langmuir model would better represent the adsorption process of PMG on the Santiago del Estero Province soil.

The dependence of the surface coverage with PMG concentration in the various horizons at constant pH = 5 is shown in Figure 6. Horizon Γ_{\max} sequence is Ap < AB < BC. This behavior is similar to those found for PZNPC.

The dependence of the surface coverage with pH in the various horizons is also shown in Figure 5. The adsorption capacity increases from pH 8 to 2. This pH effect was normally observed during the adsorption of anionic species. Consequently, PMG interaction with the surface occurs throughout the anionic chemical groups (carboxylate or phosphonate) and not through the amine group (pKa = 10.14) that is positively charged at the studied pH range (Figure 7).

The surface coverage decrease, $\Delta\Gamma_{\max}$, for horizon Ap is around 41 % for this pHs range (Table 5). This difference is lower for horizons BC, 27 %, and AB, 12 %.

The different composition and properties of the soils (Table 1), will affect sorption of the sorbate. This variability may also be linked to the organic matter content (Table 1) because of the chemical groups present on the OM (carboxyl, hydroxyl, amine, phenoxy, etc) that can coordinate the solid inorganic surface active sites or block the access of PMG to the inorganic surface. In consequence, OM presence can reduce PMG adsorbed on the solid surface. In fact, Γ_{\max} varies linearly with OM content with determination coefficients among $R^2_{pH=2} = 0.9548$ and $R^2_{pH=8} = 1.0000$. The affinities of the substances involved in the adsorption process are dependent on the identity and

Table 3. Mineralogical Composition of agriculture soils profile from Santiago del Estero, Argentina. Values in parenthesis represent estimated standard deviations.

Horizon	Mineralogical Composition (%)				
	Quartz	Sanidine Feldspar	Andesine Feldspar	Illite	Magnetite
Ap	45.2 (0.4)	9.6 (0.9)	24.7 (0.8)	18.6 (1.4)	1.3 (0.2)
AB	39.8 (0.5)	9.6 (0.8)	23.5 (0.7)	25.9 (1.5)	1.2 (0.3)
BC	46.2 (0.4)	9.3 (0.9)	19.9 (0.9)	24.7 (1.3)	1.2 (0.3)




Figure 3. XRD of the three soil horizons. Q: Quartz, Ar: Clay, F: Feldspar, Mt: Magnetite.




Figure 4. Potentiometric titration curves of the dispersions of the BC horizon at three ionic strengths ($I = \frac{1}{2} \sum c_i z_i^2$).

Table 4. Freundlich parameters ($\ln \mu\text{mol}^{1-1/n} \cdot \text{m}^{-2}$) for glyphosate adsorption on Santiago del Estero Province soils.

Horizon	Ap			AB			BC		
pH	$K_F \cdot 10^3$	$1/n$	R^2	$K_F \cdot 10^3$	$1/n$	R^2	$K_F \cdot 10^3$	$1/n$	R^2
2	7.3	0.46	0.9449	19.7	0.44	0.9745	5.3	0.64	0.9842
3	6.7	0.47	0.9953	18.4	0.45	0.9894	4.9	0.65	0.9759
4	5.9	0.47	0.9646	16.2	0.46	0.9654	4.7	0.63	0.9852
5	5.6	0.46	0.9729	16.2	0.45	0.9652	4.6	0.63	0.9892
6	5.0	0.48	0.7578	15.1	0.46	0.9862	4.3	0.62	0.9812
8	3.8	0.47	0.9453	14.3	0.45	0.9749	3.0	0.64	0.9893




Figure 6. Adsorption isotherm of PMG on horizon Ap, AB and BC at pH 5.




Figure 5. Adsorption isotherm of PMG on horizon Ap, AB and BC. Solid lines are calculated using Langmuir model with constants and maximum surface coverage detailed in Table 5. ●: pH 2, ■: pH 3, ◆: pH 4, ▲: pH 5, ▼: pH 6 and ★: pH 8.

Table 5. Langmuir parameters for PMG adsorption on soils of Santiago del Estero Province, Argentina

Horizon	Ap			AB			BC		
pH	Γ_{\max} , $\mu\text{mol.m}^{-2}$	K_L , L.mmol^{-1}	R^2	Γ_{\max} , $\mu\text{mol.m}^{-2}$	K_L , L.mmol^{-1}	R^2	Γ_{\max} , $\mu\text{mol.m}^{-2}$	K_L , L.mmol^{-1}	R^2
2	0.41	0.91	0.9572	0.81	0.93	0.9936	1.50	0.40	0.9331
3	0.40	0.87	0.9723	0.82	0.93	0.9833	1.49	0.42	0.9664
4	0.38	0.73	0.9874	0.78	0.92	0.9925	1.36	0.41	0.9955
5	0.34	0.67	0.9769	0.76	0.92	0.9932	1.27	0.40	0.9854
6	0.31	0.63	0.9633	0.74	0.91	0.9974	1.23	0.39	0.9858
8	0.24	0.61	0.9424	0.72	0.90	0.9926	1.10	0.28	0.9830




Figure 7. PMG acid-base equilibrium.

number of chemical groups present on the OM.

The highest adsorption capacity is obtained by horizon BC followed by horizon AB, and the lowest for horizon Ap. A similar sequence was obtained for PZNPC (Table 1), indicating that the horizon with higher positive surface charge presents higher PMG surface coverage.

The ratio of the Γ_{\max} of the horizons ($R_{H1/H2}$) was calculated as follow

$$R_{H1/H2} = \frac{\Gamma_{\max H1}}{\Gamma_{\max H2}} * 100 \quad (4)$$

where H1 and H2 denote two different horizons, $\Gamma_{\max H1}$ and $\Gamma_{\max H2}$ indicate the maximum coverage of H1 and H2 horizons, respectively. This ratio between the horizons BC and AB was $R_{BC/AB} = 46\%$, between horizons BC and Ap was $R_{BC/Ap} = 72\%$ and between horizon AB and Ap was $R_{AB/Ap} = 50\%$. These percentages are opposed to the phosphate content that follows the order of $Ap > AB > BC$. The high adsorption on deep horizons with regard to horizon Ap could be due to the competition of herbicide with phosphate groups for surface sites (Dion *et al.*, 2001; Gimsing *et al.*, 2007). Adsorption of glyphosate and phosphate in soil is similar to that which occurs on clay minerals (Dion *et al.*, 2001; Gimsing and dos-Santos-Afonso, 2005).

The highest adsorption constants correspond to horizon AB (Table 5). The changes in the adsorption affinity between horizon BC and AB reach $\Delta K_L = 46\%$ while horizon BC decreases 73 % in respect to horizon Ap.

The relatively high phosphate and low surface area (Table 1) of the Ap horizon, could be the cause for which it reaches its maximum at relatively low concentrations of PMG adsorption. The greater slope of the adsorption curves in the AB horizon indicate that PMG binds more strongly to the active sites of this horizon. Previous studies indicated that PMG bound iron oxides surfaces more strongly than clay minerals (dos-Santos-Afonso *et al.*, 2004). Thus, the

active site of PMG adsorption on the AB horizon could be the surface iron atoms and the higher adsorption in this horizon is directly related to higher iron content.

The adsorption on horizon BC does not reach maximum coverage in the experimental conditions. The adsorption isotherms with a low initial slope describe an adsorption process with characteristic adsorption constants of low energy interaction (Figure 5).

Note that in the working pH range an acid-base dissociation of the PMG molecule that contains amine, carboxylate, and phosphonate functional groups take place. The constant and the equilibrium reactions of acid-base dissociation of glyphosate (Barja and dos-Santos-Afonso, 1998) are shown in Figure 7, where I, II and III are the main species presents in the studied pH range. Previous studies suggested that the adsorption process occurs by surface complex formation via phosphonate group coordination to the mineral surface (Barja and dos-Santos-Afonso, 2005; Khoury *et al.*, 2010). Similar behavior should be expected for the adsorption of PMG on soils where coordination would occur on the surface of the minerals that compose them.

4. Conclusions

The major factor in PMG adsorption on soil samples is given by the pH, which could be due to the influence of this parameter on the PMG molecule and on the surface charge of the soil particles. PMG adsorption increase with acidity, and this increase correspond to the adsorption of a ligand with a negative net charge.

Sorption of glyphosate in soils is similar to the adsorption of the organic molecule on the soil components such as clay minerals, iron oxides and OM. For these soils with a low organic matter contents and/or similar amounts of clay in the various horizons, the adsorption would be determined by the content of phosphorus, iron oxide and the specific surface. Regarding the relative adsorption capacity of the soil, the adsorption process has a different behavior profile, where the deeper horizon (BC) has a higher capacity retention for this herbicide. The lower adsorption in the AB and Ap horizons could be influenced by the higher content of phosphorus. However, the strength of the interaction, as given by the Langmuir Model Constant K_L is larger on horizon AB and would be linked to the illite and iron oxide content that have a better distribution in AB.

It should be noted that the Langmuir adsorption model is the best fit to the adsorption experimental results in these soils, although the Freundlich model has a good fit for some pHs.

Given the adsorption extent found in this study, it is expected that pesticides will be retained in these soils. This strong interaction could prevent the pesticides movement into the groundwater. On the other hand, this retention rate could result in the release of the herbicide on the

environment due to displacement by runoff.

Acknowledgements

The authors acknowledge the Universidad de Buenos Aires, Secretaría de Ciencia y Técnica and MINCyT-ANPCyT- FONCyT for financial support. The authors are also grateful to Susana Conconi and Jorge Maggi from Centro de Tecnología de Recursos Minerales y Cerámica (CETMIC) for Rietveld analysis of soils.

References

- Barja, B., dos-Santos-Afonso, M., 1998, An ATR-FTIR study of glyphosate and its Fe(III) complex in aqueous solution: *Environmental Science and Technology*, 32, 3331-3335.
- Barja, B., dos-Santos-Afonso, M., 2005, Aminomethylphosphonic acid and Glyphosate adsorption onto goethite: a comparative study: *Environmental Science and Technology*, 39, 585-592.
- Barja, B., Herszage, J., dos-Santos-Afonso, M., 2001, Iron (III)-phosphonate complexes: *Polyhedron*, 20, 1821-1830.
- Damonte, M., Torres-Sánchez, R., dos-Santos-Afonso, M., 2007, Some aspects of the glyphosate adsorption on montmorillonite and its calcined form: *Applied Clay Science*, 1-3(36), 86-94.
- Dion, H., Harsh, J., Hill Jr., H., 2001, Competitive sorption between glyphosate and inorganic phosphate on clay minerals and low organic matter soils: *Journal of Radioanalytical and Nuclear Chemistry*, 249(2), 385-390.
- dos-Santos-Afonso, M., Barja, B., Pessagno, R., Tevez, H., 2004, Glyphosate adsorption on soils, in *Division of Environmental Chemistry*, New York, USA.: American Chemical Society.
- Gimsing, A., Borggaard, O., 2001, Effect of KCl and CaCl₂ as background electrolytes on the competitive adsorption of glyphosate and phosphate on goethite, *Clays Clay Minerals*, 49, 270-275.
- Gimsing, A., dos-Santos-Afonso, M., 2005, Glyphosate, in Nowack, B., VanBriesen, J. (ed.), *Biogeochemistry of Chelating Agents: ACS Symposium Series*, USA, 910, 263-277.
- Gimsing, A., Szilas, C., Borggaard, O., 2007, Sorption of glyphosate and phosphate by variable-charge tropical soils from Tanzania, *Geoderma*, 138, 127-132.
- Hendershot W.H., Lalonde, H., Duquette, M., 1993, Soil reaction and exchangeable acidity: In Carter M.R. (ed), *Soil sampling and methods of analysis*: Lewis Publishers, Boca Raton, 141-145.
- Holmgren, G., 1967, A rapid citrate-dithionite extractable iron procedure: *Soil Science Society of America Journal*, 31, 210-211.
- Jacobsen, C.S., van der Keur, P., Ivensen, B.V., Rosenberg, P., Barlebo, H.C., Torp, S., Vosgerau, H., Juhler, R.K., Ernstsén, V., Rasmussen, J., Brinch, U.C., Jac, O.H., 2008, Variation of MCPA, metribuzine, methyltriazine-amine and glyphosate degradation, sorption, mineralization and leaching in different soil horizons: *Environmental Pollution*, 3(156), 794-802.
- Khoury, G., Gehris, T., Tribe, L., Torres-Sánchez, R.M., dos-Santos-Afonso, M., 2010, Glyphosate adsorption on montmorillonite: An experimental and theoretical study of surface complexes: *Applied Clay Science*, 2(50), 167-175.
- Lorenz, G., Bonelli, C., Roldán, S., Araya, C., 2000, Soil quality changes due to land use in a Kastanozem-Phaeozemsoilscape of semiarid Chaco: *Mitteilungen der Deutschen Bodenkundlichen Gesellschaft*, 92, 169-172.
- McKeague, J.A., 1967, An evaluation of 0.1 M pyrophosphate and pyrophosphate-dithionite in comparison with oxalate as extractants of the accumulation products in podzols and some other soils: *Canadian Journal of Soil Science*, 47, 2, 95-99.
- McBride, M., Kung, K., 1989, Complexation of glyphosate and related ligands with iron (III): *Soil Science Society of America Journal*, 53, 1668-1673.
- McConnell, J., Hossner, L., 1985, pH-dependent adsorption isotherms of glyphosate: *Journal of Agricultural Food Chemistry*, 33, 1075-1078.
- Morillo, E., Undabeytia, T., Maqueda, C., Ramos, A., 2000, Glyphosate adsorption on soils of different characteristics. Influence of copper addition: *Chemosphere*, 40, 103-107.
- Nomura, N., Hilton, H., 1977, The adsorption and degradation of glyphosate in five. Hawaiian sugarcane soils: *Weed Research*, 17, 113-121.
- Nowack, B., Stone, A., 1999, Adsorption of phosphonates onto the goethite-water interface: *Journal of Colloid Interface Science*, 214, 23-30.
- Olsen, S., Cole, C., Watanabe, F., Dean, L., 1954, Estimation of available phosphorous in soils by extraction with sodium bicarbonate, United States of America, Department of Agriculture.
- Page, A., Miller, R., Keeney, D., 1982, Part 2: Chemical and microbiological properties, in *Methods of soil analysis*, Madison, Wisconsin, USA.
- Pérez-Carrera, A., Moscuza, C., Fernández-Cirelli, A., 2008, Efectos socioeconómicos y ambientales de la expansión agropecuaria. Estudio de caso: Santiago del Estero, Argentina. *Ecosistemas* 17(1), 5-15.
- Pessagno, R., dos-Santos-Afonso, M., Torres-Sánchez, R., 2005, N-(phosphonomethyl)glycine interactions with soil: The Journal of Argentine Chemical Society. *Anales de la Asociación Química Argentina*, 4-6(93), 97-108.
- Pessagno, R., Torres-Sánchez, R., dos-Santos-Afonso, M., 2008, Glyphosate behavior at soil and mineral-water interfaces: *Environmental Pollution*, 153, 53-59.
- Piccolo, A., Celano, G., Conte, P., 1996, Adsorption of glyphosate by humic substances: *Journal of Agricultural and Food Chemistry*, 44, 2442-2446.
- Rietveld, H.M., 1969, A profile refinement method for nuclear and magnetic structures: *Journal Applied Crystallography*, 2, 65-71.
- Schlichting, E., Blume, H., Stahr, K., 1995, *Soils Practical*, Wissenschafts-Verlag Blackwell, Berlin, Germany, 295 p.
- Sheals, J., Sjöberg, S., Persson, P., 2002, Adsorption of glyphosate on goethite: molecular characterization of surface complexes: *Environmental Science and Technology*, 36, 3090-3095.
- Sposito, G., 1984, *The surface chemistry of soils*: Oxford University Press, New York, 234 p.
- Tévez, H., dos-Santos-Afonso, M., Alvarez-Escalada, F.C., Serrano, M., Ruiz-Mostacero, N., 2008, Glifosato en fracciones de suelo de la provincia de Santiago del Estero, in *V Congreso Iberoamericano de Física y Química Ambiental*, Mar del Plata, Buenos Aires, Argentina.
- Torres-Bruchman, E., 1981, *Climatología general y agrícola de la Provincia de Santiago del Estero*, Ed. Universidad Nacional de Tucumán.
- Torres-Sánchez, R., Falasca, S., 1997, Specific surface area and surface charges of some argentinian soils: *Zeitschrift für Pflanzenernährung und Bodenkunde*, 160, 223-226.
- Veiga, F., Zapata, J., Marcos, M., Alvarez, E., 2001, Dynamics of glyphosate and aminomethylphosphonic acid in a forest soil in Galicia, north-west Spain: *Science of the Total Environment*, 271, 135-144.
- Zhu, Y., Zhang, F., Tong, C., Liu, W., 1999, Determination of glyphosate by ion chromatography: *Journal of Chromatography*, 850, 297-301.

Manuscript received: October 27, 2014

Corrected manuscript received: February 7, 2015

Manuscript accepted: February 10, 2015